



**HAL**  
open science

## **Molecular analysis of two local falciparum malaria outbreaks on the French Guiana coast confirms the msp1 B-K1/varD genotype association with severe malaria.**

Eric Legrand, Beatrice Volney, Anne Lavergne, Caroline Tournegros, Loïc Florent, Doris Accrombessi, Micheline Guillotte, Odile Mercereau-Puijalon, Philippe Esterre

### ► To cite this version:

Eric Legrand, Beatrice Volney, Anne Lavergne, Caroline Tournegros, Loïc Florent, et al.. Molecular analysis of two local falciparum malaria outbreaks on the French Guiana coast confirms the msp1 B-K1/varD genotype association with severe malaria.. *Malaria Journal*, 2005, 4, pp.26. 10.1186/1475-2875-4-26 . pasteur-00590991

**HAL Id: pasteur-00590991**

**<https://riip.hal.science/pasteur-00590991>**

Submitted on 30 Jun 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Research

Open Access

## Molecular analysis of two local falciparum malaria outbreaks on the French Guiana coast confirms the *msp1* B-K1/*varD* genotype association with severe malaria

Eric Legrand\*<sup>1</sup>, Beatrice Volney<sup>1</sup>, Anne Lavergne<sup>1</sup>, Caroline Tournegros<sup>1</sup>, Loïc Florent<sup>1</sup>, Doris Accrombessi<sup>1</sup>, Micheline Guillotte<sup>2</sup>, Odile Mercereau-Puijalon<sup>2</sup> and Philippe Esterre<sup>1</sup>

Address: <sup>1</sup>Centre National de Référence sur la Chimiorésistance du Paludisme dans la région Antilles – Guyane, Institut Pasteur de la Guyane, BP 6010, F-97306 Cayenne-Cedex, France and <sup>2</sup>Unité d'Immunologie Moléculaire des Parasites, Unité de Recherche Associée 2581 du Centre National de la Recherche Scientifique, Institut Pasteur, Paris, France

Email: Eric Legrand\* - [elegrand@pasteur-cayenne.fr](mailto:elegrand@pasteur-cayenne.fr); Beatrice Volney - [bvolney@pasteur-cayenne.fr](mailto:bvolney@pasteur-cayenne.fr); Anne Lavergne - [alavergne@pasteur-cayenne.fr](mailto:alavergne@pasteur-cayenne.fr); Caroline Tournegros - [cnrcp@pasteur-cayenne.fr](mailto:cnrcp@pasteur-cayenne.fr); Loïc Florent - [cnrcp@pasteur-cayenne.fr](mailto:cnrcp@pasteur-cayenne.fr); Doris Accrombessi - [cnrcp@pasteur-cayenne.fr](mailto:cnrcp@pasteur-cayenne.fr); Micheline Guillotte - [mguillot@pasteur.fr](mailto:mguillot@pasteur.fr); Odile Mercereau-Puijalon - [omp@pasteur.fr](mailto:omp@pasteur.fr); Philippe Esterre - [pesterre@pasteur-cayenne.fr](mailto:pesterre@pasteur-cayenne.fr)

\* Corresponding author

Published: 21 June 2005

Received: 30 March 2005

*Malaria Journal* 2005, 4:26 doi:10.1186/1475-2875-4-26

Accepted: 21 June 2005

This article is available from: <http://www.malariajournal.com/content/4/1/26>

© 2005 Legrand et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** *Plasmodium falciparum* outbreaks can occur in the coastal area of French Guiana, where the population is essentially non-immune. Two sporadic outbreaks were observed, including one with severe malaria cases. To characterize these outbreaks and verify previous observations of specific genotype characteristics in severe malaria in this area, all cases from each outbreak were studied.

**Methods:** *P. falciparum* genotypes for six genetic loci were determined by PCR amplification from peripheral blood parasites. The *msp1*/block2 and *msp2* genotypes were determined by DNA sequencing. Microsatellite and *varD* genotyping was based on size polymorphism and locus-specific amplification.

**Results:** The outbreak including severe malaria cases was associated with a single genotype. The other mild malaria outbreak was due to at least five distinct genotypes.

**Conclusion:** Two distinct types of outbreak occurred despite systematic and sustained deployment of malaria control measures, indicating a need for reinforced vigilance. The *varD*/B-K1 *msp1* linkage and its association with severe malaria in this area was confirmed.

### Background

The annual malaria incidence in French Guiana has increased ten-fold during the last 30 years, reaching nowadays approximately 3%, with 60% due to *Plasmodium falciparum* and 40% due to *Plasmodium vivax*. French Guiana

is an area with multidrug resistant *P. falciparum* malaria. Transmission occurs in isolated foci located in settlement pockets within the Amazonian forest and along the rivers. The main malaria-endemic areas are located along the Maroni and Oyapock rivers, which serve as natural

frontiers with Surinam and Brazil, respectively [1]. As a result of three-monthly insecticide spraying campaigns, there are virtually no malaria cases along the coast, where 80% of the population resides. However, outbreaks, which usually are of short duration and affect a small number of patients, are occasionally reported in the coastal area.

The *P. falciparum* parasite population of French Guiana presents a remarkably low degree of polymorphism, with a clonal type structure [2]. The parasite diversity is so restricted that the circulation of specific genotypes could be followed within the area, making it possible to investigate the clinical impact of specific parasite genotypes. A comparative analysis of parasite genetic characteristics in isolates collected from mild and severe malaria patients has highlighted a significant linkage disequilibrium between a particular *msp1*/block2 allele that was called B-K1 and a particular *var* gene called *varD* in isolates from patients with severe *P. falciparum* malaria [3].

To verify this association and better understand the possible cause of local outbreaks in the coastal area located away from the main endemic sites, two *P. falciparum* outbreaks were studied, one with five cases, including two cases of severe malaria and one death, the other with nine mild malaria cases. The isolates from each case were characterized for both outbreaks using a six-locus genotyping approach, including *varD*, *msp1*/block2, *msp2* [2-7], as well as three microsatellite loci [8].

## Methods

### Parasite isolates and patient information

The outbreaks occurred in Macouria and Matoury, both located on the French Guiana coast, an area with unstable, low transmission, away from the main endemic areas along the Maroni and the Oyapock (Fig. 1). The international airport close to Cayenne, the main city, is in Matoury. The isolates were collected from patients with *P. falciparum* malaria in health centres and sent to the laboratory for routine drug susceptibility phenotyping and genotyping analysis. The policy of the malaria control services is to get an *in vitro* susceptibility profile for each diagnosed *P. falciparum* malaria case. To this end, a 5 mL blood sample was systematically collected by venepuncture in EDTA tubes and maintained at 4°C during shipment to the laboratory. The samples were immediately processed for *in vitro* susceptibility assays and a 0.5 mL aliquot was frozen at -20°C for subsequent DNA preparation. The five Macouria isolates were collected in June and July 2001 from three mild (E57, E64 and E72) and two severe cases (E61, a fatal case, and E62). Patient E57 had visited the endemic Macapa region in Brazil one month before the malaria attack. The other four Macouria patients did not report having left the area. The nine

Matoury isolates, collected from February to June 2002 originated from mild malaria cases, none of whom reported a recent visit to an endemic area.

### DNA extraction and genotyping

DNA was extracted as described [4]. Genotyping was performed by PCR using the primers listed in Figure 2. For *msp1* block2, *msp2* and *varD*, PCR was performed in a 50 µL final volume containing the DNA template, 1.5 mM MgCl<sub>2</sub>, 2 µM each primer, 200 µM dNTP, and 1.5 units of AmpliTaq Gold polymerase (Applied Biosystems, Foster City, CA, USA). The PCR conditions for *msp1* block2 and *msp2* were: one cycle at 95°C for 5 min, followed by 40 cycles at 94°C for 1 min, 56°C for 2 min, 72°C for 2 min, and final extension at 72°C for 10 min. The *varD* PCR conditions were: one cycle at 94°C for 5 min, followed by 35 cycles at 94°C for 10 sec, 61°C for 1 min 30 sec, 72°C for 2 min, and one cycle at 72°C for 10 min. The PCR products were analysed by agarose gel electrophoresis and directly sequenced (Qbiogen, Every, France). Sequences were aligned using the Clustal W program, followed by manual analysis using the ED editor of the MUST package [9].

Microsatellite typing was done using the C1M4, C3M27 and C4M69 loci, located on chromosomes 1, 3 and 4, respectively. The locus-specific primers described by Su *et al.* [8] were used (see Figure 2). Amplification was done in 15 µL reaction volume containing the DNA template, 1.5 mM MgCl<sub>2</sub>, 2 µM each primer, 200 µM each dNTP, and 0.5 Units *Taq* polymerase (Promega, Madison, WI, USA). The samples were subjected to 1 cycle at 94°C for 7 min, followed by 35 cycles at 94°C for 30 sec, 52°C for 30 sec, 47°C for 30 sec, 72°C for 30 sec, and a final extension step at 72°C for 15 min. The PCR products were analysed by 3% Metaphor agarose gel (FMC Bioproduct, Rockland, Maine, USA) electrophoresis and stained with ethidium bromide. Fragment size was calculated using the Taxotron software (P.A.D. Grimont, Institut Pasteur, Paris).

### Results and discussion

PCR amplification of the *msp1*/block2 and *msp2* *P. falciparum* loci generated a single band for all isolates (Fig. 3). Each PCR band was sequenced. This showed three distinct alleles for each locus. All Macouria samples carried the same 464-bp *msp1* block2 allele, which belonged to the K1-type allelic family, corresponding to the allele called B-K1 in a previous study [3]. A smaller (431 bp) K1-type allele, previously called A-K1 [3], and a 398 pb RO33-type allele were observed in six and three Matoury isolates, respectively. All three RO33-type sequences were identical to the RO33 allele (accession number Y00087). Alignment of the B-K1 and A-K1 deduced protein sequence with sequences available in the databases identified several alleles observed in other regions of the world showing



**Figure 1**  
 Map of French Guiana. Triangles indicate the localization of the severe malaria cases with K1-B/var D described by Arley et al. [3]. The towns of Macouria and Matoury are indicated.

up to 99% and 97% identity, respectively. However, no isolate in the database had precisely similar tri-peptide repeats as B-K1 or A-K1. This is in line with recent evidence indicating that *msp1*/block 2 (and *msp2*) repeats evolve at a faster rate than SNPs in the non-repeated domains [10]. Numerous isolates showed in addition sin-

gle nucleotide polymorphisms in the non-repeated sequences (Fig. 4).

All three *msp2* alleles belonged to the 3D7-type family and were arbitrarily called 3D7a (591 bp), b (561 bp) and c (603 bp). The 3D7a allele was observed in all five

locus	forward primer	reverse primer
<b>MSP1 block2</b> conserved A +B K1-type specific MAD20-type specific RO33-type specific	AAGCTTTAGAAGATGCGATTGAC GAA ATTACTACA AAAGGTGCA AGTG GAACAAGTCGAACAGCTGTTA GCA AATACTCAAGTTGTTGCA AAGC	ATTCATTAATTTCTTCATATCCATC AGATGAAGTATTTGAACGAGGTTAAAGTG TGAATTATCTGAAGGATTTGTACGCTTGA AGGATTTGCAGCACCTGGAGATCT
<b>MSP2</b> conserved 2+3 3D7-type specific FC27-type specific	AACGAATTCATA AACAATGCTTATAATATGAGT GCA GAA AGTAAGCCTTCTACTGGTGCT GCAAATGAAGTTCTAATACTAATAG	GATGAATTCTAGAACCATGCATATGTCCATGT GATTTGTTTCGGCATTATTATGA GCTTTGGGTCTTCTCAGTTGATTC
<b>VarD</b> VarD 5' + 3'	AGTTCACCTGGTTTCCCGC	CCCTGAAGATTTTAAGCGTC
<b>microsatellites</b> C1M4 C3M27 C4M69	ATATCCTACAACGGTAAGCA AGTATCATATTTGGTTAGATC GAAATGGAGATAAACTATTAC	GGCACATAAATAATACATAC TTTGGTTAACAAATTCCTAC AATTACACAACAGATGTGAA

**Figure 2**  
Sequence of primers.

Isolate	Clinical attack	Outbreak	Date access	Parasitaemia Percent inf RBC	<i>msp1</i> <i>block2</i> allele <sup>1</sup>	<i>msp2</i> allele <sup>2</sup>	C1M4 bp	C3M27 bp	C4M69 bp	<i>varD</i> amplified	<i>genotype</i>
E57*	mild	Macouria	22/05/2001	0.5	B-K1	3D7a	140	150	520	Yes	Macouria
E61 <sup>+</sup>	severe	Macouria	03/06/2001	0.35	B-K1	3D7a	140	150	520	Yes	Macouria
E62	severe	Macouria	04/06/2001	1	B-K1	3D7a	140	150	520	Yes	Macouria
E64	mild	Macouria	11/06/2001	0.05	B-K1	3D7a	140	150	520	Yes	Macouria
E72	mild	Macouria	28/06/2001	4	B-K1	3D7a	140	150	520	Yes	Macouria
F33	mild	Matoury	14/02/2002	1.5	A-K1	3D7a	140	150	<u>430</u>	Yes	Matoury 1
F40	mild	Matoury	17/02/2002	8	A-K1	3D7a	140	150	<u>430</u>	No	Matoury 2
F52	mild	Matoury	19/02/2002	20	A-K1	3D7a	140	150	<u>430</u>	Yes	Matoury 1
F53	mild	Matoury	18/02/2002	6.5	A-K1	3D7a	140	150	<u>430</u>	No	Matoury 2
F79	mild	Matoury	29/03/2002	0.1	RO33	3D7c	nil	130	360	No	Matoury 4
F89	mild	Matoury	03/04/2002	1	RO33	3D7c	nil	130	360	No	Matoury 4
F96	mild	Matoury	17/04/2002	3	A-K1	3D7b	140	150	<u>430</u>	No	Matoury 3
F105	mild	Matoury	18/04/2002	3	A-K1	3D7a	140	150	<u>430</u>	No	Matoury 2
F113	mild	Matoury	06/05/2002	0.05	RO33	3D7c	nil	130	<u>nil</u>	No	Matoury 5

\* The patient declared having traveled to Macapa (Brazil) one month before the malaria attack.

<sup>+</sup> Fatal case

<sup>1</sup> Sequence shown in Figure 2

<sup>2</sup> Sequence shown in Figure 3

**Figure 3**  
*msp1*//*block2*, *msp2*, microsatellite and *varD* *P. falciparum* genotypes observed in the Macouria and Matoury outbreaks.

Macouria isolates and in five of nine Matoury isolates. The 3D7b and 3D7c allele were observed in one and three Matoury isolates, respectively (Fig. 3). Alignment of the deduced protein sequence showed that allele 3D7c clus-

tered with a specific group of isolates showing typical particularities in the repeat region in particular the NPPA tetrapeptide interspersed within G-, A-, S- rich repeats, mainly GAGASG (Fig. 5). However, the 3D7c allele was

K1-A	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGP	SGT	SGT	SGT	SGP	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---		
K1-B	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGP	SGT	SGP	SGT	SGP	SGT	SGP	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---		
AF061134	LEDAVLGYSLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGP	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	---		
AF061126	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGT	SAQ	SGT	SGT	SGT	SGT	SGT	SGP	SGT	---	---	---	---	---	---	---		
AF061123	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	---		
AF061129	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGP	SGT	SGP	SGT	SGP	SGT	SGP	SGT	SGP	SGP	SGT	---	---	---	---	---	---	---	---		
AF061130	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGP	SGT	SGP	SGT	SGP	SGT	SGP	SGT	SGP	SGP	SGT	---	---	---	---	---	---	---	---		
M77733	LEDAVLGYSLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---		
AF061127	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---		
AF061132	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---		
AB116596	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGP	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---		
M77732	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGP	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---		
AF061133	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGP	SGT	SGP	SGT	SGP	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---		
AF061128	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SAQ	SGT	SGT	SAQ	SGT	SGT
AF061131	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGP	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
AF061124	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGP	SGP	SGT	---	---	---	---	---	---	---	---	
RO71	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGP	SGP	SGT	---	---	---	---	---	---	---	---	---	
AF061121	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	
FCB1	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGT	SAQ	SGT	SGT	SAQ	SGT	SGT	SGT	SAQ	SGT	SGT	SAQ	SGT	SGT	SGT	SGT	SGT	---
M77729	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGT	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	
K1	LEDAVLGYSLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	
AF061135	LEDAVLGYSLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	
Palo Alto	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGT	SAQ	SGT	SGT	SAQ	SGT	SGT	SAQ	SGT	SGT	SAQ	SGT	SGT	SGT	SGT	SGT	---	
M77734	LEDAVLGYGLFHKEKMI LNNEEII TTKGA	SAQ	SGT	SGT	SGT	SGT	SGT	SGT	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	---	---	
NF7	LEDAVLGYSLFGQEKRMV LNNEEII TTKGA	SAQ	SGA	SAQ	SGA	SAQ	SGA	SAQ	SGA	SGT	SGP	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	
AF061125	LEDAVLGYSLFGQEKRMV LNNEEII TTKGA	SAQ	SGA	SAQ	SGA	SAQ	SGA	SAQ	SGA	SGP	SGP	SGT	---	---	---	---	---	---	---	---	---	---	---	---	
3D7	LEDAVLGYSLFGQEKRMV LNNEEII TTKGA	SAQ	SGA	SAQ	SGA	SAQ	SGA	SAQ	SGA	SAQ	SGA	SAQ	SGT	SGP	SGP	SGT	---	---	---	---	---	---	---	---	

K1-A	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLLTIKELKYPQLFDTLNHMLTLC DN IHGFKYLIDGYEEI
K1-B	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
AF061134	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLLTIKELKYPQLFDTLN HMLTLC DN IHGFKYLIDGYEEI
AF061126	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
AF061123	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
AF061129	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
AF061130	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
M77733	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLLTIKELKYPQLFDTLN HMLTLC DN IHGFKYLIDGYEEI
AF061127	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
AF061132	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLLTIKELKYPQLFDTLN HMLTLC DN IHGFKYLIDGYEEI
AB116596	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
M77732	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
AF061133	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLLTIKELKYPQLFDTLN HMLTLC DN IHGFKYLIDGYEEI
AF061128	SGT	SGT	SGT	SGT	SPSSRSNTLPRS ILHLGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
AF061131	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLLTIKELKYPQLFDTLN HMLTLC DN IHGFKYLIDGYEEI
AF061124	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	PPADAS	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
RO71	----	----	----	----	SPSSRSNTLPRSN TSSG I PPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
AF061121	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
FCB1	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
M77729	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
K1	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLLTIKELKYPQLFDTLN HMLTLC DN IHGFKYLIDGYEEI
AF061135	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLLTIKELKYPQLFDTLN HMLTLC DN IHGFKYLIDGYEEI
Palo Alto	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
M77734	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
NF7	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
AF061125	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI
3D7	----	----	----	----	SPSSRSNTLPRSN TSSGASPPADAS	-----	DSDAKSYADLKHVRVRYLFTIKELKYPELFDLTN HMLTLC DN IHGFKYLIDGYEEI

**Figure 4**  
 Deduced amino acid sequence of the French Guiana A-K1 and B-K1 *msp1*/block2 alleles and alignment with alleles described in other areas. Single letter codes are used for amino acids. The name of the reference laboratory strains are indicated. The accession number for K1, FCB1, Palo Alto, NF7 and RO71 were X03371, AF286876, M37213, M19144, X61930. For 3D7, the sequence of the genome sequence (locus PFI1475w) is shown. For field isolates the Genbank accession number was used. The reference sequence used was K1. Deletions are indicated by a dash and mutant residues are in red, except for the 3D7-types which are in blue.

peculiar within this group in having the largest number of both types of repeats NNPA and presenting GGSA repeats as well. The 3D7a and 3D7b also presented each a specific arrangement of repeats as well as SNPs in the non repeated regions.

To further explore the genetic make-up of the *P. falciparum* isolates, three microsatellite loci were studied. A single 140-bp C1M4 product was observed for all isolates, except F79, F89 and F113, where no band was visualized (Fig. 3). A 150-bp C3M27 product was amplified from all Macou-

ria isolates and from the F33, F40, F52, F53, F96 and F105 Matoury isolates. A distinct 130-bp C3M27 allele was detected in isolates F79, F89 and F113. For the C4M69 locus, a single 520-bp band was obtained for all Macouria isolates, a 430-bp band for Matoury isolates F33, F40, F52, F53, F96 and F105 and a 360-bp fragment for the Matoury isolates F79 and F89. No product was obtained for this locus from isolate F113 (Fig. 3). This data indicated that for all five single copy loci investigated, all isolates generated a single, monomorphic PCR fragment. We can thus reasonably consider them as being monoclonal



**Figure 5**  
 Deduced amino acid sequence of 3D7-a, -b and -c *msp2* alleles and alignment with alleles described in other areas. Single letter codes are used for amino acids. The name of the reference laboratory strains are indicated. The 3D7 sequence (locus PFBO300c) was used as reference sequence. For field isolates the Genbank accession number was used. The Genbank accession number for Palo Alto was M61122. Deletions are indicated by a dash and mutant residues as compared to 3D7 are in red. The various repeat sequence types are underline or color-coded.

infections. This confirms previous findings in this area, where a very high percentage of single infections was observed [2,3], and is in line with reports from other areas of South America [5,6].

French Guiana is a hypoendemic area, where the population has limited, if any, immunity. There are between 6,000–7,000 mild malaria cases and only between four and 20 severe malaria cases each year [1,3]. Previous studies have highlighted specific genotype characteristics in *P. falciparum* isolates from severe cases in this setting, with the B-K1 *msp1* allele in strong linkage disequilibrium with a particular *var* gene, called *varD* [3]. The *var* genes code for the variant adhesins that mediate the cytoadhesion of

parasitized erythrocytes to specific host receptors that undoubtedly contribute to severe malaria. As highlighted in Figure 3, all five Macouria isolates, including the two severe cases, had the same 5-loci genotype, suggesting that a single virulent isolate was present. These isolates were also tested for the presence of the *varD* gene. A typical *varD* product was amplified from all five Macouria isolates. Only two Matoury isolates (isolates F33 and F52) carried the *varD* locus, none of which harboured the B-K1 *msp1* allele (Fig. 3). Overall, the same genotype was observed in all Macouria isolates while there were at least five distinct genotypes in Matoury. The Matoury 1 and 2 genotypes differed at the *varD* locus. The Matoury 2 and Matoury 3 genotypes had a different *msp2* allele. Furthermore, isolates

F40 and F96 had a different drug susceptibility profile (data not shown). The Matoury 4 and Matoury 5 genotypes differed by the C4M69 microsatellite locus.

These data confirm the *varD*/B-K1 *msp1* linkage and its association with severe malaria in this area. It is important to note that this association was previously detected in isolates collected in 1994-6, more than six years before the Macouria epidemic. Such a stability over time in a species with a high recombination rate [8] is consistent with previous data pointing to a high selfing rate in this area [3]. The severe malaria cases studied here originated from a geographic area quite distinct from the previous cases reported where the *varD*/B-K1 association was observed, as illustrated in Figure 1. As observed previously [3], the *varD*/B-K1 *msp1* association was not strictly specific for severe malaria, as the same genotype was also observed in the non-severe Macouria cases. Whether the patients with mild malaria received an earlier treatment compared to those with severe malaria or were less susceptible due to their genetic or immune make up is unclear. The occurrence of two severe cases at one day interval in patients infected with this strain suggests a particular inherent virulence. Whether this is due to the particular *msp1* and/or *varD* allele present is unknown. It may reflect physical association with another locus implicated in virulence. It is worth noting that isolate F52, which harboured *varD* together with the A-K1 *msp1* allele, was collected from a patient with very high peripheral parasitaemia (20%), also considered a sign of severity [11]. *VarD* is one of approximately 60 members of the *P. falciparum* *var* genomic repertoire [12]. The presence of a particular *var* gene is not synonymous with its expression. Expression of *varD* was demonstrated in a patient with fatal *P. falciparum* malaria in a previous study [3], but could not be studied here. The data indicate that future investigations on *varD* expression in severe and non-severe malaria are warranted. Work is in progress to characterize the full *varD* gene sequence.

The different genetic profiles of the isolates involved in the two outbreaks reveal distinct onset and dynamics scenarios. Genotyping strongly suggests that the Macouria outbreak was due to one single parasite strain, but the origin of this strain is uncertain. Transmission from E57, the first registered case, to the other cases is unlikely, due to the short delay between the clinical attack and the attack experienced by E61, E62 and E64 (12, 13 and 20 days later, respectively). E67 may possibly have subsequently transmitted the strain to E72, who experienced a malaria attack 37 days later. The diversity of the Matoury isolates indicates that the outbreak was certainly caused by several distinct strains. Matoury accommodates the largest airport in the country and consequently may serve as an occasional transmission focus for parasites originating in

neighbouring endemic malaria areas. The possible cause and mode of transmission in this city at that time have not been identified.

## Conclusion

The results point to two distinct types of outbreak in a region where malaria control measures are systematically deployed and sustained. Reinforced vigilance and rapid case notification are needed to ensure rapid deployment of vector control and personal protection measures to prevent such sporadic epidemics. The Macouria outbreak provoked two severe cases, including one death, a rare event in this area, where health facilities are well-equipped and treatment policy is regularly updated. Parasite genotyping confirmed the association of the B-K1 *msp1*/*varD* genotype with severe malaria, reinforcing the notion that some *P. falciparum* strains might cause more severe infections than others.

## Authors' contributions

A. Lavergne and E. Legrand did the sequences analysis. B. Volney, C. Tournegros, D. Accombrassi and L. Florent did the laboratory experiments. M. Guillotte and O. Mercereau-Puijalon adapted the microsatellite typing to field isolates. O. Mercereau-Puijalon did the database search. All authors participated in the analysis and interpretation of data. E. Legrand and O. Mercereau-Puijalon wrote the manuscript.

## Acknowledgements

This project was supported by the French Ministry of Health (InVS agency, Paris) and the French Academy of Sciences (Prix Louis D., 2000). The authors thank Prof. B. Carne and colleagues from the Parasitology Unit of the Central Hospital, Cayenne and Drs. P. Duval and B. Maubert, from the Biomedical Unit of the Institut Pasteur de Guyane, Cayenne for providing the blood samples.

## References

1. Esterre P, Cordoliani G, Germanetto P, Robin Y: **Epidémiologie du paludisme en Guyane française.** *Bull Soc Pathol Exot* 1990, **83**:193-205.
2. Arieu F, Chalvet W, Hommel D, Peneau C, Hulin A, Mercereau-Puijalon O, Duchemin J-B, Sarthou J-L, Reynes J-M, Fandeur T: **Plasmodium falciparum parasites in French Guiana: limited genetic diversity and high selfing rate.** *Am J Trop Med Hyg* 1999, **61**:978-985.
3. Arieu F, Hommel D, LeScanf C, Duchemin JB, Peneau C, Hulin A, Sarthou JL, Reynes JM, Fandeur T, Mercereau-Puijalon O: **Association of severe malaria with a specific Plasmodium falciparum genotype in French Guiana.** *J Infect Dis* 2001, **184**:237-241.
4. Contamin H, Fandeur T, Bonnefoy S, Skouri F, Ntoumi F, Mercereau-Puijalon O: **PCR typing of field isolates of Plasmodium falciparum.** *J Clin Microbiol* 1995, **33**:944-951.
5. Sallenave-Sales S, Ferreira-da-Cruz MF, Faria CP, Cerruti C Jr, Daniel-Ribeiro CT, Zalis MG: **Plasmodium falciparum : limited genetic diversity of MSP-2 in isolates circulating in Brazilian endemic areas.** *Exp Parasitol* 2003, **103**:127-135.
6. Tami A, Grundmann H, Sutherland C, McBride JS, Cavanagh DR, Campos E, Snounou G, Barnabe C, Tibayrenc M, Warhurst DC: **Restricted genetic and antigenic diversity of Plasmodium falciparum under mesoendemic transmission in the Venezuelan Amazon.** *Parasitology* 2002, **124**:569-581.



7. Jelinek T, Kilian AH, Westermeier A, Proll S, Kabagambe G, Nothdurft HD, von Sonnenburg F, Loscher T: **Population structure of recrudescence *Plasmodium falciparum* isolates from western Uganda.** *Trop Med Int Health* 1999, **4**:476-480.
8. Su X, Ferdig MT, Huang Y, Huynh CQ, Liu A, You J, Wootton JC, Welles TE: **A genetic map and recombination parameters of the human malaria parasite *Plasmodium falciparum*.** *Science* 1999, **286**:1351-1353.
9. Philippe H: **MUST, a computer package of management utilities for sequence and tree.** *Nucleic Acids Res* 1993, **21**:5264-5272.
10. Tanabe K, Sakihama N, Kanejo A: **Stable SNPs in malaria antigen genes in isolated populations.** *Science* 2004, **303**:493.
11. World Health Organization: **Severe falciparum malaria.** *Trans R Soc Trop Med Hyg* 2000, **94(Suppl 1)**:1-90.
12. Gardner MJ, Hall N, Fung E, White O, Berriman M, Hyman RW, Carlton JM, Pain A, Nelson KE, Bowman S, Paulsen IT, James K, Eisen JA, Rutherford K, Salzberg SL, Craig A, Kyes S, Chan M-S, Nene V, Shalom SJ, Suh B, Peterson J, Angiuoli S, Pertea M, Allen J, Selengut J, Haft D, Mather MW, Vaidya AB, Martin DMA, Fairlamb AH, Fraunholz MJ, Roos DS, Ralph SA, McFadden GI, Cummings LM, Subramanian GM, Mungall C, Venter JC, Carucci DJ, Hoffman SL, Newbold C, Davis RW, Fraser CM, Borel B: **Genome sequence of the human malaria parasite *Plasmodium falciparum*.** *Nature* 2002, **419**:498-511.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

