



High Prevalence of Shared International Type 53 among Mycobacterium tuberculosis Complex Strains in Retreated Patients from Côte d'Ivoire.

Timothée Ouassa, Emanuele Borroni, Guillaume Yao Loukou, Hortense Faye-Kette, Jacquemin Kouakou, Hervé Menan, Daniela Maria Cirillo,
Timothée Ouassa

► To cite this version:

Timothée Ouassa, Emanuele Borroni, Guillaume Yao Loukou, Hortense Faye-Kette, Jacquemin Kouakou, et al.. High Prevalence of Shared International Type 53 among Mycobacterium tuberculosis Complex Strains in Retreated Patients from Côte d'Ivoire.. PLoS ONE, 2012, 7 (9), pp.e45363. 10.1371/journal.pone.0045363 . pasteur-00738645

HAL Id: pasteur-00738645

<https://riip.hal.science/pasteur-00738645>

Submitted on 4 Oct 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

High Prevalence of Shared International Type 53 among Mycobacterium tuberculosis Complex Strains in Retreated Patients from Côte d'Ivoire.

[Ouassa T](#), [Borroni E](#), [Loukou GY](#), Faye-Kette H, [Kouakou J](#), [Menan H](#), [Cirillo DM](#).

Source

Department of Bacteriology and Virology, Faculty of Pharmacy, University of Cocody, Abidjan, Côte d'Ivoire ; Centre for Diagnosis and Research on AIDS and opportunistic infections, Teaching Hospital of Treichville, Abidjan, Côte d'Ivoire.

Abstract

BACKGROUND:

Genotyping methods are useful tools to provide information on tuberculosis epidemic. They can allow a better response from health authorities and the implementation of measures for tuberculosis control. This study aimed to identify the main lineages and clades of Mycobacterium tuberculosis complex strains circulating in Côte d'Ivoire.

METHODS/MAIN FINDINGS:

Strains isolated from sputum samples of patients ongoing retreatment from all the country were characterized by spoligotyping and by MIRU-VNTR. Profiles obtained by spoligotyping were first compared to the SITVIT/SpolDB4 database for family assignment. Of 194 strains analysed, 146 (75.3%) belonged to the T lineage. The most predominant spoligotype was the shared international type 53 with 135 strains (69.6%). In contrast with neighbouring countries, LAM (11 strains, 5.7%) and H (9 strains 4.6%) lineages were slightly represented. Only 3 Beijing strains (1.5%) and 4 strains of Mycobacterium africanum (2%) were found. Analysis of the results obtained with MIRU-VNTR revealed also a high level of clustering.

CONCLUSION/SIGNIFICANCE:

The population of Mycobacterium tuberculosis complex strains among retreatment cases in Côte d'Ivoire exhibits a low diversity, allowing to assume recent transmission and locally based infection.