



The jarisch-herxheimer reaction in leptospirosis: a systematic review.

Gilles Guerrier, Eric d'Ortenzio

► To cite this version:

Gilles Guerrier, Eric d'Ortenzio. The jarisch-herxheimer reaction in leptospirosis: a systematic review.. PLoS ONE, 2013, 8 (3), pp.e59266. 10.1371/journal.pone.0059266 . pasteur-00809318

HAL Id: pasteur-00809318

<https://riip.hal.science/pasteur-00809318>

Submitted on 8 Apr 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Jarisch-Herxheimer Reaction in Leptospirosis: A Systematic Review

Gilles Guerrier^{1*}, Eric D'Ortenzio²

1 Département d'Anesthésie-Réanimation, Centre Hospitalier Territorial, Noumea, Nouvelle Calédonie, **2** Unité d'Epidémiologie des Maladies Infectieuses, Institut Pasteur de Nouvelle-Calédonie, Réseau International des Instituts Pasteur, Noumea, Nouvelle Calédonie

Abstract

Background: Leptospirosis is an endemo-epidemic zoonotic disease associated with potentially fatal renal, cardiovascular or pulmonary failure. Recommended treatment includes antibiotics, which may induce a Jarisch-Herxheimer reaction (JHR). Since little information on the importance of this adverse event is available, we performed this review to quantify frequency and impact of JHR in leptospirosis management.

Methodology/Principal Findings: This review systematically summarizes the literature on the JHR in leptospirosis. To approach the broader aspects of the subject, articles considering the treatment of leptospirosis, national leptospirosis guidelines and textbook and technical reports of the World Health Organisation were reviewed. Publications describing JHR in leptospirosis are very limited and consist mainly of single case reports and small case series. A single randomized control trial specifically assessed the JHR occurrence, but it has never been systematically investigated in large trials. Not all guidelines and not all literature on leptospirosis mention this reaction which can be fatal.

Conclusions/Significance: Although generally assumed to be a rare event, the true prevalence of JHR in leptospirosis is unknown and the awareness of this event is insufficient. All leptospirosis guidelines and local leptospirosis protocols should stress on systematic monitoring for clinical status early after antibiotic administration. Large well designed studies are required to precise the incidence and the impact of JHR as well as the severity and rates between various antibiotics.

Citation: Guerrier G, D'Ortenzio E (2013) The Jarisch-Herxheimer Reaction in Leptospirosis: A Systematic Review. PLoS ONE 8(3): e59266. doi:10.1371/journal.pone.0059266

Editor: Odir A. Dellagostin, Federal University of Pelotas, Brazil

Received: December 17, 2012; **Accepted:** February 13, 2013; **Published:** March 26, 2013

Copyright: © 2013 Guerrier, D'Ortenzio. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Institut Pasteur de Nouvelle Calédonie funded this work. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: guerriergilles@gmail.com

These authors contributed equally to this work.

Introduction

Leptospirosis is a widespread zoonotic disease of worldwide distribution caused by pathogenic *Leptospira*, a spirochetal organism that is transmitted to humans by exposure to urine of infected mammalian reservoirs such as rodents or wild and domestic animals [1]. The disease has a wide-ranging clinical spectrum from asymptomatic forms to severe presentations. The latter are estimated to occur in 5–15% of all human infections and commonly combine jaundice, renal failure, myocarditis and/or haemorrhage [2] with a significant mortality rate. While a single experimental study demonstrated the relevance of antibiotic administration even in late stages of leptospirosis [3] the benefit of antibiotic therapy on mortality lacks of clinical evidence [4]. However, the current treatment guidelines still rely on antibiotic administration regardless of stage or severity of the disease [5]. Initiation of chemotherapy in spirochetal diseases may precipitate a febrile inflammatory reaction [6,7], known as the Jarisch-Herxheimer reaction (JHR), originally described in patients with syphilis receiving mercury treatment [8,9]. This reaction is characterised by an acute inflammatory response associated with the release of large amounts of cytokines, resulting from clearance

of spirochetes from the circulation [6,7]. Prevalence, clinical manifestations and outcome of JHR have been well studied in syphilis [10], Lyme disease [11], tick-born relapsing fever [12] and louse-born relapsing fever [13–15]. However, a clear idea of importance JHR in leptospirosis is lacking. In order to quantify frequency and impact of JHR in leptospirosis, we systematically reviewed the published literature and put it in a broader perspective, identifying data gaps that should be addressed.

Methods

Search Strategy and Selection Criteria

The inclusion criteria were as follows: all publications on leptospirosis treated with antibiotics providing information about the occurrence of early adverse events including JHR were selected. When the original article was not obtainable but the abstract containing the requested information was, the publication was included in the analysis. To avoid multiple counting (duplication) of identical procedures and cases, follow-up publications on identical procedures and cases were traced and excluded. The electronic database PubMed was searched with the keywords “leptospirosis” and “Leptospira”. A second search was performed

Table 1. Case reports and case series of Jarisch-Heirxheimer reaction after admistration of antibiotics for the treatment of leptospirosis.

References	Country	Study design	Treatment administered	Delay before JHR	Symptoms	Outcome
Lau CL et al. Emerg Infect Dis 2012	American Samoa	Case report (n = 1)	penicillin	1 hour	Increase in fever Rigors Severe Headache	Discharged
Markham R et al. Med J Aust 2012	Australia	Case report (n = 1)	benzylpenicillin and ceftriaxone	2 hours	Tachycardia Tachypnea Hypertension Severe rigors	Discharged
Hashimoto T et al. J Thorac Imaging 2012	Japan	Case report (n = 1)	ceftriaxone	2 days	Pulmonary deterioration	Discharged
Masuda K et al. Kansenshogaku Zasshi 2010	Japan	Case report (n = 1)	cefepime	Unknown	Chills Fever Hypotension	Discharged
Narita M et al. Am J Trop Med Hyg 2005	Japan	Case series (n = 6)	ampicillin (n = 6)	Unknown	Rigors Hypotension	Unknown
Tattevin P et al. Eur J Clin Microbiol Infect Dis 2003	France	Case report (n = 1)	amoxicillin	4 hours	Headache Hypotension Tachycardia Fever Nuchal rigidity	Discharged
Swiader L et al. La Presse Med 1995	France	Case report (n = 1)	penicillin	8 hours	Headache Photophobia Nuchal rigidity	Unknown
Vaughan C et al. Postgrad Med J 1994	Ireland	Case series (n = 3)	benzylpenicillin (n = 2) ampicillin (n = 1)	4 to 5 hours	Increase in fever Hypotension	Died (n = 1) Discharged (n = 2)
Emmanouilides CE et al. Clin Infect Dis 1991	USA	Case report (n = 1)	penicillin	Few hours	Fever Chills Hypotension Respiratory distress	Discharged
Friedland JS et al. Rev Infect Dis 1991	UK	Case series (n = 2)	benzylpenicillin (n = 2)	5 hours 4 hours	Fever Severe rigor Hypotension Abdominal pain Headache Fever Profuse vomiting	Discharged
Winearls CG et al. Q J Med 1984	UK	Cases series (n = 3)	penicillin (n = 3)	Unknown	Sharp rise in temperature Rigor	Discharged
Mackay-Dick J et al. J Royal Army Medical Corps 1957	Malaysia	Case series (n = 70)	penicillin (n = 70)	Unknown	Fever (n = 59) Aggravation of classical symptoms (n = 31) Hypotension (n = 58) Oligo-anuric (all)	Discharged
Crooks J et al. Br Med J	Scotland	Case report (n = 1)	penicillin	2 ½ hours	Rigor Fever	Discharged

Table 1. Cont.

References	Country	Study design	Treatment administered	Delay before JHR	Symptoms	Outcome
1955					Weakness	
					Low blood pressure	

doi:10.1371/journal.pone.0059266.t001

for “penicillin”, “amoxicillin”, “ampicillin”, “cefepime”, “ceftriaxone” and “antibiotics”. A final search was done with “Jarisch-Herxheimer reaction”. Search results were assessed using different combinations. The search was not limited by study design, language or date of publication. A similar search was conducted with Embase. The Cochrane Library was also searched for leptospirosis publications. World Health Organisation (WHO) websites and international or accessible national guidelines in endemic countries were searched, and the latest editions of five standard textbooks in infectious and tropical diseases were studied [16–20]. Titles and abstracts were screened, and potentially relevant case reports, case series and trials were further evaluated. Reports were included if they assessed the occurrence of JHR in leptospirosis. Reference lists of included articles were screened for further relevant publications. Study identification and data extraction were conducted independently and cross-checked for accuracy by both authors.

Data Abstraction and Methodological Assessment

The primary outcome was JHR occurrence or rates measured at time points described by investigators. Secondary outcomes include renal failure rates and mortality. Serovars were also recorded. It was impossible to retrospectively grade the severity of the JHR due to the lack of a standardized definition of the events.

Results

Twenty eight publications met our inclusion criteria, with a total number of 976 leptospirosis cases treated with antibiotics. One publication was a follow-up report on identical case series and therefore excluded from analysis [21]. Abstract or full text was not found for an additional publication which was also excluded [22]. One previous review dedicated to the topic was published in 1991 without systematically assessing the literature [23]. Among the selected studies, 11 patients presenting with concomitant infections were excluded [24].

Studies Characteristics and JHR Occurrence

Eight case reports [25–32] and five case series [23,33–36] published between 1955 and 2012 were reviewed and analyzed. All selected patients had a laboratory confirmed leptospirosis without concomitant infection and all were treated with antibiotics. A JHR was reported in 92 patients from 1 to 48 hours after administration of the first dose of antibiotics. The nature of antibiotics regimen given varied considerably: penicillin ($n=81$), ampicillin ($n=8$), ceftriaxone ($n=1$), cefepime ($n=1$) and penicillin and ceftriaxone ($n=1$). The most common features of described JHR were sudden onset of shivering or rigors ($n=6$), with rise in temperature ($n=9$), with ($n=8$) or without ($n=1$) a fall in blood pressure, occurring after administration of the first dose of antibiotics (Table 1). The largest case series accounted for 70 cases, out of 84 case-patients treated with penicillin in Malaya in 1957 [36].

Seven randomized control trials (RCT) [21,24,37–42], six non randomized control trials (NRCT) [43–48] were reviewed and analyzed (Table 2). With the exception of two RCTs [41,42], all studies monitoring the efficacy of antibiotics were not designed to assess adverse events linked to antibiotics including the JHR occurrence. The single study designed to monitor adverse events focusing on leptospirosis-related JHR was performed in 1986 by Watt et al. [42]. In this study, a single Herxheimer-like reaction was observed in a patient receiving saline placebo. The authors concluded that fear of a Herxheimer-like reaction should not dissuade clinicians from administering antibiotics to patients with leptospirosis. However, studied patients received up to four doses of parenteral antibiotic or had completed up to two days of an oral antibiotic regimen before inclusion.

JHR in National and International Guidelines, Textbooks and WHO/ILS Guideline

Among 3 screened national or international guidelines, all described JHR as a potential adverse event, although occurring with penicillin only [5,49,50]. Three textbooks out of the five consulted mentioned the possibility of JHR early after penicillin initiation [18–20].

JHR Management and Outcome

JHR management was not specified in 82 cases, but reported with no details in five cases (supportive care). In the remaining cases, the management consisted in fluid infusion, vasopressors ($n=1$), corticosteroids and inotropic support ($n=1$) and transient dialysis ($n=1$). Overall, eleven studies mentioned immediate outcome after JHR occurrence. Among them, all cases fully recovered with the exception of one JHR-related death [34].

JHR and Leptospira Serovars

Out of 13 case reports or case series reporting JHR, the pathogen serovar was stated in 18 patients. Nine different types of serovar were involved in JHR-patients, the most prevalent being Icterohaemorrhagiae ($n=6$).

Discussion

To the best of our knowledge, this systematic review is the first to formally appraise the measurement of JHR occurrence in leptospirosis treated with antibiotics. Publications describing JHR consist mainly of single case reports or case series. All RCTs included in our review but one were not designed to assess JHR incidence and therefore judged to be of poor methodological quality. Moreover, methods used to monitor JHR were not described in enough detail to reach predefined criteria and to determine whether there was a link between JHR occurrence and organ failure or death. Most reports were targeted at criteria such as duration of fever, clearance of spirochetes and length of stay. The single appropriately designed and detailed RCT had major

Table 2. Clinical trials for leptospirosis treatment in which adverse events were searched.

References	Country of study	Study design	Treatment	Any adverse events reported	Systematic assessment of JHR	Observation of JHR	Delay before JHR	Symptoms of JHR	Outcome
Phimda K et al. Antimicrob Agents Chemother 2003–2005	Thailand	RCT	doxycycline (n = 34) vs azythromycin (n = 35)	Rash Nausea Vomiting Diarrhea	Unknown	No	NA	NA	No death
Suputtamongkol Y et al. Clin Infect Dis 2001–2002	Thailand	RCT	penicillin (n = 52) vs cefotaxime (n = 59) vs doxycycline (n = 53)	Skin rash Dizziness	Unknown	No	NA	NA	4 deaths
Panaphut T et al. Clin Infect Dis 2000–2001	Thailand	RCT	ceftriaxone (n = 87) vs penicillin (n = 86)	No	Unknown	No	NA	NA	10 deaths
Costa E et al. Rev Inst Med Trop Sao Paulo 1997–1999	Brazil	RCT	penicillin (n = 125) vs placebo (n = 128)	No	Unknown	No	NA	NA	15 deaths
Daher EdF et al. Rev Inst Med Trop Sao Paulo 1996–1998	Brazil	NRCT	penicillin (n = 16) vs untreated (n = 19)	No	Yes	No	NA	NA	1 death
Marotto PCF et al. Am J Trop Med Hyg 1989–1995	Brazil	NRCT (retrospective analysis of children)	penicillin or ampicillin (n = 28) vs untreated (15)	No	Unknown	No	NA	NA	1 death
Watt G et al. J Infect Dis 1985–1986	Philippines	RCT	penicillin (n = 24) vs placebo (n = 16)	No	Yes	Yes (n = 1 in patient receiving placebo)	Unknown	Unknown	No death
Edwards CN et al. Am J Trop Med Hyg 1983–1986	Barbados	RCT	penicillin (n = 38) vs placebo (n = 41)	No	Unknown	Yes (n = 1 in patient receiving penicillin)	3 hours	Unknown	4 deaths
McClain BL et al. Ann Intern Med 1984	Panama Year not specified	RCT	doxycycline (n = 14) vs placebo (n = 15)	No	Yes	No	NA	NA	No death

Table 2. Cont.

References	Country Year of study	Study design	Treatment	Any adverse events reported	Systematic assessment of JHR	Observation of JHR	Delay before JHR	Symptoms of JHR	Outcome
Russel RW et al. Lancet 1958	Malaya Year not specified	NRCT	oxytetracycline (n = 27) vs placebo (n = 25)	Pyrexia Sore throat Erythema Urticaria	No	No	NA	NA	Unknown
Doherty RL et al. Australas annals of Medicine 1955	Australia Year not specified	NRCT	penicillin (n = 71) vs chloramphenicol (n = 12)	No	No	No	NA	NA	Unknown
Fairburn AC et al. Lancet 1956	Malaya Year not specified	NRCT	chloramphenicol+penicillin (n = 20) penicillin (n = 10) vs chloramphenicol (n = 14)	No	No	No	NA	NA	No death
Hall HE et al. Annals of Internal Medicine 1951	Puerto Rico Year not specified	NRCT	placebo (n = 22) penicillin (n = 5) vs chloramphenicol (n = 18)	No	No	No	NA	NA	2 deaths
			vs aureomycin (n = 13)						
			vs terramycin (n = 8)						
			vs streptomycin (n = 12)						
			vs aureomycin+streptomycin (n = 9)						
			vs aureomycin+cortison (n = 2)						
			vs untreated (n = 12)						

RCT: Randomized Control Trial; NRCT: Non Randomized Control Trial.
doi:10.1371/journal.pone.0059266.t002

limitations, including a small number of subjects and a lack of JHR confirmation.

However, this work will help to better estimate the potentially detrimental impact of antibiotics in leptospirosis and inform future management guidelines. Since there is little doubt that renal damage can be aggravated by vascular hypotension, particular attention should be paid to monitor blood pressure after initiation of antibiotics in suspected leptospirosis, especially in particular conditions such as pregnant, hypotensive or chronic renal insufficiency presenting patients. Of note, JHR is not reported in studies describing prognostic factors of leptospirosis but clinicians are usually facing complicated management [51,52].

Interestingly, a higher proportion of JHR occurred in early stage leptospirosis, suggesting a higher probability of the event before the natural clearance of spirochetes. It is postulated that the inflammatory process results from activation of the cytokine cascade during the degeneration of spirochetes. Therefore, the apparent lower proportion of JHR in patients with leptospirosis compared with patients diagnosed with other spirochetal diseases may be explained by a lower bacteraemia. This hypothesis requires further investigation for validation. JHR unobserved or unreported by clinicians is an additional reason for this ostensible reduced frequency. Conversely, a confusion of JHR symptoms with the aggravation of the leptospirosis itself may lead to overestimate the incidence of this event.

Interestingly, international guidelines for the management of leptospirosis briefly mention the occurrence of JHR with penicillin, omitting the details of its prevention, management or outcome. Moreover, antibiotics other than penicillin (e.g. ceftriaxone) are not evoked. Similarly, infectious diseases textbooks mentioning the event overlook this reaction with non-penicillin antibiotics. Noteworthy, recommended antibiotics for the management of leptospirosis vary considerably according to settings. For example, most commonly used drugs in the Philippines for mild leptospirosis is doxycycline while penicillin G is recommended for treating severe cases [50]; amoxicillin and cefotaxim are preferentially used in French Caribbean or in New Caledonia in mild and severe cases, respectively [53].

Some limitations in the current review should be acknowledged. First, potential publication bias is impossible to completely

exclude. Moreover, unpublished reports or reports that were not referenced in databases could have been missed despite the comprehensive search. Second, patients presenting with undetected co-infections may have been included in the study. Finally and most importantly, the diagnosis of JHR was not supported by any dosage of biological markers. In addition, the definition of JHR was not uniform, including in the largest case series published by Mackay-Dick [36] which lacks of details such as the onset of reaction. Therefore, although most of the cases presented a genuine JHR, some might have had a clinical aggravation related to the spirochetal disease regardless of the antibiotherapy.

Conclusions

The prevalence of JHR in leptospirosis treated with antibiotics is unknown and the awareness of this adverse event is insufficient. Although this review suggests that antibiotic treatment in patients with leptospirosis may result in less common and less severe JHR than in patients with other spirochetal diseases, all leptospirosis guidelines and local leptospirosis protocols should stress on systematic monitoring for clinical status early after antibiotic administration. We strongly recommend that patients receiving penicillin or other antibiotics for the management of leptospirosis to be monitored early after initiation of treatment to prevent any detrimental effects of a potential JHR. Since it is still controversial whether antimicrobials produce a beneficial effect in mild human leptospirosis, large well designed studies are required integrating a specific monitoring to precise the incidence and the impact of JHR as well as the severity and rates between various antibiotics.

Acknowledgments

The authors greatly acknowledge Patrick Lefèvre, MD, for sharing his experience on management of leptospirosis.

Author Contributions

Conceived and designed the experiments: GG ED. Performed the experiments: GG ED. Analyzed the data: GG ED. Contributed reagents/materials/analysis tools: GG ED. Wrote the paper: GG ED.

References

- Bharti AR, Nally JE, Ricaldi JN, Matthias MA, Diaz MM, et al. (2003) Leptospirosis: a zoonotic disease of global importance. *The Lancet Infectious Diseases* 3: 757–771.
- McBride AJ, Athanazio DA, Reis MG, Ko AI (2005) Leptospirosis. *Curr Opin Infect Dis* 18: 376–386.
- Spichler A, Ko AI, Silva EF, De Brito T, Silva AM, et al. (2007) Reversal of renal tubule transporter downregulation during severe leptospirosis with antimicrobial therapy. *Am J Trop Med Hyg* 77: 1111–1119.
- Brett-Major DM, Coldren R (2012) Antibiotics for leptospirosis. *Cochrane Database Syst Rev* 2: CD008264.
- World Health Organization WHO, International Leptospirosis Society ILS (2003) Human Leptospirosis: guidance for diagnosis, surveillance and control; Society WHO-IL, editor: World Health Organization. 122 p.
- Bryceson AD (1976) Clinical pathology of the Jarisch-Herxheimer reaction. *J Infect Dis* 133: 696–704.
- Pound MW, May DB (2005) Proposed mechanisms and preventative options of Jarisch-Herxheimer reactions. *J Clin Pharm Ther* 30: 291–295.
- Jarisch A (1895) Therapeutische Versuche bei Syphilis. *Wien Med Wochenschr* 45: 721–724.
- Herxheimer K, Krause D (1902) Ueber eine bei Syphilitischen vorkommende Quecksilberreaktion. *Dtsch Med Wochenschr* 28: 895–897.
- Shenep JL, Feldman S, Thornton D (1986) Evaluation for endotoxemia in patients receiving penicillin therapy for secondary syphilis. *JAMA* 256: 388–390.
- Maloy AL, Black RD, Seguro RJ, Jr. (1998) Lyme disease complicated by the Jarisch-Herxheimer reaction. *J Emerg Med* 16: 437–438.
- Dworkin MS, Anderson DE, Jr., Schwan TG, Shoemaker PC, Banerjee SN, et al. (1998) Tick-borne relapsing fever in the northwestern United States and southwestern Canada. *Clin Infect Dis* 26: 122–131.
- Bryceson AD, Cooper KE, Warrell DA, Perine PL, Parry EH (1972) Studies of the mechanism of the Jarisch-Herxheimer reaction in louse-borne relapsing fever: evidence for the presence of circulating *Borrelia* endotoxin. *Clin Sci* 43: 343–354.
- Warrell DA, Perine PL, Krause DW, Bing DH, MacDougall SJ (1983) Pathophysiology and immunology of the Jarisch-Herxheimer-like reaction in louse-borne relapsing fever: comparison of tetracycline and slow-release penicillin. *J Infect Dis* 147: 898–909.
- Fekade D, Knox K, Hussein K, Melka A, Lalloo DG, et al. (1996) Prevention of Jarisch-Herxheimer reactions by treatment with antibodies against tumor necrosis factor alpha. *N Engl J Med* 335: 311–315.
- Cook GC, Zumla AI (2008). *Manson's Tropical diseases*, 22nd edition. Edinburgh, United Kingdom: Elsevier Science Limited and W.B. Saunders. 1161.
- Warrell D, Cox TM, Firth J, Torok ME (2012) *Oxford Textbook of Medicine: Infection*: Oxford University Press. 864 p.
- Guerrant RL, Walker DH, Weller PF (2011). *Tropical Infectious Diseases*, 3rd Edition: Elsevier Science Limited and W. B. Saunders. 307.
- Mandell GL, Bennett JE, Dolin R (2009). *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases*, 7th Edition: Elsevier Science Limited. 3102.
- Gorbach SL, Bartlett JG, Blacklow NR (2004). *Infectious diseases*: Lippincott Williams and Wilkins. 1469.
- Watt G, Padre LP, Tuazon ML, Calubaquib C, Santiago E, et al. (1988) Placebo-controlled trial of intravenous penicillin for severe and late leptospirosis. *Lancet* 1: 433–435.
- Munnich D, Lakatos M (1972) Treatment of leptospirosis with Semicillin. *Ther Hung* 20: 152–155.

23. Friedland JS, Warrell DA (1991) The Jarisch-Herxheimer reaction in leptospirosis: possible pathogenesis and review. *Rev Infect Dis* 13: 207–210.
24. Phimda K, Hoontrakul S, Suttinont C, Chareonwat S, Losuwanaluk K, et al. (2007) Doxycycline versus azithromycin for treatment of leptospirosis and scrub typhus. *Antimicrob Agents Chemother* 51: 3259–3263.
25. Markham R, Slack A, Gerrard J (2012) The Jarisch-Herxheimer reaction in a patient with leptospirosis: a foreseeable problem in managing spirochaete infections. *Med J Aust* 197: 276–277.
26. Hashimoto T, Akata S, Park J, Harada Y, Hirayama Y, et al. (2012) High-resolution computed tomography findings in a case of severe leptospira infection (Weil disease) complicated with Jarisch-Herxheimer reaction. *J Thorac Imaging* 27: W24–26.
27. Masuda K, Uehara Y, Ono H, Furukawa K (2010) [A case of severe leptospirosis infection (Weil's disease) in Tokyo]. *Kansenshogaku Zasshi* 84: 59–64.
28. Tattevin P, Jaureguiberry S, Michelet C (2003) Meningitis as a possible feature of the Jarisch-Herxheimer reaction in leptospirosis. *Eur J Clin Microbiol Infect Dis* 22: 449.
29. Emmanouilides CE, Kohn OF, Garibaldi R (1994) Leptospirosis complicated by a Jarisch-Herxheimer reaction and adult respiratory distress syndrome: case report. *Clin Infect Dis* 18: 1004–1006.
30. Swiader L, Disdier P, Retornaz F, Pauzier F, Harle JR, et al. (1995) [Jarisch-Herxheimer reaction in leptospirosis]. *Presse Med* 24: 1753.
31. Lau CL, JM D (2012) Leptospirosis, American Samoa. *Emerg Infect Dis* [Internet].
32. Crooks J, Blair W (1955) L. canicola infection treated by penicillin. *Br Med J* 1: 885–887.
33. Narita M, Fujitani S, Haake DA, Paterson DL (2005) Leptospirosis after recreational exposure to water in the Yaeyama islands, Japan. *Am J Trop Med Hyg* 73: 652–656.
34. Vaughan C, Cronin CC, Walsh EK, Whelton M (1994) The Jarisch-Herxheimer reaction in leptospirosis. *Postgrad Med J* 70: 118–121.
35. Winckles CG, Chan L, Coghlan JD, Ledingham JG, Oliver DO (1984) Acute renal failure due to leptospirosis: clinical features and outcome in six cases. *Q J Med* 53: 487–495.
36. Mackay-Dick J, Robinson JF (1957) Penicillin in the treatment of 84 cases of leptospirosis in Malaya. *J R Army Med Corps* 103: 186–197.
37. Suputtamongkol Y, Niwattayakul K, Suttinont C, Losuwanaluk K, Limpiboon R, et al. (2004) An open, randomized, controlled trial of penicillin, doxycycline, and cefotaxime for patients with severe leptospirosis. *Clin Infect Dis* 39: 1417–1424.
38. Panaphut T, Domrongkitchaiporn S, Vibhagool A, Thinkamrop B, Sussaengrat W (2003) Ceftriaxone compared with sodium penicillin G for treatment of severe leptospirosis. *Clin Infect Dis* 36: 1507–1513.
39. Costa E, Lopes AA, Sacramento E, Costa YA, Matos ED, et al. (2003) Penicillin at the late stage of leptospirosis: a randomized controlled trial. *Rev Inst Med Trop Sao Paulo* 45: 141–145.
40. Edwards CN, Nicholson GD, Hassell TA, Everard CO, Callender J (1988) Penicillin therapy in icteric leptospirosis. *Am J Trop Med Hyg* 39: 388–390.
41. McClain JB, Ballou WR, Harrison SM, Steinweg DL (1984) Doxycycline therapy for leptospirosis. *Ann Intern Med* 100: 696–698.
42. Watt G, Padre LP, Tuazon M, Calubaquib C (1990) Limulus lysate positivity and Herxheimer-like reactions in leptospirosis: a placebo-controlled study. *J Infect Dis* 162: 564–567.
43. Russell RW (1958) Treatment of leptospirosis with oxytetracycline. *Lancet* 2: 1143–1145.
44. Doherty RL (1955) A clinical study of leptospirosis in North Queensland. *Australas Ann Med* 4: 53–63.
45. Fairburn AC, Semple SJ (1956) Chloramphenicol and penicillin in the treatment of leptospirosis among British troops in Malaya. *Lancet* 270: 13–16.
46. Hall HE, Hightower JA, Diaz Rivera R, Byrne RJ, Smadel JE, et al. (1951) Evaluation of antibiotic therapy in human leptospirosis. *Ann Intern Med* 35: 981–998.
47. Marotto PC, Marotto MS, Santos DL, Souza TN, Seguro AC (1997) Outcome of leptospirosis in children. *Am J Trop Med Hyg* 56: 307–310.
48. Daher EF, Nogueira CB (2000) Evaluation of penicillin therapy in patients with leptospirosis and acute renal failure. *Rev Inst Med Trop Sao Paulo* 42: 327–332.
49. Disease control division Doph, Ministry of Health, Malaysia (2011) Guidelines for the diagnosis, management, prevention and control of leptospirosis in Malaysia. 48 p.
50. Philippine College of Physicians (2011) Philippine Critical Practice Guidelines on the Diagnosis, Management and Prevention of leptospirosis. 66.
51. Dupont H, Dupont-Perdrizet D, Perie JL, Zehner-Hansen S, Jarrige B, et al. (1997) Leptospirosis: prognostic factors associated with mortality. *Clin Infect Dis* 25: 720–724.
52. Spichler AS, Vilaca PJ, Athanazio DA, Albuquerque JO, Buzzar M, et al. (2008) Predictors of lethality in severe leptospirosis in urban Brazil. *Am J Trop Med Hyg* 79: 911–914.
53. Collège de Maladies Infectieuses et Tropicales (2012) E. Pilly - Maladies infectieuses et tropicales, 23ème édition: Cmit Vivactis. 610 p.