



**HAL**  
open science

## **Terminin: a protein complex that mediates epigenetic maintenance of *Drosophila* telomeres.**

Grazia D Raffa, Laura Ciapponi, Giovanni Cenci, Maurizio Gatti

► **To cite this version:**

Grazia D Raffa, Laura Ciapponi, Giovanni Cenci, Maurizio Gatti. Terminin: a protein complex that mediates epigenetic maintenance of *Drosophila* telomeres.. *Nucleus*, 2011, 2 (5), pp.383-91. 10.4161/nucl.2.5.17873 . pasteur-00977757

**HAL Id: pasteur-00977757**

**<https://riip.hal.science/pasteur-00977757>**

Submitted on 11 Apr 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

This article was downloaded by: [Institut Pasteur]

On: 01 December 2014, At: 08:42

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Nucleus

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/kncl20>

### Terminin: A protein complex that mediates epigenetic maintenance of *Drosophila* telomeres

Grazia D. Raffa, Laura Ciapponi, Giovanni Cenci & Maurizio Gatti

Published online: 25 Oct 2011.

To cite this article: Grazia D. Raffa, Laura Ciapponi, Giovanni Cenci & Maurizio Gatti (2011) Terminin: A protein complex that mediates epigenetic maintenance of *Drosophila* telomeres, *Nucleus*, 2:5, 383-391, DOI: [10.4161/nucl.2.5.17873](https://doi.org/10.4161/nucl.2.5.17873)

To link to this article: <http://dx.doi.org/10.4161/nucl.2.5.17873>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Versions of published Taylor & Francis and Routledge Open articles and Taylor & Francis and Routledge Open Select articles posted to institutional or subject repositories or any other third-party website are without warranty from Taylor & Francis of any kind, either expressed or implied, including, but not limited to, warranties of merchantability, fitness for a particular purpose, or non-infringement. Any opinions and views expressed in this article are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor & Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

It is essential that you check the license status of any given Open and Open Select article to confirm conditions of access and use.

# Terminin

## A protein complex that mediates epigenetic maintenance of *Drosophila* telomeres

Grazia D. Raffa,<sup>1</sup> Laura Ciapponi,<sup>1</sup> Giovanni Cenci<sup>2</sup> and Maurizio Gatti<sup>1,\*</sup>

<sup>1</sup>Istituto Pasteur-Fondazione Cenci Bolognetti and Istituto di Biologia e Patologia Molecolari del CNR; Dipartimento di Biologia e Biotecnologie, Sapienza, Università di Roma; Roma, Italy; <sup>2</sup>Dipartimento di Biologia di Base ed Applicata; Università dell'Aquila; Coppito, L'Aquila, Italy

**Key words:** terminin, telomeres, telomere fusion, *Drosophila*

In most organisms, telomeres are extended by telomerase and contain GC-rich repeats. *Drosophila* telomeres are elongated by occasional transposition of specialized retroelements rather than telomerase activity, and are assembled independently of the sequence of the DNA termini. Recent work has shown that *Drosophila* telomeres are capped by a complex, we call terminin, which includes HOAP, HipHop, Moi and Ver; these are fast-evolving proteins that prevent telomere fusion, directly interact with each other, and appear to localize and function only at telomeres. With the possible exception of Ver that contains an OB fold domain structurally similar to the Stn1 OB fold, none of the terminin proteins is evolutionarily conserved outside the *Drosophila* species. Human telomeres are protected by the shelterin complex, which comprises six proteins that bind chromosome ends in a sequence-dependent manner. Shelterin subunits are not fast-evolving proteins and are not conserved in flies, but localize and function only at telomeres like the terminin components. Based on these findings, we propose that concomitant with telomerase loss *Drosophila* rapidly evolved terminin to bind chromosome ends in a sequence-independent fashion, and that terminin is functionally analogous to shelterin.

### Introduction


Telomeres of all eukaryotes are nucleoprotein complexes that protect the extremities of linear chromosomes from degradation and fusion, and counterbalance incomplete replication of terminal DNA. When telomeres are not properly capped, they are not recognized as natural chromosome ends but are sensed as double stranded DNA breaks (DSBs), triggering the DNA damage response that arrests the cell cycle until the DNA lesion is repaired. Uncapped telomeres may also undergo inappropriate DNA repair, leading to end-to-end fusions that ultimately result in chromosome breakage.<sup>1-3</sup> Telomeres also cope with the inability of DNA polymerase to fully replicate the DNA of chromosome termini, often referred to as the end replication problem. In most organisms, this problem is solved by telomerase, a

specialized reverse transcriptase that adds short GC-rich repeats to chromosome ends using an internal species-specific RNA template. In *Drosophila*, telomerase is absent<sup>4</sup> and telomeres are elongated by transposition of three specialized non-long-terminal repeat (LTR) retrotransposons, called *HeT-A*, *TART* and *TAHRE* (collectively abbreviated as HTT). These three elements transpose independently of each other and target individual telomeres at rates ranging from 10<sup>-2</sup> to 10<sup>-4</sup> per fly generation. Thus, most *Drosophila* chromosomes do not terminate with GC-rich repeats but carry HTT arrays of variable length (reviewed in refs. 5–7).

Several studies indicate that *Drosophila* telomeres are assembled independently of the HTT array and, more generally, independently of the sequence of the DNA termini (reviewed in ref. 8 and 9). In 1938 H.J. Muller observed that following X irradiation of males, terminal deletions could not be recovered. He concluded that chromosome ends are capped by special structures that he called telomeres and that are essential for chromosome stability and transmission.<sup>10,11</sup> However, subsequent studies showed that terminal deletions (TDs) can be recovered in several ways. For example, TDs were recovered following irradiation of females carrying a mutation in the mutator gene *mu2* and to lesser extent of wild type females.<sup>12-14</sup> These TDs can be transmitted for many generations without reacquiring HTT elements even if they undergo a progressive loss of terminal DNA due to the end replication problem.<sup>15</sup> It is now clear that TDs that do not end with HTT elements are capped by a neotelomere, which appears to have the same characteristics of the telomeres of intact chromosomes (reviewed in refs. 8 and 9). TDs with neotelomeres have also been recovered from mutational events occurred in the male germline. These events include X-ray induced breaks in the *Dp(1;f)1187* mini-chromosome,<sup>16</sup> the mobilization of a P element located near the telomere,<sup>17</sup> breakage of dicentric chromosomes generated by site specific recombination,<sup>18,19</sup> and induction of an enzymatic cut in an I-SceI site placed within a P element construct inserted near the telomere.<sup>20</sup> Collectively, these results demonstrate that the HTT elements are not required for fly telomere assembly and that virtually any DNA sequence has the ability to form the nucleoprotein complex that protects the ends of *Drosophila* chromosomes.

The structural features of the HTT elements, the molecular mechanisms underlying their transposition and their regulation

\*Correspondence to: Maurizio Gatti; Email: maurizio.gatti@uniroma1.it  
Submitted: 07/01/11; Revised: 08/24/11; Accepted: 08/25/11  
<http://dx.doi.org/10.4161/nucl.2.5.17873>



**Figure 1.** Mutations in genes encoding terminin components display frequent TFs. (A) wild type control metaphase. (B–D) Examples of TFs observed in larval brains from *ver* (B) and *moi* (C and D) mutants. Mutants in *cav*, *ver* and *moi* show two types of TFs: single TFs (STFs), in which a single telomere associates with either its sister or a nonsister telomere; and double TFs (DTFs), wherein a pair of sister telomeres joins with another pair. STFs and DTFs are likely to be generated during the S-G2 and the G<sub>1</sub> phase, respectively. In wild type cells, the frequency of TFs is less than 0.01/cell, whereas *cav*, *ver* and *moi* mutants exhibit approximately 5 TFs/cell, most of which are DTFs. (B) Metaphase containing 2–2 (arrow) and 4–4 (arrowhead) dicentric chromosomes and a dicentric ring involving both X chromosomes (asterisk), all generated by DTFs. (C) Metaphase showing a 4–4 DTA (asterisk), a 2–2 dicentric ring chromosome (arrowhead) and a 3–3 DTF (diamond); the XL and 3R telomeres exhibit sister union STFs (arrows). (D) Metaphase with a multi-centric chromosome (arrow) containing 3–3 and XR-XR DTFs and 3 STFs involving both XL telomeres and individual telomeres of chromosomes 2 and 3 (the metaphases shown are from figures in ref. 31 and 32).

by small RNAs have been reviewed in recent excellent articles.<sup>5-7</sup> Here, we describe the genes/proteins required for *Drosophila* telomere capping, with a focus on terminin, a multiprotein complex that evolved after telomerase loss to bind fly telomeres in a sequence-independent fashion.

### Telomere Capping Complexes in Organisms with Telomerase

In organisms with telomerase, telomeres associate with capping complexes that specifically bind the telomeric repeats generated by telomerase activity. In *S. cerevisiae*, telomeres are protected by the Rap1-Rif1-Rif2 complex that associates with the telomeric DNA duplex (dsDNA) through its Rap1 subunit, and by the Cdc13-Stn1-Ten1 complex (CST) that interacts with the 3' single stranded telomere overhang.<sup>21,22</sup> The three subunits of the CST complex all contain OB-fold domains and interact with each other to form an RPA-like complex that binds the 3' single stranded overhang via its Cdc13 subunit.<sup>22-25</sup>

Human telomeres terminate with a single stranded overhang of tandem TTAGGG repeats, which loops back invading the

anterior TTAGGG duplex, thus creating a telomeric DNA loop (t-loop). A complex of six proteins, called shelterin, specifically associates with the TTAGGG repeats (reviewed in ref. 1). Three of the shelterin subunits directly interact with the TTAGGG repeats; TRF1 and TRF2 bind the TTAGGG duplex, and POT1 binds the 3' overhang. TRF1, TRF2 and POT1 are interconnected by TIN2 and TPP1, and TRF2 interacts with hRap1, a distant homolog of *S. cerevisiae* Rap1 with no DNA binding ability.<sup>1</sup> The shelterin subunits share three properties that distinguish them from the non-shelterin telomere-associated proteins: they specifically localize to telomeres; they are abundant at telomeres throughout the cell cycle; and their functions are limited to telomere maintenance.<sup>1</sup>

The CST and shelterin complexes are evolutionarily conserved, even if they vary in composition and architecture in different phyla. The Stn1 and Ten1 subunits of the CST complex are conserved in *S. pombe*, plants and humans, while shelterin-like elements are found in *S. pombe* and plants but not in *S. cerevisiae*.<sup>26-29</sup> *S. pombe* and plants have both shelterin-like and CST-like complexes, both of which are required for telomere protection. The two complexes are present also in humans and are thought to collaborate in telomere protection. However, the human CST complex does not share the shelterin properties and appears to have a relatively minor role in telomere capping.<sup>28,29</sup>

The shelterin and CST components of yeast, plant and mammalian telomeres interact with several conserved polypeptides required for telomere function. These polypeptides include many proteins involved in the DNA damage response and in DNA repair such as the ATM and Chk2 kinases, the Ku70/80 heterodimer, the MRE11/RAD50/NBS1 (MRN) complex, Rad51, the ERCC1-XPF and MUS81 endonucleases, the Apollo exonuclease and the RecQ family members WRN and BLM, which are mutated in the Werner and Bloom syndromes, respectively (reviewed in refs. 1, 2, 26 and 30). In addition, yeast and mammalian telomeres are enriched in proteins that are homologous to *Drosophila* HP1. All non-shelterin and non-CST proteins localize and function not only at telomeres but also elsewhere in the cell.<sup>1,2,26,30</sup>

### *Drosophila* Telomeres are Protected by the Terminin Complex

The identification of *Drosophila* proteins required for telomere protection has mainly relied on the isolation of mutants that display frequent telomeric fusions (TFs) in larval brain cells (Fig. 1). The molecular characterization of the genes specified by these mutants identified ten loci that are required to prevent end-to-end fusion (Table 1). These are *Su(var)205* and *caravaggio* (*cav*) that encode HP1 and HOAP (HP1/ORC-associated protein), respectively;<sup>33,34</sup> *UbcD1* that encodes an E2 enzyme involved in protein ubiquitination;<sup>35</sup> the *Drosophila* homologs of the *ATM*, *RAD50*, *MRE11* and *NBS1* genes;<sup>36-43</sup> *without children* (*woc*) that specifies a putative transcription factor;<sup>44</sup> *modigliani* (*moi*; also called *DTL*) that encodes a nonconserved HOAP-binding protein;<sup>31,45</sup> and *verrocchio* (*ver*) that specifies an OB-fold containing protein structurally homologous to

**Table 1.** *Drosophila* genes required to prevent telomere fusion

Gene name	Protein name	Protein full name	Function outside of telomeres	References
<i>cav</i>	HOAP	HP1-ORC-Associated Protein	None known	34
<i>hiphop</i>	HipHop	HP1-HOAP-interacting protein	None known	20
<i>moi</i>	Moi	Modigliani	None known	31, 45
<i>ver</i>	Ver	Verrocchio	None known	32
<i>Su(var)205</i>	HP1	Heterochromatin Protein 1	Heterochromatin regulation; transcription factor	33, 46
<i>eff</i>	UbcD1	Ubiquitin Conjugating Enzyme D1	E2 ubiquitin conjugating enzyme	35, 59
<i>woc</i>	Woc	Without Children	Transcription factor	44
<i>mre11</i>	Mre11	Meiotic recombination 11	DNA repair; Component of the MRN complex	36, 37
<i>rad50</i>	Rad50	Radiation sensitive 50	DNA repair; Component of the MRN complex	37
<i>nbs</i>	Nbs	Nijmegen breakage syndrome	DNA repair; Component of the MRN complex	41–43
<i>tefu</i>	ATM	Ataxia Telangiectasia Mutated	Kinase; DNA damage response	36, 38–41, 61
<i>mei-41</i> (1)	ATR	Ataxia Telangiectasia Related	Kinase; DNA damage response	41–43
<i>mus-304</i> (1)	ATRIP	ATR Interacting Protein	DNA helicase; DNA damage response	41, 43
<i>armi</i> (2)	Armi	Armitage	helicase; piRNA biogenesis	69
<i>aub</i> (2)	Aub	Aubergine	piRNA biogenesis	69

(1) Mutations in *mei-41* or *mus-304* do not cause TFs but genetically interact with mutations in *tefu*, so that *mei-41 tefu* and *mus-304 tefu* double mutants exhibit TF frequencies that are much higher than those seen in *tefu* single mutants. (2) Mutations in *armi* and *aub* cause TFs only during embryogenesis.


STN1.<sup>32</sup> An additional protein required to prevent telomere fusion, called HP1-HOAP interacting protein (or HipHop), was recently identified among the polypeptides that co-precipitate with HOAP.<sup>20</sup>

Mutations in *caravaggio* (HOAP), *modigliani* and *verrocchio* cause very high frequencies of TFs (~5 per cell), often producing multicentric linear chromosomes that resemble little “trains” of chromosomes. The genes specified by these mutations have been named to reflect this phenotype, as the three Italian trains that are dubbed with the names of these famous painters. The HOAP, Ver and Moi proteins directly interact with each other in GST pull-down assays; HOAP and Moi also bind HP1 but Ver does not. Immunolocalization experiments have shown that HOAP is specifically associated with the telomeres of both mitotic and polytene chromosomes. An analysis of GFP-Moi and Ver-GFP localization on polytene chromosomes showed that these proteins are exclusively enriched at telomeres where they precisely colocalize with HOAP and HP1. However, GFP-Moi and Ver-GFP could not be detected at mitotic chromosome ends, probably due to their very low abundance.<sup>31,32</sup> These results indicate that HOAP, Moi and Ver form a complex that accumulates only at telomeres of both interphase (polytene) and mitotic chromosomes. In addition, available data indicate that HOAP, Moi and Ver function primarily if not exclusively at telomeres. Thus the HOAP-Moi-Ver complex, which has been named terminin (after the name of Rome’s train station), has the same properties as human shelterin and is likely to be a functional analog of shelterin.<sup>31,32</sup> HipHop directly interacts with both HOAP and HP1, although it is currently unknown whether it also binds Moi and Ver. HipHop specifically localizes at both mitotic and polytene chromosome telomeres and appears to function only at telomeres.<sup>20</sup> Thus, HipHop is likely to be an additional terminin component. Remarkably, HOAP, HP1 and HipHop all localize


to the extremities of various types of terminally deleted chromosomes demonstrating that these proteins bind chromosome ends independently of the sequence of terminal DNA.<sup>19,20,33,34</sup> It should be noted that despite its direct interaction with HOAP, HipHop and Moi, HP1 should not be considered as a terminin component, because it does not localize exclusively at telomeres and has multiple telomere-unrelated functions (reviewed in refs. 46 and 47).

The structural and functional information on the terminin proteins is still rather limited; the architecture of the complex and the possible roles of its components are not well defined. Biochemical analyses showed that HOAP and HipHop are mutually dependent for their stability, as loss of one protein reduces the amount of the other.<sup>20</sup> In contrast, HOAP, Moi and Ver do not appear to be interdependent for stability, as loss of one protein does not destabilize the others. However, recruitment of terminin proteins to the telomeres is governed by precise dependencies: localization of Moi and Ver at chromosome ends requires HOAP, while Moi and Ver are mutually dependent for their association with telomeres. HOAP, HipHop and Moi do not exhibit structural features that help define their roles in telomere protection. BLAST searches did not detect HOAP, HipHop or Moi homologs outside the *Drosophilidae* insects. In addition, these proteins do not appear to contain known functional domains, although HOAP is thought to carry an HMG-like domain.<sup>48</sup> Previous studies showed that HOAP binds double stranded DNA of different sequence, although with different affinities.<sup>48</sup> Ver contains an OB-fold domain that shares structural similarity with the OB fold domain of Rpa2/Stn1 and binds single stranded DNAs of different sequence (reviewed in ref. 32; our unpublished results). Substitution of four critical amino acids in the Ver OB fold abolished the DNA binding ability of the protein (our unpublished results). When this mutated Ver version was expressed in flies,





**Figure 2.** Distribution of HOAP and HipHop over the terminal DNA of a *Drosophila* chromosome ending with the *white* gene sequence (see text and ref. 20 for detailed explanation). Protein localization was determined by chromatin immunoprecipitation using anti-HOAP or anti-HipHop antibodies. Note that HOAP and HipHop exhibit similar distributions with the amount of each protein increasing from -11 to -1 kb from the end of the chromosome to drop in the terminal kb (adapted from Fig. 6 in ref. 20; with permission of the authors).



**Figure 3.** A tentative model for the molecular organization of *Drosophila* telomeres. We propose that the very ends of the chromosomes are capped by terminin, which includes HOAP, HipHop (HHop), Moi and Ver. HOAP and Ver bind double-stranded and single-stranded DNA, respectively, but it is currently unknown whether HipHop and HOAP directly associate with DNA. Moi and Ver would be absent from the telomeric DNA duplex proximal to the single stranded overhang; this duplex, however, would be associated with HOAP-HipHop. *Drosophila* telomeres are also enriched in HP1 and Woc, but these proteins are not terminin components because they associate with multiple polytene bands and play functions outside the telomeres. The UbcD1 (Ub) protein is enriched at the telomere region but appears to concentrate near the TAS rather than at the ends of the chromosomes (Cipressa and Cenci unpublished results). The MRN complex and the ATM kinase exhibit a rather uniform distribution along the *Drosophila* mitotic chromosomes and do not accumulate at polytene chromosome ends. Thus, UbcD1, Mre11, Rad50, Nbs and ATM were not included in the telomere region of the scheme, even if these proteins must function at telomeres, as mutations in the genes they specify cause telomeric fusions.

it was recruited at telomeres but was unable to prevent telomere fusion, suggesting that the DNA binding activity of the Ver OB fold is crucial for telomere protection.<sup>32</sup> It is not currently known whether Moi and HipHop can directly bind DNA. However, chromatin immunoprecipitation (ChIP) experiments using TD chromosomes terminating with the *white* gene sequence revealed that HOAP and HipHop are highly enriched within an 11-kb stretch of telomeric DNA. An analysis of the distribution of these proteins further showed their concentration increases from -11 kb to reach a peak at -1 kb but then drops in the terminal kb of telomeric DNA (Fig. 2). Although these results are rather fragmentary, they permit us to conceive a possible model for the interaction of terminin with telomeric DNA (Fig. 3). We would like to propose that HOAP and HipHop are primarily bound to the telomeric DNA duplex while Ver and Moi are associated with the single stranded overhang, which might span the terminal region of reduced HOAP/HipHop binding.

Although the terminin components are all required to prevent telomere fusion, they do not play identical roles at *Drosophila* telomeres. Previous work showed that HOAP-depleted telomeres trigger both the DNA damage response (DDR) and the spindle assembly checkpoint (SAC).<sup>49,50</sup> The SAC appears to be mediated by the BubR1 kinase, which accumulates at the uncapped telomeres in almost all *cav* mutant cells, but is never targeted to wild type telomeres. In the absence of Moi, Ver or HP1, telomeres have little or no ability to activate the DDR and trigger the SAC.<sup>31,32,49-51</sup> It is currently unknown whether HipHop depletion triggers the DDR and the SAC response. These results suggest that HOAP is crucial for protecting chromosome ends so as to prevent both telomere fusion and checkpoint responses. In contrast, Ver and Moi are not required to prevent checkpoint responses but are essential to hide chromosome ends from the DNA repair machineries that mediate telomere fusion. Interestingly, recent studies have shown that dysfunctional mouse telomeres also recruit BubR1, but it is unclear whether telomere-associated BubR1 can activate the SAC response.<sup>52</sup>

### The Roles of *Drosophila* Nonterminin Proteins Required for Telomere Protection

*Drosophila* telomere capping is not only ensured by terminin but also by a number of proteins that do not share the terminin properties; namely, proteins that do not localize or function only at telomeres. To date, we know 7 non-terminin proteins required for telomere protection from fusion events: HP1a, UbcD1, Mre11, Rad50, Nbs, ATM and Woc.

The most characterized nonterminin telomere-capping factor is HP1a. Besides HP1a, the *Drosophila* genome harbors 5 additional HP1 paralogs (HP1b, HP1c, HP1d, HP1e and Umbrea/HP6), none of which has been unambiguously shown to be required for telomere protection (reviewed in ref. 47; our unpublished results). In polytene chromosomes, HP1a is enriched at the telomeres, the chromocenter, the fourth chromosome and many euchromatic bands.<sup>33,53,54</sup> Consistent with this localization pattern,

HP1a binds diverse proteins involved in a variety of processes including telomere capping, gene silencing, DNA replication and repair, the maintenance of proper chromosome structure, and transcriptional regulation (reviewed in refs. 46, 47, 55 and 56). In *Drosophila*, HP1a is not only required for telomere protection but it is also involved in the control of telomere length. In stocks heterozygous for lethal mutations in *Su(var)205* (which encodes HP1a) the telomeres are much longer than those of wild type flies due to a dramatic elongation of the HTT array. This is a likely consequence of an increased *HetA* and *TART* transcription, which has been observed in both *Su(var)205* heterozygotes and homozygotes.<sup>57,58</sup> We never observed telomere elongation in stocks heterozygous for mutations in *cav*, *moi* or *ver*. In addition, our real time RT-PCR experiments did not detect substantial increases in *HetA* and *TART* transcription in homozygous *cav*, *moi* or *ver* mutant larvae (our unpublished results). Thus, the extant observations suggest that terminin is not implicated in the control of *Drosophila* telomere elongation.

*effete/UbcD1* was the first *Drosophila* gene shown to be required for prevention of telomere fusion.<sup>35</sup> *UbcD1* is an essential gene that encodes a highly conserved E2 ubiquitin conjugating enzyme implicated in several *Drosophila* cellular processes.<sup>35</sup> The UbcD1 protein associates with many polytene chromosome bands and is enriched at the telomere region of polytene chromosomes (Cipressa and Cenci G, unpublished observations). This suggests that failure to ubiquitinate one or more telomere proteins leads to fusogenic telomeres. However, we do not know whether UbcD1 interacts with terminin, and the telomere-associated target(s) of UbcD1 remain to be identified. Given that mutations in some of the proteasome components do not cause TFs (our unpublished results), we suspect that UbcD1-mediated ubiquitination of proteins involved in telomere protection is not required for their degradation but is instead a post-translational modification that ensures their proper capping function. Consistent with this idea, polytene chromosomes from *UbcD1* mutants and those from wild type controls display comparable telomeric concentrations of both HP1 and HOAP.<sup>59</sup> These results indicate that UbcD1 is required neither for HP1 or HOAP localization at telomeres nor for proteolysis of these proteins. However these findings do not exclude the possibility that either one or both of these proteins are ubiquitinated by UbcD1.

Studies performed in the last few years have shown that several proteins involved in DNA repair are also needed to prevent telomere fusion. Mutants in the *Drosophila mre11*, *rad50* and *nbs* genes die at late larval stages and exhibit both TFs and chromosome breakage in brain cells.<sup>36,37,41-43</sup> The *Drosophila* Mre11, Rad 50 and Nbs proteins are the fly orthologs of human MRE11, RAD50 and NBS1, which form the highly conserved MRN complex involved in both double-strand breaks (DSBs) repair and telomere maintenance. This complex mediates DSB repair by participating in both the homologous recombination (HR) and the nonhomologous end joining (NHEJ) pathways.<sup>60</sup> In humans, the MRN complex also associates with the TRF2 subunit of shelterin, facilitating telomerase recruitment and participating in detection and signaling of uncapped telomeres (reviewed in ref. 60). It is currently unknown whether the

components of the *Drosophila* MRN complex interact with terminin. However, mutations in the *rad50*, *mre11* and *nbs* genes strongly reduce HOAP accumulation at mitotic telomeres. These mutations also inhibit HOAP and HP1 localization at polytene chromosome ends.<sup>36,37,41-43</sup> Consistent with these findings, mutations in *mre11* prevent *Moi* localization at polytene chromosome telomeres.<sup>31</sup> Collectively, these results strongly suggest that terminin recruitment to telomeres requires the wild type function of the MRN complex. However, even in the absence of MRN activity, mitotic chromosomes retain the ability to recruit low levels of HOAP<sup>36,37,41-43</sup> and, presumably, of the other terminin components. These low amounts of terminin are likely to provide a partial protection of telomeres, as suggested by the finding that *rad50*, *mre11* and *nbs* mutants display much fewer TFs than *cav* mutants.<sup>34,36,37,41-43</sup>

Another DNA repair protein involved in *Drosophila* telomere protection is the ATM kinase. Null mutations in the *tefu/atm* gene result in lethality at late larval stages and cause both chromosome breakage and TFs in larval brain cells.<sup>36,38-40,61</sup> *tefu* mutations do not substantially affect HOAP localization at mitotic telomeres, although they might reduce HOAP accumulation at polytene telomeres.<sup>36,38</sup> Thus, loss of *Drosophila* ATM does not appear to cause telomere fusion by preventing terminin localization at chromosome ends. Null mutants in the ATR-encoding *mei-41* gene or in the *mus-304* gene that encodes the ATR-interacting protein ATRIP are both viable and their larval brain cells exhibit chromosome aberrations but not TFs.<sup>62-64</sup> However, *tefu mei-41* and *tefu mus-304* double mutants fail to recruit HOAP at telomeres and have significantly higher frequencies of TFs than those observed in *tefu* single mutants.<sup>41-43</sup> This suggests that ATM and ATR/ATRIP have partially redundant roles in telomere protection and that failure to phosphorylate a common target leads to deprotected telomeres. The nature of this target is currently unknown and it does not appear to be HOAP (ref. 41; our unpublished results).

The mechanism by which the combined action of the MRN complex, ATM and ATR-ATRIP leads to terminin recruitment to telomeres is unclear. The MRN complex plays a central role in detection and repair of DNA double strand breaks and mediates recruitment of ATM at the site of DNA damage (reviewed in ref. 60). Although MRN preferentially associates with the ends of linear DNA molecules *in vitro*,<sup>65</sup> the Rad50, Mre11 and Nbs protein are uniformly distributed along *Drosophila* chromosomes (ref. 37; our unpublished results), making it unlikely that MRN directly tethers terminin to telomeres. Thus, it has been hypothesized that interactions of DNA ends with the MRN complex and ATM-ATR result in conformational changes that facilitate terminin recruitment at telomeres.<sup>9,37,49</sup>

Protecting *Drosophila* telomeres from fusion also requires the wild type activity of the *without children (woc)* gene. *woc* encodes a zinc finger protein that interacts with HP1c and functions both in transcriptional regulation and telomere capping.<sup>44,66,67</sup> *Woc* is enriched at polytene chromosome telomeres and co-localizes with all euchromatic bands that associate with the initiating form of Pol II.<sup>44</sup> We do not know whether *Woc* interacts with terminin. However, *Woc* localization at telomeres is not affected by

mutations in *Su(var)205*, *cav*, *atm* or *rad50*, and mutations in *woc* do not affect HOAP and HP1 localization at chromosome ends. In addition, *woc* mutations do not dominantly affect telomere length.<sup>44</sup> These results indicate that the Woc function at telomeres is independent of those played by HOAP, HP1, ATM or Rad50, and that Woc is a transcription factor with telomere-capping properties, as is also the case for yeast Rap1.<sup>68</sup>

The *armitage* (*armi*) and *aubergine* (*aub*) genes play a tissue-specific role in Drosophila telomere stability.<sup>69</sup> *armi* and *aub* mutants are viable but female-sterile, as they cause maternal-effect embryonic lethality. Both genes are involved in the biogenesis of piRNAs; *armi* encodes an RNA helicase and *aub* a piRNA-binding Argonaute-like protein (reviewed in ref. 70). In embryos from *armi* and *aub* homozygous mothers, there are frequent anaphase bridges that are probably caused by telomeric fusions. Consistent with this interpretation, chromatin immunoprecipitation studies showed that mutation in both genes disrupt telomere binding of HOAP in the embryo.<sup>69</sup> Mutations in *armi* and *aub* also reduced an embryonic subpopulation of piRNAs that share some homology with telomeric retrotransposons, raising the possibility that these small RNAs may facilitate HOAP binding to the HTT array.<sup>69</sup> However, these piRNAs or other Armi- and Aub-dependent piRNAs (or the Armi and Aub proteins) must also facilitate HOAP binding to telomeres devoid of the HTT array, as diverse TD chromosomes are regularly transmitted during embryogenesis (see Introduction). It should also be noted that the Armi- and Aub-based mechanism of telomere protection is likely to be restricted to embryogenesis, as homozygous *armi* and *aub* individuals generated by heterozygous mothers are viable and do not exhibit TFs (our unpublished results).

### Evolution of Drosophila Telomere Proteins


There is a general consensus that the crucial event that led to the evolution of Drosophila telomeres was progressive loss of telomerase accompanied by the development of a regulated transposon-based mechanism for telomere elongation. It has also been suggested that the two mechanisms of telomere maintenance might have coexisted for some time.<sup>5,71</sup> This scenario is not difficult to envisage as in the silkworm *Bombyx mori*, which retains telomerase, there are two classes of retrotransposons that specifically insert into the telomere regions.<sup>72,73</sup> It is logical to assume that the transition from a telomerase-driven to a transposon-driven telomere elongation mechanism resulted in a divergence of terminal DNA sequences, accompanied by a strong selective pressure toward the evolution of sequence-independent telomere-binding factors. In agreement with this idea, none of the shelterin or the CST components is conserved in flies, and none of the terminin proteins, with the possible exception of Ver, has obvious homologs in yeasts, mammals or plants.<sup>20,31,32,34</sup> Thus, we hypothesized that concomitant with telomerase loss, Drosophila lost the shelterin and the CST homologs that bind DNA in a sequence-specific fashion, and evolved terminin to bind chromosome ends independently of the DNA sequence.<sup>31,32</sup> Ver exhibits a very limited amino acid sequence homology with Stn1, but contains an OB fold domain that is structurally similar to the Stn1

OB fold.<sup>32</sup> We speculate that Drosophila evolved Ver exploiting an Stn1 ancestor similar to the RPA proteins that bind ssDNA with no sequence specificity. The origin of HOAP, Moi and HipHop is unknown, as no conserved proteins that share amino acid sequence homology with these terminin components have been so far identified.

It is conceivable that following telomerase loss selective pressure on terminin proteins was much stronger than that exerted on other telomere proteins not specifically involved in capping. Proteins involved in diverse cellular processes are indeed expected to have functional constraints that prevent them from responding to selection with the same high rate of amino acid substitutions as polypeptides that are solely involved in the protection of chromosome ends. Therefore, one would hypothesize that proteins directly and exclusively involved in telomere capping evolved more rapidly than the other telomere-associated proteins. This hypothesis is verified by two findings. First, all Drosophila nonterminin proteins required to prevent telomere fusion have clear mammalian homologs, most of which have been implicated in telomere maintenance (HP1a, ATM, MRE11, RAD50, NBS1).<sup>31,32</sup> Second, HOAP, Moi, Ver and HipHop all exhibit a very high rate of nonsynonymous substitution per nonsynonymous site, and are therefore fast-evolving proteins. In contrast, none of the nonterminin telomere proteins, including HP1, appears to be a fast-evolving polypeptide.<sup>20,32,74</sup> Consistent with these findings, sequence analysis of proteins from *D. melanogaster* and 11 recently sequenced species<sup>75</sup> revealed that the terminin components are substantially more divergent than non-terminin proteins (Fig. 4). In conclusion, the rate of nonsynonymous substitutions in HOAP, Moi, Ver and HipHop is so high and distinctive that we propose to use it as an additional identifying criterion for both the extant terminin components and for terminin proteins that might emerge from future screens.

Recent work has shown that the terminin complex can vary in composition in different Drosophila tissues. In male meiotic and postmeiotic cells, including the sperm nuclei, HipHop is substituted by the product of the *ms(3)k81* gene (henceforth referred as *k81*).<sup>76,77</sup> *k81* is a duplication of the *hiphop* gene, presumably originated through a retroposition mechanism.<sup>78</sup> Phylogenetic analysis suggests that *k81* is a relatively young gene, as it is present only in the *melanogaster* subgroup of Drosophila species, although an independent *hiphop* duplication with male-biased expression was also found in *D. willistoni*.<sup>76</sup> K81 associates with the sperm telomeres and is specifically required for telomeric protection of male derived-chromosomes during embryogenesis; in embryos fathered by *k81* mutants, the paternal chromosomes display frequent TFs, ultimately leading to embryonic death.<sup>76,77</sup> The relationships between K81 and the other terminin proteins are unclear. In one study, K81 was found to colocalize with HOAP and HP1 in sperm nuclei of wild type males. HP1 and HOAP were absent from the sperm nuclei of *k81* mutants, suggesting that K81 is needed for the maintenance of capping complexes at the sperm telomeres.<sup>76</sup> In another study, mature sperm of wild type males were found to be devoid of both HP1 and HOAP signals, suggesting that the telomere-capping machinery of the sperm does not include HOAP and HP1.<sup>77</sup> It would be interesting





**Figure 4.** Identity percentages between the *D. melanogaster* proteins required for telomere capping and the homologous proteins from 11 *Drosophila* species (*D. mel*, *D. melanogaster*; *D. sim*, *D. simulans*; *D. sec*, *D. sechellia*; *D. yak*, *D. yakuba*; *D. ere*, *D. erecta*; *D. ana*, *D. ananassae*; *D. pse*, *D. pseudoobscura*; *D. per*, *D. persimilis*; *D. wil*, *D. willistoni*; *D. moj*, *D. mojvensis*; *D. vir*, *D. virilis*; *D. grim*, *D. grimshawi*). The identity percentage (id%; represented by the indicated color) is the percentage of identical matches between two amino acid sequences, calculated using the pairwise alignment EMBOSS Needle software.

to determine whether this machinery includes Moi and Ver and other proteins that, like K81, are required for telomere capping in sperm cells. Whatever the composition of the sperm telomere capping complex, the studies on K81 have clearly shown that during early embryogenesis there is substantial remodeling of the telomere structure; during this period K81 is substituted by HipHop and other telomere proteins are possibly recruited. It is also logical to speculate that the K81-HipHop transition period might be crucial for the formation of neotelomeres at the ends of TD chromosomes. It would be interesting to learn whether TD chromosomes, produced in males by means different from irradiation (see introduction), acquire a K81 cap in the sperm or are occasionally capable to escape from fusion events in the embryo and to recruit a normal capping complex during the K81-HipHop transition.

### Conclusions and Perspectives

We have described the main characteristics of 15 *Drosophila* proteins that are necessary to protect telomeres from fusion events (Table 1). In a review published in 2005, we estimated that the *Drosophila* genome contains at least 40 genes required to prevent telomere fusion.<sup>8</sup> The current results do not alter this conclusion and we still believe that there are many *Drosophila* telomere-capping genes that await to be discovered. The molecular and genetic analyses of the genes so far identified have shown that *Drosophila* telomeres are capped by terminin, a complex composed of fast-evolving proteins that specifically bind the telomeres. An important issue that needs to be elucidated by

future work is how terminin is recruited at the telomeres independently of the DNA sequence. We have learned that terminin recruitment requires the function of the MRN complex and the partially redundant activities of the ATM and ATR kinases. However the molecular mechanism underlying terminin recruitment remains elusive.

We have hypothesized that during the transition from a telomerase-based to transposon-based telomere elongation mechanism, *Drosophila* rapidly evolved terminin to bind chromosome ends independently of the DNA sequence. Consistent with this hypothesis, the *Drosophila* terminin proteins are not conserved in humans, while the shelterin proteins have no obvious *Drosophila* homologs. In contrast, *Drosophila* nonterminin and human nonshelterin telomere proteins are largely conserved from flies to mammals, and many of them play telomere-related functions in both organisms. These findings indicate that the main difference between *Drosophila* and human telomeres is in the protective complexes that specifically associate with the DNA termini. Thus, apart from the different mechanisms of elongation, *Drosophila* and human telomeres might not be as different as it is generally thought. The conservation in humans of the nonterminin telomere proteins further suggests that the identification of additional proteins of this type may lead the discovery of novel components of human telomeres.

### Acknowledgements

We are grateful to John C. Lucchesi for a critical reading of the manuscript. This work was supported in part by a grant from AIRC (Italian Association for Cancer Research) to M.G.

## References

- Palm W, de Lange T. How shelterin protects mammalian telomeres. *Annu Rev Genet* 2008; 42:301-34; PMID:18680434; DOI:10.1146/annurev.genet.41.110306.130350.
- Jain D, Cooper JP. Telomeric strategies: means to an end. *Annu Rev Genet* 2010; 44:243-69; PMID:21047259; DOI:10.1146/annurev-genet-102108-134841.
- O'Sullivan RJ, Karlseder J. Telomeres: protecting chromosomes against genome instability. *Nat Rev Mol Cell Biol* 2010; 11:171-81; PMID:20125188.
- Sasaki T, Fujiwara H. Detection and distribution patterns of telomerase activity in insects. *Eur J Biochem* 2000; 267:3025-31; PMID:10806402; DOI:10.1046/j.1432-033.2000.01323.x.
- Mason JM, Frydrychova R, Biessmann H. *Drosophila* telomeres: an exception providing new insights. *Bioessays* 2008; 30:25-37; PMID:18081009; DOI:10.1002/bies.20688.
- Capkova Frydrychova R, Biessmann H, Mason JM. Regulation of telomere length in *Drosophila*. *Cytogenet Genome Res* 2008; 122:356-64; PMID:19188706; DOI:10.1159/000167823.
- Pardue ML, DeBaryshe PG. *Drosophila* telomeres: A variation on the telomerase theme. *Fly (Austin)* 2008; 2:101-10; PMID:18820466.
- Cenci G, Ciapponi L, Gatti M. The mechanism of telomere protection: a comparison between *Drosophila* and humans. *Chromosoma* 2005; 114:135-45; PMID:16012858; DOI:10.1007/s00412-005-0005-9.
- Rong YS. Telomere capping in *Drosophila*: dealing with chromosome ends that most resemble DNA breaks. *Chromosoma* 2008; 117:235-42; PMID:18193446; DOI:10.1007/s00412-007-0144-2.
- Muller HJ. The remaking of chromosomes. *Collecting Net* 1938; 8:182-95.
- Muller HJ. An analysis of the process of structural change in chromosomes of *Drosophila*. *J Genet* 1940; 40:1-66; DOI:10.1007/BF02982481.
- Mason JM, Strobel E, Green MM. mu-2: mutator gene in *Drosophila* that potentiates the induction of terminal deficiencies. *Proc Natl Acad Sci USA* 1984; 81:6090-4; PMID:6435123; DOI:10.1073/pnas.81.19.6090.
- Mason JM, Champion LE, Hook G. Germ-line effects of a mutator, mu2, in *Drosophila melanogaster*. *Genetics* 1997; 146:1381-97; PMID:9258681.
- Dronamraju R, Mason JM. Recognition of double strand breaks by a mutator protein (MU2) in *Drosophila melanogaster*. *PLoS Genet* 2009; 5:1000473; PMID:19424425; DOI:10.1371/journal.pgen.1000473.
- Biessmann H, Carter SB, Mason JM. Chromosome ends in *Drosophila* without telomeric DNA sequences. *Proc Natl Acad Sci USA* 1990; 87:1758-61; PMID:2308935; DOI:10.1073/pnas.87.5.1758.
- Tower J, Karpen GH, Craig N, Spradling AC. Preferential transposition of *Drosophila* P elements to nearby chromosomal sites. *Genetics* 1993; 133:347-59; PMID:8382177.
- Levis RW. Viable deletions of a telomere from a *Drosophila* chromosome. *Cell* 1989; 58:791-801; PMID:2548737; DOI:10.1016/0092-8674(89)90112-8.
- Ahmad K, Golic KG. The transmission of fragmented chromosomes in *Drosophila melanogaster*. *Genetics* 1998; 148:775-92; PMID:9504924.
- Titen SW, Golic KG. Healing of euchromatic chromosome breaks by efficient de novo telomere addition in *Drosophila melanogaster*. *Genetics* 2010; 184:309-12; PMID:19897748; DOI:10.1534/genetics.109.109934.
- Gao G, Walser JC, Beaucher ML, Morciano P, Wesolowska N, Chen J, et al. HipHop interacts with HOAP and HP1 to protect *Drosophila* telomeres in a sequence-independent manner. *EMBO J* 2010; 29:819-29; PMID:20057353; DOI:10.1038/emboj.2009.394.
- Shore D, Bianchi A. Telomere length regulation: coupling DNA end processing to feedback regulation of telomerase. *EMBO J* 2009; 28:2309-22; PMID:19629031; DOI:10.1038/emboj.2009.195.
- Gao H, Cervantes RB, Mandell EK, Otero JH, Lundblad V. RPA-like proteins mediate yeast telomere function. *Nat Struct Mol Biol* 2007; 14:208-14; PMID:17293872; DOI:10.1038/nsmb1205.
- Mitton-Fry RM, Anderson EM, Hughes TR, Lundblad V, Wuttke DS. Conserved structure for single-stranded telomeric DNA recognition. *Science* 2002; 296:145-7; PMID:11935027; DOI:10.1126/science.1068799.
- Gelinas AD, Paschini M, Reyes FE, Heroux A, Batey RT, Lundblad V, et al. Telomere capping proteins are structurally related to RPA with an additional telomere-specific domain. *Proc Natl Acad Sci USA* 2009; 106:19298-303; PMID:19884503; DOI:10.1073/pnas.0909203106.
- Sun J, Yu EY, Yang Y, Confer LA, Sun SH, Wan K, et al. Stn1-Ten1 is an Rpa2-Rpa3-like complex at telomeres. *Genes Dev* 2009; 23:2900-14; PMID:20008938; DOI:10.1101/gad.1851909.
- Linger BR, Price CM. Conservation of telomere protein complexes: shuffling through evolution. *Crit Rev Biochem Mol Biol* 2009; 44:434-46; PMID:19839711; DOI:10.3109/10409230903307329.
- Lue NF. Plasticity of telomere maintenance mechanisms in yeast. *Trends Biochem Sci* 2009.
- Miyake Y, Nakamura M, Nabetani A, Shimamura S, Tamura M, Yonehara S, et al. RPA-like mammalian Ctc1-Stn1-Ten1 complex binds to single-stranded DNA and protects telomeres independently of the Pot1 pathway. *Mol Cell* 2009; 36:193-206; PMID:19854130; DOI:10.1016/j.molcel.2009.08.009.
- Surovtseva YV, Churikov D, Boltz KA, Song X, Lamb JC, Warrington R, et al. Conserved telomere maintenance component 1 interacts with STN1 and maintains chromosome ends in higher eukaryotes. *Mol Cell* 2009; 36:207-18; PMID:19854131; DOI:10.1016/j.molcel.2009.09.017.
- Oganesian L, Karlseder J. Telomeric armor: the layers of end protection. *J Cell Sci* 2009; 122:4013-25; PMID:19910493; DOI:10.1242/jcs.050567.
- Raffa GD, Siriaco G, Cugusi S, Ciapponi L, Cenci G, Wojcik E, et al. The *Drosophila* modigliani (moi) gene encodes a HOAP-interacting protein required for telomere protection. *Proc Natl Acad Sci USA* 2009; 106:2271-6; PMID:19181850; DOI:10.1073/pnas.0812702106.
- Raffa GD, Raimondo D, Sorino C, Cugusi S, Cenci G, Cacchione S, et al. Verrocchio, a *Drosophila* OB fold-containing protein, is a component of the terminin telomere-capping complex. *Genes Dev* 2010; 24:1596-601; PMID:20679394; DOI:10.1101/gad.574810.
- Fanti L, Giovinozzo G, Berloco M, Pimpinelli S. The heterochromatin protein 1 prevents telomere fusions in *Drosophila*. *Mol Cell* 1998; 2:527-38; PMID:9844626; DOI:10.1016/S1097-2765(00)80152-5.
- Cenci G, Siriaco G, Raffa GD, Kellum R, Gatti M. The *Drosophila* HOAP protein is required for telomere capping. *Nat Cell Biol* 2003; 5:82-4; PMID:12510197; DOI:10.1038/ncb902.
- Cenci G, Rawson RB, Belloni G, Castrillon DH, Tudor M, Petrucci R, et al. UbcD1, a *Drosophila* ubiquitin-conjugating enzyme required for proper telomere behavior. *Genes Dev* 1997; 11:863-75; PMID:9106658; DOI:10.1101/gad.11.7.863.
- Bi X, Wei SD, Rong YS. Telomere protection without a telomerase: the role of ATM and Mre11 in *Drosophila* telomere maintenance. *Curr Biol* 2004; 14:1348-53; PMID:15296751; DOI:10.1016/j.cub.2004.06.063.
- Ciapponi L, Cenci G, Ducau J, Flores C, Johnson-Schlitz D, Gorsky MM, et al. The *Drosophila* Mre11/Rad50 Complex is required to prevent both telomeric fusion and chromosome breakage. *Curr Biol* 2004; 14:1360-6; PMID:15296753; DOI:10.1016/j.cub.2004.07.019.
- Oikemus SR, McGinnis N, Queiroz-Machado J, Tukachinsky H, Takada S, Sunkel CE, et al. *Drosophila* atm/telomere fusion is required for telomeric localization of HP1 and telomere position effect. *Genes Dev* 2004; 18:1850-61; PMID:15256487; DOI:10.1101/gad.1202504.
- Silva E, Tiong S, Pedersen M, Homola EM, Royou A, Fasulo B, et al. ATM is required for telomere maintenance and chromosome stability during *Drosophila* development. *Curr Biol* 2004; 14:1341-7; PMID:15296750; DOI:10.1016/j.cub.2004.06.056.
- Song YH, Mirey G, Betson M, Haber DA, Sletten J. The *Drosophila* ATM ortholog, dATM, mediates the response to ionizing radiation and to spontaneous DNA damage during development. *Curr Biol* 2004; 14:1354-9; PMID:15296752; DOI:10.1016/j.cub.2004.06.064.
- Bi X, Srikanta D, Fanti L, Pimpinelli S, Badugu R, Kellum R, et al. *Drosophila* ATM and ATR checkpoint kinases control partially redundant pathways for telomere maintenance. *Proc Natl Acad Sci USA* 2005; 102:15167-72; PMID:16203987; DOI:10.1073/pnas.0504981102.
- Ciapponi L, Cenci G, Gatti M. The *Drosophila* Nbs protein functions in multiple pathways for the maintenance of genome stability. *Genetics* 2006; 173:1447-54; PMID:16648644; DOI:10.1534/genetics.106.058081.
- Oikemus SR, Queiroz-Machado J, Lai K, McGinnis N, Sunkel C, Brodsky MH. Epigenetic telomere protection by *Drosophila* DNA damage response pathways. *PLoS Genet* 2006; 2:71; PMID:16710445; DOI:10.1371/journal.pgen.0020071.
- Raffa GD, Cenci G, Siriaco G, Goldberg ML, Gatti M. The putative *Drosophila* transcription factor woc is required to prevent telomeric fusions. *Mol Cell* 2005; 20:821-31; PMID:16364909; DOI:10.1016/j.molcel.2005.12.003.
- Komonyi O, Schauer T, Papai G, Deak P, Boros IM. A product of the bicistronic *Drosophila melanogaster* gene CG31241, which also encodes a trimethylguanosine synthase, plays a role in telomere protection. *J Cell Sci* 2009; 122:769-74; PMID:19240120; DOI:10.1242/jcs.035097.
- Fanti L, Pimpinelli S. HP1: a functionally multifaceted protein. *Curr Opin Genet Dev* 2008; 18:169-74; PMID:18329871; DOI:10.1016/j.gde.2008.01.009.
- Vermaak D, Malik HS. Multiple roles for heterochromatin protein 1 genes in *Drosophila*. *Annu Rev Genet* 2009; 43:467-92; PMID:19919324; DOI:10.1146/annurev-genet-102108-134802.
- Shareef MM, King C, Damaj M, Badugu R, Huang DW, Kellum R. *Drosophila* heterochromatin protein 1 (HP1)/origin recognition complex (ORC) protein is associated with HP1 and ORC and functions in heterochromatin-induced silencing. *Mol Biol Cell* 2001; 12:1671-85; PMID:11408576.
- Cenci G, Ciapponi L. Telomere capping and cellular checkpoints: clues from fruit flies. *Cytogenetic and Genome Research* 2008.
- Musarò M, Ciapponi L, Fasulo B, Gatti M, Cenci G. Unprotected *Drosophila melanogaster* telomeres activate the spindle assembly checkpoint. *Nat Genet* 2008; 40:362-6; PMID:18246067; DOI:10.1038/ng.2007.64.
- Cenci G. *Drosophila* cell cycle under arrest: uncapped telomeres plead guilty. *Cell Cycle* 2009; 8:990-5; PMID:19270525; DOI:10.4161/cc.8.7.7960.
- Muñoz P, Blanco R, de Carcer G, Schoeffner S, Benetti R, Flores JM, et al. TRF1 controls telomere length and mitotic fidelity in epithelial homeostasis. *Mol Cell Biol* 2009; 29:1608-25; PMID:19124610; DOI:10.1128/MCB.01339-08.
- James TC, Eissenberg JC, Craig C, Dietrich V, Hobson A, Elgin SC. Distribution patterns of HP1, a heterochromatin-associated nonhistone chromosomal protein of *Drosophila*. *Eur J Cell Biol* 1989; 50:170-80; PMID:2515059.

54. Fanti L, Berloco M, Piacentini L, Pimpinelli S. Chromosomal distribution of heterochromatin protein 1 (HP1) in *Drosophila*: a cytological map of euchromatic HP1 binding sites. *Genetica* 2003; 117:135-47; PMID:12723693; DOI:10.1023/A:1022971407290.
55. Piacentini L, Fanti L, Negri R, Del Vecovo V, Fatica A, Altieri F, et al. Heterochromatin protein 1 (HP1a) positively regulates euchromatic gene expression through RNA transcript association and interaction with hnRNPs in *Drosophila*. *PLoS Genet* 2009; 5:1000670; PMID:19798443; DOI:10.1371/journal.pgen.1000670.
56. Chiolo I, Minoda A, Colmenares SU, Polyzos A, Costes SV, Karpen GH. Double-strand breaks in heterochromatin move outside of a dynamic HP1a domain to complete recombinational repair. *Cell* 2011; 144:732-44; PMID:21353298; DOI:10.1016/j.cell.2011.02.012.
57. Savitsky M, Kravchuk O, Melnikova L, Georgiev P. Heterochromatin protein 1 is involved in control of telomere elongation in *Drosophila melanogaster*. *Mol Cell Biol* 2002; 22:3204-18; PMID:11940677; DOI:10.1128/MCB.22.9.3204-18.2002.
58. Perrini B, Piacentini L, Fanti L, Altieri F, Chichiarelli S, Berloco M, et al. HP1 controls telomere capping, telomere elongation and telomere silencing by two different mechanisms in *Drosophila*. *Mol Cell* 2004; 15:467-76; PMID:15304225; DOI:10.1016/j.molcel.2004.06.036.
59. Cenci G, Siriacò G, Gatti M. The role of HeT-A and TART retrotransposons in *Drosophila* telomere capping. *Genetica* 2003; 117:311-8; PMID:12723710; DOI:10.1023/A:1022972902263.
60. Lamarche BJ, Orazio NI, Weitzman MD. The MRN complex in double-strand break repair and telomere maintenance. *FEBS Lett* 2010; 584:3682-95; PMID:20655309; DOI:10.1016/j.febslet.2010.07.029.
61. Queiroz-Machado J, Perdigo J, Simoes-Carvalho P, Herrmann S, Sunkel CE. *tef*: a mutation that causes telomere fusion and severe genome rearrangements in *Drosophila melanogaster*. *Chromosoma* 2001; 110:10-23; PMID:11398972; DOI:10.1007/s004120000116.
62. Gatti M. Genetic control of chromosome breakage and rejoining in *Drosophila melanogaster*: spontaneous chromosome aberrations in X-linked mutants defective in DNA metabolism. *Proc Natl Acad Sci USA* 1979; 76:1377-81; PMID:108678; DOI:10.1073/pnas.76.3.1377.
63. Hari KL, Santerre A, Sekelsky JJ, McKim KS, Boyd JB, Hawley RS. The *mei-41* gene of *D. melanogaster* is a structural and functional homolog of the human ataxia telangiectasia gene. *Cell* 1995; 82:815-21; PMID:7671309; DOI:10.1016/0092-8674(95)90478-6.
64. Brodsky MH, Sekelsky JJ, Tsang G, Hawley RS, Rubin GM. *mus304* encodes a novel DNA damage checkpoint protein required during *Drosophila* development. *Genes Dev* 2000; 14:666-78; PMID:10733527.
65. de Jager M, van Noort J, van Gent DC, Dekker C, Kanaar R, Wyman C. Human Rad50/Mre11 is a flexible complex that can tether DNA ends. *Mol Cell* 2001; 8:1129-35; PMID:11741547; DOI:10.1016/S1097-2765(01)00381-1.
66. Font-Burgada J, Rossell D, Auer H, Azorin F. *Drosophila* HP1c isoform interacts with the zinc-finger proteins WOC and Relative-of-WOC to regulate gene expression. *Genes Dev* 2008; 22:3007-23; PMID:18981478; DOI:10.1101/gad.481408.
67. Abel J, Eskeland R, Raffa GD, Kremmer E, Imhof A. *Drosophila* HP1c is regulated by an auto-regulatory feedback loop through its binding partner Woc. *PLoS ONE* 2009; 4:5089; PMID:19352434; DOI:10.1371/journal.pone.0005089.
68. Shore D. Telomeric chromatin: replicating and wrapping up chromosome ends. *Curr Opin Genet Dev* 2001; 11:189-98; PMID:11250143; DOI:10.1016/S0959-437X(00)00178-7.
69. Khurana JS, Xu J, Weng Z, Theurkauf WE. Distinct functions for the *Drosophila* piRNA pathway in genome maintenance and telomere protection. *PLoS Genet* 2010; 6:1001246; PMID:21179579; DOI:10.1371/journal.pgen.1001246.
70. Senti KA, Brennecke J. The piRNA pathway: a fly's perspective on the guardian of the genome. *Trends Genet* 2010; 26:499-509; PMID:20934772; DOI:10.1016/j.tig.2010.08.007.
71. Louis EJ. Are *Drosophila* telomeres an exception or the rule? *Genome Biol* 2002; 3:7.
72. Takahashi H, Okazaki S, Fujiwara H. A new family of site-specific retrotransposons, SART1, is inserted into telomeric repeats of the silkworm, *Bombyx mori*. *Nucleic Acids Res* 1997; 25:1578-84; PMID:9092665; DOI:10.1093/nar/25.8.1578.
73. Fujiwara H, Osanai M, Matsumoto T, Kojima KK. Telomere-specific non-LTR retrotransposons and telomere maintenance in the silkworm, *Bombyx mori*. *Chromosome Res* 2005; 13:455-67; PMID:16132811; DOI:10.1007/s10577-005-0990-9.
74. Schmid KJ, Tautz D. A screen for fast evolving genes from *Drosophila*. *Proc Natl Acad Sci USA* 1997; 94:9746-50; PMID:9275195; DOI:10.1073/pnas.94.18.9746.
75. Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, et al. Evolution of genes and genomes on the *Drosophila* phylogeny. *Nature* 2007; 450:203-18; PMID:17994087; DOI:10.1038/nature06341.
76. Dubruielle R, Orsi GA, Delabaere L, Cortier E, Couble P, Marais GA, et al. Specialization of a *Drosophila* capping protein essential for the protection of sperm telomeres. *Curr Biol* 2010; 20:2090-9; PMID:21093267; DOI:10.1016/j.cub.2010.11.013.
77. Gao G, Cheng Y, Wesolowska N, Rong YS. Paternal imprint essential for the inheritance of telomere identity in *Drosophila*. *Proc Natl Acad Sci USA* 2011; 108:4932-7; PMID:21383184; DOI:10.1073/pnas.1016792108.
78. Loppin B, Lepetit D, Dorus S, Couble P, Karr TL. Origin and neofunctionalization of a *Drosophila* paternal effect gene essential for zygote viability. *Curr Biol* 2005; 15:87-93; PMID:15668163; DOI:10.1016/j.cub.2004.12.071.