

Molecular characterization and antibiotic resistance of Salmonellain children with acute gastroenteritis in Abuja, Nigeria

Casmir Ifeanyichukwu Cajetan Ifeanyi, Bassey Enya Bassey, Nkiruka Florence Ikeneche, Nazek Al-Gallas

▶ To cite this version:

Casmir Ifeanyichukwu Cajetan Ifeanyi, Bassey Enya Bassey, Nkiruka Florence Ikeneche, Nazek Al-Gallas. Molecular characterization and antibiotic resistance of Salmonellain children with acute gastroenteritis in Abuja, Nigeria. Journal of Infection in Developing Countries, 2014, 8 (06), pp.712-9. 10.3855/jidc.4185. pasteur-01375049

HAL Id: pasteur-01375049 https://riip.hal.science/pasteur-01375049

Submitted on 2 Oct 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Original Article

Molecular characterization and antibiotic resistance of *Salmonella* in children with acute gastroenteritis in Abuja, Nigeria

Casmir Ifeanyichukwu Cajetan Ifeanyi¹, Bassey Enya Bassey², Nkiruka Florence Ikeneche³, Nazek Al-Gallas⁴

¹ Department of Veterinary Microbiology, Faculty of Veterinary Medicine, University of Abuja, Abuja, Nigeria

² World Health Organization Office, Abuja, Nigeria

³Department of Medical Microbiology, College of Health Science, University of Abuja, Abuja, Nigeria

⁴ Laboratoire de Controle des Eaux et Denrées Alimentaires, Institut Pasteur de Tunis, Tunisie

Abstract

Introduction: In Nigeria, acute gastroenteritis in children under five years of age is a major cause of mortality and morbidity; identification and characterization of microbial agents of acute gastroenteritis, including *Salmonella*, remains a powerful tool for effective management, surveillance, and control.

Methodology: Diarrheal stool samples were directly plated onto differential and selective media to isolate *Salmonella*. Extended-spectrum beta-lactamases were screened using the double disk diffusion technique and by PCR targeting the *bla*_{TEM} and *bla*_{CTX-M} genes. Pulsed-field gel electrophoresis (PFGE) was performed using the PulseNet Canada Laboratory protocol for molecular subtyping using the restriction enzymes *Xba*I and *Bln*I.

Results: The serotypes identified were *Salmonella enterica* serovar Zanzibar (n = 5), *Salmonella* Brancaster (n = 3), and one isolate of *Salmonella* Enteritidis (phage type 1). The following levels of resistance were found among the *Salmonella* strains: amoxicillin, five strains (55.6%); amoxicillin-clavulanic acid, two strains (22.2%); cephalexin, five strains (55.6%); and cefuroxime, five strains (55.6%). Intermediate resistance was found in five strains (55.6%) only to amoxicillin-clavulanic acid. All isolates were susceptible to nalidixic acid, ciprofloxacin, and ceftriaxone, and no ESBL-producing *Salmonella* were detected.

Conclusions: Our findings demonstrated the involvement of three *Salmonella* serovars in acute gastroenteritis; resistance to penicillins and cephalosporins was common.

Key words: acute gastroenteritis in children; pulse-field gel electrophoresis; Salmonella; antibiotic resistance; Nigeria

J Infect Dev Ctries 2014; 8(6):712-719. doi:10.3855/jidc.4185

(Received 08 September 2013 - Accepted 25 March 2014)

Copyright © 2014 Ifeanyi *et al.* This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Salmonella infections are associated with acute gastroenteritis, one of the most common alimentary diseases; they are caused by the consumption of contaminated water and food, especially meat [1]. The prevalence rate of acute diarrhea in Nigeria is 18.8%, one of the worst in sub-Saharan Africa. It accounts for over 16% of child deaths in Nigeria and an estimated 150,000 deaths – chiefly amongst children under five years of age - occur annually due to this disease, mainly caused by poor sanitation and hygiene practices. These diseases are especially common in developing countries with poor hygiene and sanitation and with limited access to safe drinking water [2]. The identification of the pathogenic organisms responsible for these diseases is highly crucial for surveillance, prevention, and control activities. TEM β-lactamases are encoded by plasmid-mediated bla_{TEM} gene clusters, which have various coding region frameworks and types of promoters [3]. The bla_{TEM} gene cluster encodes by plasmid mediation resistance mechanisms such as penicillinase hyper production, extendedspectrum beta-lactamases (ESBL), or inhibitorresistant TEM beta-lactamase [4]. Variations in sugar content of the O-polysaccharide contribute to the wide variety of antigenic types between species and even strains of Gram-negative bacteria [5]. The variability defined by immunologic characterization of two surface structures, O-polysaccharide (O antigen) and flagellin protein (H antigen), provides the basis for serotyping Salmonella spp. into serogroups [6]. Polymorphism in the bla_{TEM} genes has already been noted for clinical isolates with > 150 variants [7]. The highly polymorphic *bla*_{TEM} gene has been targeted as a

molecular marker to detect antimicrobial resistance associated with *Salmonella* serovars in acute gastroenteritis in Abuja.

Antimicrobial resistance in enteric pathogens is of great importance in developing countries, where the rate of diarrheal disease is high. The progressive increase in antimicrobial resistance among enteric pathogens in developing countries is becoming a critical area of concern. Among the bacterial causes of diarrheal illnesses, Salmonella spp. continues to be a major public health problem [8]. Although most Salmonella infections are self-limiting, serious sequelae, including systemic infection and death, can occur [9]. In potentially life-threatening cases, the antibiotics of choice are fluoroquinolones and extended-spectrum cephalosporins [10]. These cephalosporins are commonly used in the treatment of Salmonella infections in children because of their pharmacodynamic properties and the very low prevalence of resistance to these agents. Very few studies have reported on the problem of ESBL in Africa in general and in Nigeria in particular. There have been reports of SHV-12 in Salmonella enterica serotype in Mali [10] and TEM-3 in S. Tyhimurium in Morocco [11].

The aim of this study was to determine serotype, antibiotic resistance, and extended-spectrum betalactamase (ESBL) gene production of *Salmonella* species associated with childhood acute gastroenteritis in Federal Capital Territory Abuja, Nigeria. In addition, the study evaluated the molecular characteristics and clonal relatedness among the strains.

Methodology

Microbiological diagnosis

Stool samples were collected in sterile plastic cups, with prior instruction given for correct sampling. The samples were obtained by spontaneous emission, and given to the investigator. They were inoculated within two hours of collection onto selective and differential media: MacConkey (MAC) agar, Salmonella Shigella (SS) agar, and xylose lysine deoxycholate (XLD) agar, using a calibrated inoculating loop in the spread plate method. The media were then incubated aerobically at 35°C for 18 to 24 hours. Samples were also inoculated into selenite cystine enrichment broth - which favors the development of potentially pathogenic microorganisms and contains substances that inhibit the native flora [12] - and incubated at 35°C for 8 to 12 hours before re-inoculation onto MCK and SS agars.

Colonies with а presumptive Salmonella morphology as per the phenotypic characteristics in the different culture media were identified biochemically according to the following procedure [13]: five probable Salmonella colonies were selected from the SS or XLD agars, inoculated into 3.5 mL brain-heart infusion broth (BHI), incubated at 35°C for 10 minutes, and then inoculated onto the different media. Biochemical tests were then carried out for: the determination of glucose and lactose; gas formation and the production of ferrous sulfite from sodium thiosulfate in Kligler agar; the presence of cytochrome oxidase; citrate utilization; urea hydrolysis in Christensen's agar; the presence of phenylalanine deaminase; the liberation of indole; acid production from sugar fermentation; the decarboxylation of lysine, ornithine, and arginine; and the utilization of malonate as the sole carbon source. In addition, a nutrient agar plate was inoculated from each BHI broth and incubated at 35°C for 24 hours; the colonies were used for the oxidase test.

Serotyping and phage typing

Serotyping and phage typing of isolates were performed at the National Microbiology Laboratory (NML), Public Health Agency (PHC) of Canada according to the Kauffmann–White scheme and the recommendations of Edwards and Ewing [14].

Molecular characterization

The serovars that were resistant to either amoxicillin-clavulanic acid, cephalexin, or cefuroxime alone or in combinations were screened by PCR using two primer sets sequence specific for bla_{TEM} and $bla_{\text{CTX-M}}$ genes to amplify the DNA templatefor genes coding for ESBL production [3].

DNA was extracted from Salmonella isolates using Wizard Genomic DNA purification Kit (Promega, USA) according to manufacturer's Fitchburg, specifications. Oligonucleotide pairs that amplified 708 bp and 593 bp products of bla_{TEM} and $bla_{\text{CTX-M}}$ genes, respectively, were used (Table 1). PCR amplification employed a final volume of 25 uL containing 1.5 mM MgCl₂ 100uM deoxyribonucleoside triphosphate, 0.2µM of each oligonucleotide, and 1U of Tag DNA polymerase. The PCR amplification reaction for the detection of bla_{TFM} consisted of 36 cycles of 30 seconds of denaturation at 94°C, 30 seconds of hybridization at 42°C, and 60 seconds of extension at 72°C, with a final extension

Primer	Primer sequence	Primer target	Ampilicon size (bp)	Reference
_{TEM} - G _{TEM} -R	5'-TTGCTCACCCAGAAACGCTGGTG-3' 5'-TACGATACGGGAGGGCTTACC-3'	bla_{TEM}	708	Boyd et al., 2004
_{CTX-M} -F _{CTX-M} -R	5'-TTTGCGATGTGCAGTACCAGTAA-3' 5'-CGATATCGTTGGTGGTGCCATA-3'	bla _{CTX-M}	593	Boyd et al., 2004

Table 1. Primers used in identifying ESBL gene production in Salmonella serogroups

step at 72°C for 10 minutes [3]. The $bla_{\text{CTX-M}}$ gene was identified with PCR conditions similar to those described for bla_{TEM} , except for the alteration of the annealing temperature to 58°C [3]. The amplified products were visualized electrophoretically on 2% agarose gel [15]. For quality control, *S. branderup* H9812 ATCC BAA-664 was used.

Pulsed-field gel electrophoresis analysis

Pulsed-field gel electrophoresis (PFGE) was performed using the standardized protocol of PulseNet Canada Laboratory for molecular subtyping by using the restriction enzymes XbaI and BlnI according to the standardized protocol for *E. coli* (O157:H7). Salmonella switch times of 2.2-63.8 seconds was used for the molecular typing of Salmonella isolates [15]. S. braenderup strain H9812 (ATCC BAA 664) was used as a reference strain. DNA macrorestriction profiles were analyzed with BioNumerics software version 2.5 (Applied Maths, Saint Martens-Latem, Belgium) using a band tolerance of 1.5% and optimized at 1.0%. Dendrograms were constructed on the basis of the unweighted pair group method of averages, position tolerance of 1%. Clusters were defined as DNA patterns sharing $\geq 85\%$ similarity [16]. PFGE pattern designations were assigned according the PulseNet Canada database.

Antimicrobial susceptibility testing

Characterized strains were tested for susceptibility to the following antibiotics: nalidixic acid (30 µg), ciprofloxacin (5 µg), cephalexin (30 µg), cefuroxime $(30 \ \mu g)$, amoxicillin $(25 \ \mu g)$, ceftriaxone $(30 \ \mu g)$, and amoxicillin-clavulanic acid $(30 \mu g)$, using the standard Kirby Bauer disk diffusion susceptibility testing method. Escherichia coli ATCC 25922 was used as a control. Multidrug resistance (MDR) for this study was defined as resistance to two or more classes of Minimum antimicrobials agents. inhibitory concentrations (MICs) of isolates were performed by the broth microdilution method (Sensititre Automated Microbiology System; Trek Diagnostic Systems Ltd., Westlake, OH, USA). Using Clinical and Laboratory

Standards Institute (CLSI) guidelines, each organism was classified either resistant or susceptible [17].

Detection of ESBL production was done by a double disk diffusion technique with the following disks: three containing ceftazidime 30 µg (CAZ30 C), cefotaxime 30 µg (CTX30C), and cefpodoxime 30µg (CPD30C), and three with a combination of the same antibiotics with the addition of clavulanate: 30 ceftazidime µg/clavulanic acid 10 μg (CAZ30/CLAV10); cefotaxime 30µg/clavulanic acid (CTX30/CLAV10); cefpodoxime 10 μg 30µg/clavulanic acid 10 µg (CPD30C/CLAV10) (CA; Oxoid, Basingstoke, UK). The zones of inhibition of each isolate were tested on Mueller-Hinton agar plates (Oxoid, Basingstoke, UK). Interpretive criteria established by the CLSI were used to categorize the results of antimicrobial susceptibility testing and ESBL production [17].

Ethical considerations

Ethical approvals were obtained from the University of Abuja Teaching Hospital Gwagwalada, Abuja, National Hospital Abuja, and Health and Human Services Secretariat of the Federal Capital Administration. Informed consent was obtained from the parents of the patients who were to participate in the study. A questionnaire was used to collect epidemiological data following the principles proposed by the model ethical protocol in accordance with the Helsinki declaration [18].

Results

The nine strains of *Salmonella* isolates were typed into three serovars with specific antigenic formulae. The serotypes identified were *Salmonella* Zanzibar (n = 5), *Salmonella* Brancaster (n = 3), and one strain of *Salmonella* Enteritidis. The *S*. Enteritidis strain was phage type 1; the phage types of *S*. Zanzibar and *S*. Brancaster were not determined (Table 2).

There was no genetic diversity in the PFGE patterns of *Xba*I digested chromosomal DNA of the five isolates of *S*. Zanzibar, as all had the same restriction pattern.

Patient identity	Locality	Age groupType of(months)diarrhea		Organism identity/serotype	Antigentic formulae	Phage type
087	Municipal	4	Watery	Salmonella Zanzibar	3,10:k:1,5	ND
099	Peri-urban	12	Watery	Salmonella Brancaster	4,12,27:z29:-	ND
155	Municipal	12	Mucoid	Salmonella Zanzibar	3,10:k:1,5	ND
159	Municipal	24	Mucoid	Salmonella Zanzibar	3,10:k:1,5	ND
167	Municipal	18	Watery	Salmonella Brancaster	4,12,27:z29:-	ND
280	Municipal	4	Watery	Salmonella Brancaster	4,12,27:z29:-	ND
298	Peri-urban	12	Mucoid	Salmonella Zanzibar	3,10:k:1,5	ND
302	Municipal	24	Watery	Salmonella Enteritidis	9,12:g,m:-	1
326	Municipal	18	Mucoid	Salmonella Zanzibar	3,10:k:1,5	ND

Table 2. Demographic and phenotypic characteristics of *Salmonella* serotypes isolated from diarrheic children younger than five years of age in FCT, Abuja, Nigeria

ND: not determined

Table 3. The PFGE patterns of *Xba*I and *Bln*I digested chromosomal DNA of *Salmonella* serovar isolates from diarrheagenic children younger than five years of age in FCT Abuja, Nigeria

Salmon alla conovon	PFGE	pattern	Information		
Saimoneua serovar	XbaI	BlnI	interence		
S. Zanzibar (5)	ZanXAI.0001	ZanBNI.0001	New XbaI, new BlnI		
	ZanXAI.0001	ZanBNI.0002	New BlnI		
	ZanXAI.0001	ZanBNI.0001	nil		
	ZanXAI.0001	ZanBNI.0001	nil		
	ZanXAI.0001	ZanBNI.0001	nil		
S. Brancaster(3)	BranXAI.0001	BranBNI.0001	nil		
	BranXAI.0001	BranBNI.0001	New XbaI, new BlnI		
	BranXAI.0001	BranBNI.0001	nil		
S. Enteritidis(1)	SENXAI.0001	SENBNI.0017	Common patterns		

Table 4a. Antimicrobial resistance pattern of *Salmonella* isolates (n = 9)

P		
Antibiotic used	Number of resistant strains (%)	
Amoxicillin	5 (55.5)	
Cephalexin	5 (55.5)	
Cefuroxime	5 (55.5)	
Ciprofloxacin	0 (0.0)	
Ceftriaxone	0 (0.0)	
Amoxicillin-clavulanic acid	2 (22.2)	
Nalidixic acid	0 (0.0)	

Table 4b. Antibiotic resistance pattern of nine strains of Salmonella serovars isolated	from children younger than five years
of age in FCT, Abuja Nigeria.	

Salmonella serovars	Antibiotic resistance profile	No of Strain	Total No
Salmonella Zanzibar (5)	AMX,CEPH	1	4
	CEPH ,CEFU	1	
	AMX,CEPH,CEFU	1	
	AMX,AUG,CEPH,CEFU	1	
Salmonella Brancaster(3)	AMX,CEPH,CEFU	1	2
	AMX,AUG,CEPH,CEFU	1	
Salmonella Enteritidis(1)	AMX,CEPH,CEFU	1	1

AMX: amoxicillin; CEPH: cephalexin; CEFU: cefuroxime; CEFTR: ceftriaxone; AUG: Augmentin (amoxicillin-clavulanic acid)

Comparatively, one isolate out the five *S*. Zanzibar isolates had a different *Bln*I restriction pattern than the other four (relating to a single band shift). The three isolates of *S*. Brancaster had the same restriction patterns for *Xba*I and *Bln*I digested chromosomal DNA, respectively. No comparison could be made for serovar Enteritidis clonality based on PFGE, as only a single patient isolate was recovered. The PFGE patterns of the five strains of *S*. Zanzibar isolated appeared identical and hence genetically related, as did the three strains of *S*. Brancaster isolates (Table 3).

The following levels of resistance were found among the *Salmonella* strains: amoxicillin, five strains (55.6%); augmentin, two strains (22.2%); cephalexin, five strains (55.6%); and cefuroxime, five strains (55.6%). Intermediate resistance was found in five strains (55.6%) only to amoxicillin-clavulanic acid (Augmentin). All the serotypes isolated were susceptible to nalidixic acid, ciprofloxacin, and ceftriaxone (Table 4a). In all, seven antimicrobial resistant phenotypes were observed among the three *Salmonella* serovars. All the *Salmonella* serovars with the exception of one *S*. Zanzibar strain were resistant to two or more antibiotics. Pan susceptibility was detected in only one strain each of *S*. Zanzibar and *S*. Brancaster. Similarly, a multidrug-resistant pattern to

Dice (Opt:0.50%) (Tol 1.0%-1.0%) (H>0.0% S>0.0%) [0.0%-100.0%]

four antibiotics (AMX, AUG, CEPH, and CEFU) was detected in only one strain each of these serovars (S. Zanzibar and S. Brancaster). Among the three Salmonella serovars, common resistant profile patterns to three antibiotics (AMX, CEPH, and CEFU) were observed (Table 4b). Although among the Salmonella serovars, multidrug resistance was observed in five (55.6%) of the Salmonella serovars (two S. Zanzibar, two S. Brancaster) and in the only S. Enteritidis serovar, none exhibited resistance MICs to tested antimicrobial agents according to the current CLSI breakpoints. In addition, all the serotypes exhibited susceptible zone inhibition diameters ranging from 32-34 mm (minimum and maximum) to ceftazidime, cefotaxime, and cefpodoxime alone or in combination with clavulanic acid. No ESBL-producing Salmonella isolates identified as either TEM or CTX-M-type ESBLs genes were detected in any of the isolates. Screening of isolates for *bla*_{SHV} and AmpC betalactamase production was not performed in this study.

Figure 1 illustrates phylogenic dendrogram of PFGE patterns obtained using the restriction enzymes *XbaI* and *BlnI* showing percent similarity calculated by the Dice similarity index of PFGE restriction endonuclease digestion profiles among the nine Salmonella isolates (*S.* Zanzibar, *S.* Brancaster, *S.*

Figure 1. Dendogram showing the cluster analysis on the basis of *Xba*I-PFGE and *BIn*I-PFGE of the three *Salmonella* serovars. ND: Not determined;NHA: National Hospital Abuja; MDH: Maitama Distict Hospital; NGH: Nyanya General Hospital

PFGE Xbal	PFGE X	bal												
-1 00 100	-1000	600.00	-400.00 -300.00 -250.00	-150.00	40.00	20.00	Reference	Туре	Phage Type	Antigen	Date (Mo/Yr)	Pat. Age (months)	Diarrhea Type	Hosp-Loc
	111						87	Zanzibar	ND	ND	Aug/08	4	Watery	MDH
	111						155	Zanzibar	ND	ND	Sept/08	12	Mucoid	NHA
	111						159	Zanzibar	ND	ND	Sept/08	24	Mucoid	NHA
	III		III		Ī		298	Zanzibar	ND	ND	Sept/08	12	Mucoid	NGH
			11				326	Zanzibar	ND	ND	Sept/08	18	Mucoid	MDH
	11			L I			302	Enteritidis	1	9,12:g,m	Sept/08	24	Watery	NHA
	I I					1	99	Brancaster	ND	ND	Aug/08	12	Watery	NGH
	11						167	Brancaster	ND	ND	Sept/08	18	Watery	NHA
1	11		11	<u> </u>		Ì	280	Brancaster	ND	ND	Sept/08	4	Watery	NHA
Dice (Opt0.50%) (Tol 1.0%-1.0%) (PFGE Bini	H>0.0% S>0.0 PFGE BI	%) [0.0% I ni	6-100.0%]											
- 40 - 80 - 100	1000	500.00	350.00 -300.00 -250.00	-200.00 -150.00 -100.00	-40.00	-20.00	Reference	Туре	Phage Type	Antigen	Date (Mo/Yr)	Pat. Age (months)	Diarrhea Type	Hosp-Loc
	1 1	1	H	. 1			99	Brancaster	ND	ND	Aug/08	12	Watery	NGH
	1 1	1	11.	1			167	Brancaster	ND	ND	Sept/08	18	Watery	NHA
		1	1				280	Brancaster	ND	ND	Sept/08	4	Watery	NHA
	1 11	I	111	Í	1		302	Enteritidis	1	9,12:g,m	Sept/08	24	Watery	NHA
	-	1	11	11			87	Zanzibar	ND	ND	Aug/08	4	Watery	MDH
	111	1	11	11			159	Zanzibar	ND	ND	Sept/08	24	Mucoid	NHA
	11	1	11	11			298	Zanzibar	ND	ND	Sept/08	12	Mucoid	NGH
		1					326	Zanzibar	ND	ND	Sept/08	18	Mucoid	MDH
		1			1		155	Zanzibar	ND	ND	Sept/08	12	Mucoid	NHA
ND: Not Determ	nined	1	NHA : I	Nationa	al Ho	spi	tal Abuja	MDH: N	/laitama Di	strict Ho	spital	NGH: Nyanya (General Hosp	oital

Enteritidis). The different patterns, sources, year of isolation, and number of strains are indicated. The Zanzibar and Brancaster isolates were the first from north-central Nigeria to be ever typed using PFGE. The five S. Zanzibar isolates all hade the same restriction patterns using XbaI (no genetic diversity was identified), but one isolate had a different BlnI restriction pattern than the other four. Four Zanzibar isolates were from mucoid-type diarrhea. All three S. Brancaster isolates assigned to the same PFGE pattern had the same XbaI and BlnI patterns. Of the three Brancaster isolates, two isolates were from the municipal area and one was from the peri-urban area. No comparative assessment of S. Enteritidis was possible, but the lone isolate was recovered from Abuja Municipality and was phenotypically phage type 1.

Discussion

This study characterized Salmonella species recovered from children with acute gastroenteritis, and determined antibiotic susceptibility patterns and extended-spectrum beta-lactamase production using polymerase chain reaction, pulse field gel electrophoresis, micro dilution, and double disk diffusion techniques. Notably, a rare serotype incriminated in human Salmonella infections, S. Zanzibar, had the highest prevalence (55.6%) in this study. The first reported human infection with the serotype S. Zanzibar was in 1976 from the stool culture of an eight-year-old boy with gastroenteritis in Ankara [19]. Thereafter, it has been documented to be associated with adult infections in North Scotland [20], and more recently in Tunis [21]. Notably, multidrug resistant S. Brancaster isolated from poultry has been reported in Dakar, Senegal [22]. The reported epidemiology of S. Brancaster in the literature is rather limited [23]. However, its occurrence in our study is in consonance with reported isolation from human cases in the Toscana region of Italy [24], in the United States [25], and in Senegal, on the West African coast [24]. Recently, resistance to spectinomycin and streptomycin by S. Zanzibar isolated from chicken muscle was observed in Dakar, Senegal [26].

The choice of antimicrobial agents evaluated in this study was based on reported resistance of *Salmonella* isolates to commonly used antimicrobials [27]. The results of this study showed that 77.8% of the *Salmonella* isolates tested for antimicrobial susceptibility were resistant to two or more antibiotics. Although a previous study in Nigeria had reported a resistance rate of 61.0% [28], other authors from Ethiopia [29] recently reported a resistance rate of 83.3%, which is, comparatively, a higher proportion of multidrug resistance. The finding of good antimicrobial activity for ciprofloxacin is in line with earlier reports from Lagos, Nigeria [28].

The identification of no ESBL-producing Salmonella serovars is consistent with the findings of some previous studies in developed and developing countries. No ESBLs were detected among human Salmonella isolates in multicenter surveys conducted by a France hospital-based network in 1994 (n=2,622) and 1997 (n=2,464) [30]. Similarly, in China, no ESBL-producing isolates were identified among human diarrheal Salmonella isolates from 28 hospitals in Henan Province in 2006 [31]. More recently, non-ESBL-producing Salmonella isolates were identified in another multicenter study conducted in India that monitored in vitro susceptibility of community- or hospital-acquired Gram-negative bacilli in intraabdominal infections to identify early changes in susceptibility patterns, with a focus on ESBL producers. [32]. Therefore, the antibiotic resistance exhibited by the Salmonella serotypes isolated as observed from the disk diffusion susceptibility tests may have been conferred by a mechanism for β -lactam resistance other than those analyzed in this study. Conversely, the identification of non-ESBL-producing Salmonella isolates may be inadvertent, but not altogether unusual.

Conclusions

Molecular characterization helps scientists to understand the nature of the epidemiology of infections caused the different *Salmonella* serotypes. Our findings have demonstrated the involvement of three *Salmonella* strains in acute gastroenteritis; ciprofloxacin, ceftriaxone, and nalidixic acid remain the antibiotics of choice in the management of acute gastroenteritis caused by the three strains of *Salmonella* isolates. The study further demonstrated the absence of extended-spectrum beta-lactamase producing *Salmonellae*.

Acknowledgements

The authors thank all those who helped and collaborated for the success of this work. We thank the staff of the Enteric Diseases Program (WHO Collaborating Centre) at the National Microbiology Laboratory (Canada) for assisting with serotyping, phage typing, and PFGE testing, including Dr. Celine Nadon, Rafiq Ahmed, Helen Tabor, and Dr. Matthew Gilmour. We are grateful to the children and their parents for kindly providing the stool samples. We thank the laboratory staff of all the hospitals and the center for their contribution in collecting the samples for study, and the staff of the Department of Biological Science, University of Abuja, for the logistic support. We also thank Laboratoire de Contrôle des Eaux et Denrées Alimentaires de l'Institut Pasteur de Tunis in Tunis, Tunisia for their valuable input into this study.

References

- Rodulfo H, De DonatoM, Luiggi J, Michelli E, Millán A, Michelli M (2012) Molecular characterization of *Salmonella* strains in individuals with acute diarrhea syndrome in the State of Sucre, Venezuela. Rev Soc Bras Med Trop 45: 329-333.
- 2. Kosek M, Bern C, Guerrant RL (2003) The magnitude of the global burden of diarrhoeal disease from studies published 1992-2000. Bull World Health Organ 81: 197-204.
- Caniça M, Dias R, Correia JD, Pomba C (2004) Genetic relatedness between human and animal polymorphic *bla*_{TEM} genes strengthens zoonotic potential among uropathogenic *Escherichia coli* strains. J Antimicrob Chemother 54: 284-286.
- Bugarel M, Granier SA, Weill F, Fach P, Brisabois A (2011) A multiplex real-time PCR assay targeting virulence and resistance genes in Salmonella enteric serotype Typhimurium. BMC Microbiol 11: 151.
- Lerouge I, Vanderleyden J (2002) O-antigen structural variation: mechanisms and possible roles in animal/plantmicrobe interactions. FEMS Microbiol Rev 26: 17-47.
- Popoff MY, Bockemuhl J, Gheesling LL (2004) Supplement 2002 (no.46) to the Kauffmann-White scheme. Res Microbiol 155: 568-570.
- 7. Jacoby GA (2006) Beta-lactamase nomenclature. Antimicrob Agents Chemother 50: 1123-1129.
- Akinyemi KO, Bamiro BS, Coker AO (2007) Salmonellosis in Lagos, Nigeria: Incidence of *Plasmodium falciparum*associated Co-infection, Patterns of Antimicrobial Resistance, and Emergence of Reduced Susceptibility to Fluoroquinolones. J Health Popul Nutr 25: 351-358.
- Tjaniadi P, Lesmana M, Subekti D, Machpud N, Komalarini S, Santoso W (2003) Antimicrobial resistance of bacterial pathogens associated with diarrheal patients in Indonesia. Am J Trop Med Hyg 68: 666-670.
- Yates C, Amyes S (2005) Extended-spectrum beta-lactamases in non-typhoidal *Salmonella* spp. isolated in the UK are now a reality: why the late arrival? J Antimicrob Chemother 56: 262-264.
- AitMhand R, Soukri A, Moustaoui N, Amarouch H, ElMdaghri N, Sirot D, Benbachir M (2002) Plasmid-mediated TEM-3 extended-spectrum beta-lactamase production in Salmonella typhimurium in Casablanca.J Antimicrob Chemother49: 169-172.
- Winn W, Allen S, Janda W, Koneman E, Procop G, Schreckenberger P (2005) Koneman's Color Atlas and Textbook of Diagnostic Microbiology, 6th edition. New York: Lippincott Williams & Wilkins.
- MacFaddin J (2000) Biochemical Tests for Identification of Medical Bacteria. 3rd edition. New York, USA: Lippincott Williams & Wilkins.
- Bopp CA, Brenner FW, Fields PI, Wells JG, Strockbine NA (2003) *Esherichia*, *Shigella*, and *Salmonella*. In. Murray PR,

Baron EJ, Jorgensen JA, Pfaller MA, Yoken RH, editors. Manual of Clinical Microbiology, 8th edition. 654-671.

- Oliveira S, Santos L, Schuch D, Silva A, Sallr S, Canal C (2002) Detection and identification of Salmonella from poutry-related samples by PCR. Vet Microbiol 87: 25-35.
- Swaminathan B, Barrett TJ, Hunter SB (2001)PulseNet: the molecular subtyping network for foodborne bacterial disease surveillance, United States. Emerg Infect Dis 7: 382-389.
- Cui S, Li J, Sun Z, Hu C, Jin S, Li F, Guo Y, Ran L, Ma Y (2009) Characterization of Salmonella enterica isolates from infants and toddlers in Wuhan, China. J Antimicrob Chemother 63: 87-94.
- Clinical and Laboratory Standards Institute (2006) Performance standards for antimicrobial susceptibility testing, 16th informational supplement. Document M100-S16. Wayne, PA: CLSI.
- Council for International Organization of Medical Sciences (2002) International Ethical Guidelines for Biomedical Research Involving Human Subjects. Geneva, Switzerland: CIOMS.
- Aksoycan N (1976) First isolation of S. Zanzibar in Turkey. Mikrobiyol Bulletin 10: 65-66.
- 21. Johnston WS (1981) An unusual sequel to imported *Salmonella* Zanzibar.J Hyg Cambri 87:525-528.
- Bada-Alambedji R, Fofana A, Seydi M, Akakpo AJ (2006) Salmonella Isolated from Poultry Carcasses in Dakar (Senegal). Braz J Microbiol 37: 510-515.
- 23. AL-Gallas N, Bahri O, Bouratbeen A, Ben Hasen A, Ben Aissa R (2007) Etiology of acute diarrhea in children and adult in Tunis-Tunisia with emphasis on diarrheagenic *E.coli*: prevalence, phenotyping, and molecular epidemiology. Am J Trop Med Hyg77: 571-582.
- 24. Cardinale E, Perrier Gros-Claude JD, Rivoal K, Rose V, Tall F, Mead GC, Salvat G (2005) Epidemiological analysis of *Salmonella enterica* ssp. *enterica* serovars Hadar, Brancaster and Enteritidis from humans and broiler chickens in Senegal using pulsed-field gel electrophoresis and antibiotic susceptibility. J Appl Microbiol99: 968-977.
- Fontin G, Sanavio G,Maini C, Giugliano I,Polli AC (1993) Isolamento di salmonelle nel triennio 1989–1991 da acque superficiali, coprocolture ed alimenti. Boll Microbiol Indagini Lab13: 15-24.
- Dione MM, Geerts S, Antonio M (2012) Characterisation of novel strains of multiply antibiotic-resistant *Salmonella* recovered from poultry in Southern Senegal.J Infect Dev Ctries 6: 436-442. doi:10.3855/jidc.1530.
- 27. Pacini R, Quagli E, Galassi R, Tozzi E,Malloggi L (1994) Dieci anni di sorveglianza epidemiologica a Livorno (I parte) Studio epidemiologico sulle salmonelle isolate da portatori sani, nell'area livornese, dal 1984 al 1993. Boll Microbiol Indagini Lab14: 9-14.
- Akinyemi KO, Bamiro BS, Coker AO (2007) Salmonellosis in Lagos, Nigeria: Incidence of *Plasmodium falciparum*associated Co-infection, Patterns of Antimicrobial Resistance, and Emergence of Reduced Susceptibility to Fluoroquinolones. J Health Popul Nutr 25: 351-358.
- Zewdu E, Cornelius P (2009) Antimicrobial resistance of Salmonella serotypes isolated from food items and personnel in Addis Ababa, Ethiopia. Trop Animal Health Prod 41: 241-249.
- Breuil J, Brisabois A, Casin L, Armand-Lefèvre L, Frémy S, Collatz E(2000) Antibiotic resistance in salmonellae isolated

from humans and animals in France: comparative data from 1994 and 1997. J Antimicrob Chemother 46: 965-971.

- 31. Zhu M, Xie Z, Zhang L, Xia S, Lu W, Ran Y, Wang Z (2009) Characterization of *Salmonella Enterica* Serotype Typhimurium from Outpatients of 28 Hospitals in Henan Province in 2006.Biomed and Environ Sci 22: 136-140.
- Chaudhuri BN, Rodrigues, C, Balaji V, Iyer R, Sekar U, Chand WD, Chitnis DS, Dhole TN, Joshi S (2011) Incidence of ESBL Producers amongst Gram-negative Bacilli isolated from intra-abdominal infections across India (Based on SMART study, 2007 Data). J Assoc Phys India 59: 287-292.

Corresponding author

Casmir Ifeanyichukwu Cajetan Ifeanyi Department of Veterinary Microbiology, Faculty of Veterinary Medicine University of Abuja P.M.B 117, Abuja, Nigeria Phone: +2348030750990 Email: ifeanyi.casmir@gmail.com

Conflict of interests: No conflict of interests is declared.