



Virome analysis of two sympatric bat species (*Desmodus rotundus* and *Molossus molossus*) in French Guiana

Vincent Lacoste, Arielle Salmier, Sourakhata Tirera, Alain Franc, Edith Darcissac, Damien Donato, Christiane Bouchier, Anne Lavergne, Benoit de Thoisy, Naomi Forrester

► To cite this version:

Vincent Lacoste, Arielle Salmier, Sourakhata Tirera, Alain Franc, Edith Darcissac, et al.. Virome analysis of two sympatric bat species (*Desmodus rotundus* and *Molossus molossus*) in French Guiana. PLoS ONE, 2017, 12 (11), pp.1-25. 10.1371/journal.pone.0186943 . pasteur-01633803

HAL Id: pasteur-01633803

<https://riip.hal.science/pasteur-01633803>

Submitted on 13 Nov 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License


RESEARCH ARTICLE

Virome analysis of two sympatric bat species (*Desmodus rotundus* and *Molossus molossus*) in French Guiana

Arielle Salmier¹, Sourakhata Tirera¹, Benoit de Thoisy¹, Alain Franc², Edith Darcissac¹, Damien Donato¹, Christiane Bouchier³, Vincent Lacoste¹, Anne Lavergne^{1*}

1 Laboratoire des Interactions Virus-Hôtes, Institut Pasteur de la Guyane, Cayenne, French Guiana, **2** UMR BIOGECO, Institut National de la Recherche Agronomique (INRA), Cestas, France, **3** Institut Pasteur, Plateforme de Génomique - Pôle Biomix, Paris, France

* alavergne@pasteur-cayenne.fr



Abstract

Environmental disturbances in the Neotropics (e.g., deforestation, agriculture intensification, urbanization) contribute to an increasing risk of cross-species transmission of microorganisms and to disease outbreaks due to changing ecosystems of reservoir hosts. Although Amazonia encompasses the greatest diversity of reservoir species, the outsized viral population diversity (virome) has yet to be investigated. Here, through a metagenomic approach, we identified 10,991 viral sequences in the saliva and feces of two bat species, *Desmodus rotundus* (hematophagous), trapped in two different caves surrounded by primary lowland forest, and *Molossus molossus* (insectivorous), trapped in forest and urban habitats. These sequences are related to 51 viral families known to infect a wide range of hosts (*i.e.*, bacteria, plants, insects and vertebrates). Most viruses detected reflected the diet of bat species, with a high proportion of plant and insect-related viral families for *M. molossus* and a high proportion of vertebrate-related viral families for *D. rotundus*, highlighting its influence in shaping the viral diversity of bats. Lastly, we reconstructed the phylogenetic relationships for five vertebrate-related viral families (*Nairoviridae*, *Circoviridae*, *Retroviridae*, *Herpesviridae*, *Papillomaviridae*). The results showed highly supported clustering with other viral sequences of the same viral family hosted by other bat species, highlighting the potential association of viral diversity with the host's diet. These findings provide significant insight into viral bat diversity in French Guiana belonging to the Amazonian biome and emphasize that habitats and the host's dietary ecology may drive the viral diversity in the bat communities investigated.

OPEN ACCESS

Citation: Salmier A, Tirera S, de Thoisy B, Franc A, Darcissac E, Donato D, et al. (2017) Virome analysis of two sympatric bat species (*Desmodus rotundus* and *Molossus molossus*) in French Guiana. PLoS ONE 12(11): e0186943. <https://doi.org/10.1371/journal.pone.0186943>

Editor: Naomi Forrester, University of Texas Medical Branch at Galveston, UNITED STATES

Received: February 7, 2017

Accepted: October 10, 2017

Published: November 8, 2017

Copyright: © 2017 Salmier et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files. All virus sequences reported in this study have been deposited in the GenBank nucleotide database under accession numbers KX812440 to KX812444, KX812446, KX812447, KX821677 and KX954092. The data from Illumina sequencing were deposited in GenBank Sequence Reads Archive under accession numbers SAMN05725475–SAMN05725481.

Introduction

Bats present a rich viral diversity as compared to other animals and their uniqueness as hosts of pathogenic viruses has animated debates in recent years [1–3]. Their ability to fly over long distances and their diverse feeding strategies seem to facilitate the acquisition and dispersal of

Funding: A. Salmier was funded by a grant from European funds (PO FSE 2007-2013) and "Investissement d'Avenir" managed by Agence Nationale de la Recherche (CEBA, Ref. ANR-10-LABEX-25-01). This study was conducted within the CAROLIA program supported by European funds (ERDF/FEDER) and assistance from Région Guyane and Direction Régionale pour la Recherche et la Technologie. It also received a European Commission "REGPOT-CT-2011-285837-STRonGer" grant within the FP7 and "Investissement d'Avenir" grants managed by the Agence Nationale de la Recherche (CEBA, Ref. ANR-10-LABEX-25-01). The Genomics Platform is a member of the France Génomique consortium (ANR10-INBS-09-08). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

viruses across remote regions, as well as cross-species transmissions [2,4,5]. Likewise, their wide variety of both social structures and behaviors contribute to viral transmissions and persistence of viruses in bat populations [2,6]. Studies have also highlighted that bats present long associations with numerous viral families and genera (e.g., *Paramyxoviridae*, *Filoviridae*, *Lyssavirus*, *Henipavirus*), with viruses detected in bats usually being older than those found in humans or other animals [7]. To date, viral metagenomics in bats has almost exclusively focused on North American and Eurasian bat communities and many novel mammalian viruses have been reported, including some that are important to public health [8–17].

Amazonia encompasses a great diversity of bat species. Numerous studies, using conventional methods or high-throughput sequencing, have allowed the identification of a number of viruses in this region [18–21]. For instance, Drexler *et al.* detected and identified a diversity of paramyxoviruses in several New World bat species [22], Tong *et al.* reported novel influenza A viruses in Peruvian flat-faced fruit bats (*Artibeus planirostris*) [23], and Lima *et al.* proposed a new genus for two newly characterized polyomaviruses in *Tadarida brasiliensis* [24].

In French Guiana, an Amazonian region, 106 bat species have been listed [25]. During the last few decades this region has undergone environmental changes (e.g., urbanization, agriculture intensification, deforestation), which have led to alterations in the composition and dynamics of bat communities [26,27]. Studies conducted in the Neotropics reported an overall decrease in species richness and relative abundance correlated with the urban influence, although species highly tolerant to environmental disturbances, such as insectivorous bats, tend to persist in large urban environments [28]. Furthermore, the diversity of bat habitats may influence both microbe transmission and persistence in bat communities. For example, de Thoisy *et al.* [29] showed that pristine forest habitats favored the circulation and maintenance of the rabies virus, compared to disturbed areas, and in Southeast Asian rainforests Gay *et al.* [30] showed that fragmentation reduces both viral and endoparasite species richness.

The aim of our study was to characterize the virome in the feces and the saliva of two abundant and opportunistic Amazonian bat species: the common vampire bat, *Desmodus rotundus* (Phyllostomidae) and Pallas's mastiff bat, *Molossus molossus* (Molossidae). Both bat species are sympatric in the forest areas of French Guiana. They both have high plasticity and tolerance to perturbations but present distinct diet and social behavior.

D. rotundus bats are hematophagous and live in groups in highly complex social systems, for which Wilkinson *et al.* showed a strong association between the mating system and genetic structure of populations [31,32]. Studies on *D. rotundus* also highlighted that their diet associated with social interactions (especially social grooming and food sharing) within colonies contribute to the transmission and maintenance of viruses such as rabies [29,33]. Furthermore, a modeling study of rabies transmission in a *Desmodus* population in Peru identified metapopulation dynamics as a major driver of rabies maintenance [34]. Turmelle and Olival also pointed out that genetic spatial structure and the host's ecology and social behavior correlate with viral richness [35]. We investigated the virome of *D. rotundus* bats trapped in two caves (F and M) located in forest areas, for which no exchange of individuals was observed through a longitudinal mark-recapture study [29].

M. molossus bats are insectivorous and live in stable social groups that practice social foraging over large spatial scales [36]. Studies on Molossidae also highlighted that members of this family tend to benefit from increased foraging resources in urban areas and persist in large urban environments for which they present high behavioral flexibility and heterogeneity [28]. We investigated the virome of *M. molossus* bats trapped in urban and forest areas.

Lastly, focusing on viral families known to infect mammals, we investigated the phylogenetic relationships of viral sequences related to five viral families (*Circoviridae*, *Herpesviridae*, *Nairoviridae*, *Papillomaviridae* and *Retroviridae* (genus *Spumavirus*)), detected in either high

proportion in both species and all habitats or only a distinct bat species and habitat. These findings allowed identifying novel vertebrate-related viral sequences and emphasize the importance of studying the role of habitats and dietary ecology in shaping viral diversity for these two Amazonian bat species.

Materials and methods

Ethics statement

All animals were captured, handled and sampled following ASM guidelines [37] under the supervision of researchers granted the French animal experimentation level 1 diploma. Bats are not protected by law in French Guiana. The project was nevertheless submitted to and approved by the Conseil Scientifique Régional pour le Patrimoine Naturel de la Guyane. Captures that occurred within protected areas (nature reserves) received approval by the Conseil Scientifique Régional du Patrimoine Naturel on 26 January 2010 and ad hoc authorizations (No. 2011–35 dated 05/30/2011, No. 35 and 59 obtained 03/21/2013 and 04/17/2013, respectively, and delivered by the Prefecture of French Guiana).

Study areas

D. rotundus bats were sampled in two caves (F and M) located 25 km apart in pristine primary lowland forests, presenting differences in physical patterns, cave-inhabiting bat species as well as population size (Table 1 and S1 Fig). Cave F is cohabited by four bat species (i.e., *Anoura geoffroyi*, *Carollia perspicillata* and *Trachops cirrhosus*), and the *D. rotundus* population ranges from 60 to 100 individuals [29]. Cave M is larger and moister than cave F, suggesting different roost characteristics and carrying capacity. Cave M is cohabited by at least nine different bat species (i.e., *Anoura geoffroyi*, *Artibeus* spp., *Carollia perspicillata*, *Pteronotus rubiginosus* and *P. sp3*, *Tonatia saurophila*, *Trachops cirrhosus* and *Xophostoma sylvicola*), and the *D. rotundus* population ranges from 120 to 150 individuals [29].

M. molossus bats were sampled in two sites located in primary lowland rainforests and under the roofs of houses in two sites located in disturbed areas (Table 1 and S1 Fig). *M. molossus* individuals trapped in forest habitats formed monospecific colonies, whereas individuals trapped in urban habitats shared artificial roosts with a congener, *Molossus coibensis*.

Bat sampling

Feces and saliva samples were collected in both dry and rainy seasons during a 2-year period (April 2012 to April 2014). All animals were trapped with mist nets erected inside roosts or in

Table 1. Characteristics of the sampling sites and number of collected feces and saliva samples for *Molossus molossus* and *Desmodus rotundus*.

Species	Habitats	Sites	GPS coordinates		n saliva swabs		n feces	
			Latitude (N)	Longitude (W)	Per site	Total	Per site	Total
<i>D. rotundus</i>	Forest	Cave F	4° 38' 59.144"	52° 17' 35.441"	0	0	75	75
		Cave M	4° 32' 6.82"	52° 9' 6.239"	50	50	66	66
					Total	50	Total	141
<i>M. molossus</i>	Forest	Paracou	5° 14' 21.282"	52° 55' 24.218"	17	58	4	14
		Saut Athanase	4° 11' 14.706"	52° 19' 11.21"	41		10	
	Urban	Cacao	4° 34' 33.771"	52° 28' 4.926"	18	30	0	5
		La Chaumière	4° 53' 5.419"	52° 21' 4.124"	12		5	
					Total	88	Total	19

<https://doi.org/10.1371/journal.pone.0186943.t001>

putative foraging courses. Species were identified on site, using external morphology. Prior to release, 138 saliva swabs were collected from *M. molossus* and *D. rotundus*, in cave M only (Table 1). Samples were preserved on ice with 500 μ l of Dulbecco's modified Eagle Medium (DMEM, Sigma).

Feces were collected in two ways: *M. molossus* bats were held in individual sacks secured to a rope line for at least 10 min, allowing enough time for the excretion of 19 fresh feces of about 100 mg each ($n = 5$ in urban habitats, $n = 14$ in forest habitats; Table 1), while clean plastic sheets were laid down on flat surfaces beneath known roosts for *D. rotundus*. Then 141 freshly produced feces of about 150 mg each were collected ($n = 75$ in cave F, $n = 66$ in cave M; Table 1) the following morning and temporarily preserved on ice. All samples were later stored at -80°C .

Sample processing

Feces and saliva samples were pooled according to the habitats and the species regardless of the season and year, corresponding to four pools for *M. molossus* (urban saliva, urban feces, forest saliva, forest feces) and three pools for *D. rotundus* (cave F feces, cave M saliva, cave M feces) (Table 1).

Samples were processed as previously described by Victoria *et al.* [38]. Briefly, feces were vigorously homogenized with 5–10 mL DMEM. Samples were cleared of debris by low-speed centrifugation (5 min, 10,000 g, 4°C). Eukaryotic and prokaryotic cell-sized particles were removed from supernatants through three successive filtrations (0.8 μm , 0.45 μm and 0.22 μm), using cellulose acetate membrane filters (Nalgene). A differential centrifugation procedure, described by Prescott *et al.* [39], was used to pellet the viral particles. Briefly, filtrates were cleared of smaller and less dense components through a 1-h ultracentrifugation (100,000 g, 4°C) procedure. Then the pellets were resuspended in nuclease-free water and cleared of persistent high-density particles with a low-speed centrifugation (15 min, 10,000 g, 4°C). Lastly, viral particles were pelleted with a 1-h ultracentrifugation step (100,000 g, 4°C).

For saliva samples, swabs were vigorously resuspended and 200 μL of suspension from each collecting tube was used to constitute the pools. Samples were cleared of debris by low-speed centrifugation (5 min, 10,000 g, 4°C). Eukaryotic and prokaryotic cell-sized particles were removed from supernatants through two successive filtrations (0.45 μm and 0.22 μm), using cellulose acetate membrane filters (Nalgene). The filtrates were cleared of persistent high-density particles with low-speed centrifugation (15 min, 10,000 g, 4°C), then viral particles were pelleted with a 1-h ultracentrifugation step (100,000 g, 4°C). All viral pellets were resuspended in 40 μL of nuclease-free water.

Nuclease treatment and viral acid nucleic extraction

Resuspended viral pellets from feces were treated with a mixture of DNases (Turbo DNase from Ambion and Benzonase from Novagen) and RNase One (Promega) to digest nonenveloped nucleic acids (*i.e.*, those not in viral capsids) [40]. Resuspended viral pellets from saliva samples were treated with the mixture of DNases. All viral nucleic acids (RNA and DNA) were then extracted using the NucliSENS easyMAG[®] bio-robot (bioMérieux).

Reverse transcription and amplification

For both feces and saliva samples, the RNA virus-only and DNA virus-only libraries were respectively constructed using a whole transcriptome (WTA) or a whole genome (WGA) amplification method previously described by Berthet *et al.* [41].

For the RNA virus-only amplification, an aliquot of the extracted viral nucleic acid collected was treated with Turbo DNase to remove viral DNA. Persistent rRNA was depleted with the GeneRead™ rRNA depletion kit (Qiagen), following the manufacturer's recommendations. Then viral RNA amplification was performed as described in the protocol of the QuantiTect® Whole Transcriptome Kit (Qiagen) except for the reverse transcription step. cDNA was synthesized using SuperScript® III Reverse Transcriptase (Invitrogen) and random hexamers (Roche), following the manufacturers' recommendations.

For DNA virus-only amplification, an aliquot of the extracted viral nucleic acid collected was treated with RNase One to remove viral RNA. Then viral DNA amplification was performed as described in the protocol of the QuantiTect® Whole Genome Kit (Qiagen). To ensure homogeneity and blunt ends at the end of the strands produced, 10 U of Klenow polymerase (Roche) was added to the WTA- and WGA-amplified nucleic acids with 8 µL of random hexamers (Roche), incubated 1 h at 37°C, followed by 10 min at 75°C. Samples were assayed with a Qubit® fluorometric quantitation (Qiagen), using the dsDNA Broad Range and dsDNA High Sensitivity Qubit® assay kits, as recommended by the manufacturer.

High-throughput sequencing

For both feces and saliva samples, 1 µg of each viral library was pooled together, whenever possible, to construct RNA plus DNA viral libraries. High-throughput sequencing was carried out at the Genomics platform at the Institut Pasteur, Paris. Shotgun libraries were prepared by standard Illumina protocols using 1 µg of total genomic DNA. Each sample (feces or saliva) was indexed according to its provenance (species and habitats) using Illumina adaptor-specific primers. Samples were sequenced on a MiSeq sequencer in 300-base paired-end reads.

Bioinformatic pipeline

The Illumina sequencing reads were grouped according to their adaptor tags and were processed individually. Sequence files were stripped of their adapter sequences using Trimmomatic v.0.32 [42]. Duplicate sequence reads were removed with FastqMcf v.1.04.676 and reads were quality-filtered using the fastq_quality_filter program from the Fastx toolkit v.0.0.13 (available at http://hannonlab.cshl.edu/fastx_toolkit/index.html), with a quality threshold of 30 and length criteria of 70%. All cleaned data sets were saved for further analysis. Simultaneously, a three-pass digital normalization was run on each sample file using the Khmer software library [43,44]. Normalized files were saved for further analysis. Following Baker's contig consolidation strategy [45], both normalized and non-normalized reads were *de novo* assembled using two assembly algorithms: Velvet [46] and SPAdes [47]. Three k-mer sizes (21, 55 and 99) were used for both assemblers. The contigs generated were compared (Fig 1) between k-mer sizes for each assembler (comparison 1), between assemblers (comparison 2) and between normalized and non-normalized data sets (comparison 3). A removal of duplicated contigs and a clustering, using USEARCH v.7.0.1090 (available at <http://www.drive5.com/usearch>, [48]) with a 95% threshold identity value, were performed following each comparison. Clustered contigs from comparison 3 (centroids) were considered as consolidated contig sets with lengths ranging from 33 to 45,057 bp and low redundancy. Centroids were submitted to BLASTn comparison with both the NCBI nt and Institut Pasteur de Paris gbvrl databases (in March 2015), (Fig 1). The best-matched reference sequence was retrieved from both databases (cutoff value: 10^{-5}). A length comparison between the results was performed using a bit score ratio ($br = \text{bitscore_blastn_nt} / \text{bitscore_blastn_gb_vrl}$), and only contigs with $br > 0.90$ were regarded as suspect-viral sequences and conserved for further analysis. Centroids not identified as suspect-viral underwent a BLASTx comparison against a custom-made database, which




Fig 1. Bioinformatic analysis pipeline for the 3,722,219 assembled contigs, based on Baker *et al.* [45]. (A) Contigs assembled from both normalized and non-normalized sequence reads by two *de novo* assemblers (SPAdes and Velvet) were consolidated by sequential comparisons (numbered shadowed areas) and removal of duplicate sequences (red arrows). (B) Centroids subjected to sequential BLAST comparison and manually controlled taxonomic classification to identify viral sequences. Proportion of sequences assembled from feces (red) and saliva swabs (blue) are shown in the stacked chart for each species

depending on habitats (caves F and M for *D. rotundus* and Urban and Forest for *M. molossus*). (C) Column charts represent the distribution of contig matches (per pooled samples and overall) and the distribution of vertebrate-related viral families by bat species and habitats.

<https://doi.org/10.1371/journal.pone.0186943.g001>

included all protein sequences of the viral kingdom (cutoff value: 10^{-3}). To reduce the mis-identification bias of false-positive sequences, identified suspect-viral sequences underwent a second BLASTx comparison with the NCBI nr database (cutoff value: 10^{-3}) (Fig 1). Sequences not retained were discarded. Taxonomic information and the kingdom of each gene id were retrieved with taxdb.

Phylogenetic analysis

Contigs from five vertebrate-related viral families: *Nairoviridae*, *Circoviridae*, *Retroviridae*, *Herpesviridae* and *Papillomaviridae* were used for phylogenetic analysis. Reference genomes or nucleotide sequences of previously identified viruses were downloaded from GenBank. The accession number of viral sequences used to infer the phylogenetic trees are given in the respective analyses. Nucleotide and protein sequence editing were performed with GENEIOUS R9 (available at <http://www.geneious.com>, [49]). Sequences were aligned using the MAFFT alignment tool [50] included in the software. Nucleotide and protein sequences were trimmed and gap-stripped prior to phylogenetic analyses. For each analysis, the best-fitted model of nucleotide or amino acid substitution was selected using jModelTest 2 [51] and ProtTest 3 [52], respectively, under corrected Akaike information criteria (AICc). Bayesian phylogenetic analyses were performed using MrBAYES 3. The Markov chain Monte Carlo (MCMC) algorithm was run with four chains with 10 million generations each, with trees sampled every 500 generations and a burn-in of 25%. Validation of the inference was assessed based on the standard deviation of split frequencies, less than the expected threshold value of 0.01 in MrBAYES and by inspecting the effective sampling size (ESS > 500) criterion in TRACER version 1.6 [53].

Nucleotide sequence accession numbers

All virus sequences reported in this study were deposited in the GenBank nucleotide database under accession numbers KX812440 to KX812444, KX812446, KX812447, KX821677 and KX954092. The data from Illumina sequencing were deposited in GenBank Sequence Reads Archive under accession numbers SAMN05725475–SAMN05725481.

Results

Illumina sequencing and assembly

Overall, we obtained 53,325,594 raw read sequences (Table A in S1 File). For *M. molossus*, 33,333,557 raw sequences were generated from both feces and saliva samples and 19,992,037 for *D. rotundus*. Following the data reduction steps, 10,339,752 non-digitally normalized reads and 4,117,544 digitally normalized reads were used for *de novo* assembly. SPAdes and Velvet assemblers generated a comparable number of contigs, even between normalized and non-normalized data. Between 50.22% and 67.66% of the contigs were saved after comparison 1 (Table B in S1 File). Following comparison 2, between 68.09% and 73.19% of the contigs from comparison 1 were saved for further analyses (Table C in S1 File). Finally, between 50.61% and 56.30% of the contigs from comparison 2 were saved to generate the centroids (Table D in S1 File). Following our consolidation steps, a total of 3,722,219 contigs were used for the taxonomic assignment. Most of the contigs retained originated from feces (Fig 1). *Eukaryota*-, *Bacteria*- and *Virus*-associated contigs accounted for 1.02%, 3.82% and 0.30% of the total of

consolidated contigs, respectively (Table E in [S1 File](#)). A total of 10,991 viral-associated sequences related to 51 known viral families were identified based on their most significant BLAST matches ([Fig 2](#)). The total number of virus-associated contigs ranged from 1,930 for *D. rotundus* in cave F to 3,768 for *M. molossus* in forest areas (Table F in [S1 File](#)).

General virome information and phage-related sequences

More than half of the viral-associated contigs matched phage sequences ([Fig 1](#) and [Table 2](#)), accounting for nine phage families belonging to the dsDNA, ssDNA and ssRNA positive-strand groups. These families were identified in both feces and saliva samples ([Fig 2](#) and [Table 2](#)). Among the nine phage families, the *Podoviridae* family of the dsDNA order *Caudovirales* accounted for greatest number of phage sequences detected in *D. rotundus* while the *Microviridae* family accounted for most of the phage sequences identified for *M. molossus* ([Fig 2](#) and [Table 2](#)). The eukaryotic viral sequences (insect, plant/protozoan and vertebrate viruses) accounted for about three-tenths of the total of the viral-associated contigs identified, representing 42 known viral families. A high proportion of these families reflected the diet of bats, especially for *M. molossus* trapped in urban areas where most of the eukaryote-associated viruses identified belonged to nonmammalian viruses ([Fig 1](#) and [Table 2](#)). Contigs matching unclassified viral sequences accounted for one-tenth of the total of viral contigs ([Fig 1](#) and [Table 2](#)).

Eukaryotic viral sequences

Insect viruses. Fourteen insect-related viral families and a group of unclassified viruses were identified ([Table 2](#)). Positive single-stranded RNA (ssRNA (+)) viruses were predominant for *M. molossus* trapped in urban habitats, with most of the viral sequences identified related to the *Dicistroviridae*, *Nodaviridae* and *Iflaviridae* families. DNA viruses detected in this sample were related to viruses from the double-stranded DNA (dsDNA) *Iridoviridae* family and the single-stranded DNA (ssDNA) *Parvoviridae* family. In contrast, ssDNA viruses (mostly from *Parvoviridae*) were slightly more numerous for *M. molossus* trapped in forest habitats compared to RNA viruses mostly from the *Dicistroviridae* and *Nodaviridae* families. The four viral families (*Baculoviridae*, *Iridoviridae*, *Nudiviridae* and *Iflaviridae*) detected for the *D. rotundus* were likely related to insects flying around feces.

Plant and protozoan viruses. Sixteen plant- and protozoan-related viral families and a group of unassigned viruses were identified ([Table 2](#)). ssRNA viruses, mostly from the unassigned group followed by the *Luteoviridae*, *Bromoviridae* and *Virgaviridae* families, were predominant for *M. molossus* trapped in urban habitats. In contrast, dsDNA viruses (*Phycodnaviridae*) and ssDNA viruses (*Geminiviridae*) were predominant for *M. molossus* and *D. rotundus* trapped in forest habitats.

Vertebrate viruses. Fourteen vertebrate-related viral families were identified ([Fig 1](#) and [Table 2](#)). The highest viral diversity was found in both feces samples and forest habitats, and DNA viruses were the most frequently found viruses for both species. The *Anelloviridae*, *Circoviridae* and *Parvoviridae* families accounted for the highest proportion of DNA viruses for both samples of *D. rotundus*, with variation in number according to the sampling sites.

The *Circoviridae* and *Anelloviridae* families were also the most abundant in the sample of *M. molossus* trapped in urban habitats, with very few viral sequences detected in saliva. In contrast, for *M. molossus* trapped in forest habitats, viral sequences related to the *Herpesviridae*, *Papillomaviridae* and *Circoviridae* families were the most numerous, with herpesviruses found for the most part in the saliva sample.




Fig 2. Heatmap based on the viral-associated contigs of 51 families of insect, phage, plant/protozoan and vertebrate viruses in each pooled sample. Location information is provided above each column (Caves F and M for *D. rotundus*, Urban and Forest for *M. molossus*). The names of the viral families are presented in alphabetical order in the middle. The boxes colored from green to dark red represent the number of contigs observed. Contigs varied between 1 and 450 for the saliva samples and between 1 and 1465 for feces. The scales are given for each type of sample.

<https://doi.org/10.1371/journal.pone.0186943.g002>

Table 2. Contingency table of the viral families identified in the feces and saliva samples of *D. rotundus* and *M. molossus*.

Order	Viral family	Genome	Desmodus rotundus						Molossus molossus						Overall total	
			Cave F			Cave M			Urban			Forest				
			Saliva	Feces	Total	Saliva	Feces	Total	Saliva	Feces	Total	Saliva	Feces	Total		
Insect virus	Nidovirales	ssRNA(+)		0	0	0	0	0	0	2	2	0	0	0	2	
	Picornavirales	ssRNA(+)		0	0	0	0	0	0	394	394	0	77	77	471	
	Picornavirales	ssRNA(+)		2	2	0	2	2	0	32	32	0	10	10	46	
	Unassigned	ssRNA(+)		0	0	0	0	0	0	20	20	0	1	1	21	
	Unassigned	dsDNA		0	0	2	1	3	0	1	1	4	0	4	8	
	Unassigned	ssRNA(+)		0	0	0	0	0	0	0	0	0	1	1	1	
	Unassigned	dsDNA		1	1	0	3	3	0	57	57	4	0	4	65	
	Unassigned	dsDNA		0	0	0	0	0	0	0	0	0	4	4	4	
	Unassigned	ssRNA(+)		0	0	0	0	0	0	105	105	0	62	62	167	
	Unassigned	dsDNA		0	0	0	1	1	0	0	0	0	1	1	2	
	Unassigned	ssDNA (+/-)		0	0	0	0	0	0	28	28	17	122	139	167	
	Unassigned	Permutotetraviridae	ssRNA(+)		0	0	0	0	0	2	2	0	0	0	2	2
	Unassigned	Polydnaviridae	dsDNA		0	0	0	0	0	0	0	2	4	6	6	6
	Unassigned	Poxviridae	dsDNA		0	0	0	0	0	0	0	0	3	3	3	3
Phage	Unassigned	-		0	0	0	0	0	0	23	23	0	11	11	34	
	Caudovirales	dsDNA		279	279	1	617	618	2	1	3	19	42	61	961	
	Caudovirales	dsDNA		435	435	0	767	767	0	15	15	8	361	369	1586	
	Caudovirales	dsDNA		285	285	2	173	175	0	5	5	16	56	72	537	
	Ligamaenvirales	dsDNA		1	1	0	0	0	0	0	0	0	0	0	1	
	Unassigned	ssDNA(+)		39	39	0	1	1	0	1	1	0	3	3	44	
	Unassigned	dsDNA		4	4	0	0	0	0	0	0	0	0	0	4	
	Unassigned	ssRNA(+)		0	0	0	0	0	0	3	3	0	0	0	3	
	Unassigned	ssDNA(+)		132	132	0	196	196	0	1402	1402	7	1465	1472	3202	
	Unassigned	dsDNA		7	7	0	3	3	0	0	0	0	0	0	10	
	Unassigned	dsDNA		14	14	0	2	2	0	0	0	0	0	0	16	
	Unassigned	dsDNA		13	13	0	20	20	0	0	0	2	0	2	35	
	Unassigned	Unclassified dsDNA phages														
	Plant and protozoan virus	Mononegavirales	ssRNA(-)		0	0	0	0	0	0	2	2	0	2	2	4
Picornavirales		ssRNA(+)		0	0	0	0	0	0	2	2	0	0	0	2	
Tymovirales		ssRNA(+)		0	0	0	0	0	0	1	1	0	7	7	8	
Unassigned		ssRNA(+)		0	0	0	0	0	0	5	5	0	0	0	5	
Unassigned		ssRNA(+)		0	0	0	0	0	0	1	1	0	0	0	1	
Unassigned		ssDNA (+/-)		9	9	0	10	10	0	7	7	0	27	27	53	
Unassigned		ssRNA(+)		0	0	0	0	0	0	9	9	0	2	2	11	
Unassigned		dsDNA		0	0	0	1	1	0	0	0	0	0	0	1	

(Continued)

Table 2. (Continued)

Order	Viral family	Genome	Desmodus rotundus						Molossus molossus						Overall total	
			Cave F			Cave M			Urban			Forest				
			Saliva	Feces	Total	Saliva	Feces	Total	Saliva	Feces	Total	Saliva	Feces	Total		
Plant and protozoan virus	Unassigned	dsDNA		1	1	0	1	1	0	0	0	0	0	0	0	2
	Unassigned	ssDNA(+)		3	3	0	1	1	0	2	2	0	7	7	13	
	Unassigned	dsRNA		0	0	0	0	0	0	17	17	0	4	4	21	
	Unassigned	dsDNA		6	6	0	2	2	0	7	7	0	9	9	24	
	Unassigned	ssRNA(+)		0	0	1	0	1	0	0	0	0	1	1	2	
	Unassigned	ssRNA(+)		0	0	0	0	0	0	0	0	0	2	2	2	
	Unassigned	dsRNA		0	0	0	0	0	0	0	0	0	2	2	2	
	Unassigned	ssRNA(+)		0	0	2	0	2	0	5	5	0	0	0	7	
	Unassigned	-		0	0	0	0	0	0	32	32	0	17	17	49	
	Bunyvirales	ssRNA(-)		2	2	0	0	0	0	4	4	0	0	0	6	
Vertebrate virus	Herpesvirales	dsDNA		4	4	3	3	6	0	2	2	450	5	455	467	
	Picornavirales	ssRNA(+)		0	0	0	0	0	0	8	8	0	1	1	9	
	Unassigned	dsDNA		78	78	0	35	35	0	0	0	18	0	18	131	
	Unassigned	ssDNA(-)		272	272	6	348	354	1	56	57	12	9	21	704	
	Unassigned	ssRNA(+)		2	2	0	0	0	0	2	2	0	3	3	7	
	Unassigned	ssDNA (+/-)		117	117	7	65	72	0	160	160	1	195	196	545	
	Unassigned	ssDNA (+/-)		10	10	0	20	20	0	5	5	1	42	43	78	
	Unassigned	ssRNA(+)		0	0	0	0	0	0	1	1	0	0	0	1	
	Unassigned	Papillomaviridae		0	0	0	0	0	1	17	18	93	124	217	235	
	Unassigned	ssDNA (+/-)		111	111	10	187	197	0	3	3	2	6	25	319	
ND	Unassigned	dsDNA		2	2	4	5	9	0	6	6	0	3	3	20	
	Unassigned	dsDNA		0	0	0	6	6	0	0	0	0	0	0	6	
	Unassigned	ssRNA-RT		1	1	1	8	9	2	0	2	69	2	71	83	
	Unassigned	-		55	55	0	49	49	2	166	168	0	235	235	507	
	Unassigned	ssDNA		45	45	0	15	15	0	78	78	4	106	110	248	
	Unassigned	-		0	0	0	12	12	0	3	3	0	5	5	20	
	Unassigned	Unclassified viruses		1 930	1 930	39	2 554	2 593	8	2 692	2 700	729	3 039	3 768	10 991	

The number of viral-associated contig sequences is given for each sample (saliva and feces), overall, as well as for each habitat (caves F and M for *D. rotundus*, urban and forest for *M. molossus*). The total number of contigs detected for each collecting site is given in bold.

<https://doi.org/10.1371/journal.pone.0186943.t002>

We found viral sequences related to *Poxviridae* in both species, but only the *D. rotundus* trapped in cave M presented mammal-related poxviruses. These sequences presented high amino acid homology (>90%) with the *bovine papular stomatitis virus*, suggesting that this virus came directly from cattle and its presence was related to the dietary habit of *D. rotundus* (i.e., *D. rotundus* feeds off the blood of cattle).

RNA viruses were found in distinct bat species or habitats and belonged to the *Astroviridae*, *Nairoviridae*, *Hepeviridae*, *Picornaviridae* and *Retroviridae* families. Astroviruses were only found in *D. rotundus* trapped in cave F but in both habitats for *M. molossus*. In contrast, nairoviruses were only found in *D. rotundus* trapped in cave F and in *M. molossus* trapped in forest habitats. Hepeviruses were only found in *M. molossus* trapped in urban areas. Picornaviruses were found in both habitats for *M. molossus*, but the only sequence detected in forest habitats was different from those detected in urban habitats.

Viral characterization and phylogenetic relationships of selected viruses

Bat circoviruses. The family *Circoviridae* is known to infect birds, mammals and insects, and comprises two genera, *Circovirus* and *Cyclovirus* (ICTV 2016). Their transmission occurs primarily through the fecal–oral route [54]. Here, *Circoviridae* viruses were detected in both *D. rotundus* and *M. molossus* samples, whatever the environment, with a total of 545 contigs (Table 2). These viruses were essentially found in the feces, except for *D. rotundus* in cave M for which a small proportion of sequences was also detected in saliva. Four complete genomes of circoviruses (CVs) were obtained for *M. molossus* ($n = 1$ in urban habitats and $n = 3$ in forest habitats), and one partial sequence corresponding to the complete replication-associated protein gene for *D. rotundus* in cave M ($n = 1$) (Genbank acc. nb.: KX812440–KX812443 and KX954092). The complete genomes presented the archetypal genome organization of CVs, with the two inversely arranged ORFs encoding the Rep and Cap proteins (Fig 3). The intergenic region was variable in length, depending on the species as well as the habitat. The full-length REP was used to perform evolutionary analyses. The phylogenetic tree revealed that CVs (identified in cave M for *D. rotundus* and forest habitats for *M. molossus*) formed two distinct monophyletic clades with other bat circoviruses detected in the gastrointestinal tract and feces samples of Chinese bat species [55] (Fig 3). In contrast, the CV identified in urban areas for *M. molossus* was closely related to CVs detected in birds, with a posterior probability of 0.84.

Bat spumaviruses. *Spumavirus* (ICTV 2016) constitutes the only genus of the *Spumaretrovirinae* subfamily, which belongs to the *Retroviridae* family. They are highly prevalent in several animal species (e.g., cats, cows, horses and nonhuman primates) and currently six species have been described in the genus. Transmission of spumaviruses between nonhuman primates and cross-species transmission to humans occurs mainly through saliva (e.g., by licking, aggressive contacts, bites) [56–58]. Nevertheless, other routes of transmission, such as vertical transmission, are being studied [59–61]. In this study, a 312-nt-length fragment (Genbank acc. nb.: KX812444), obtained with two contigs detected in the saliva sample of *M. molossus* in urban habitats, showed 54.88% and 48.48% nucleotide and amino acid identity, respectively, with the *pol* gene of *Rhinolophus affinis foamy virus 1* (Genbank acc. nb.: JQ814855) detected in the feces of Chinese bats [13]. Phylogenetic analysis revealed that the foamy virus (FV) identified was close to *Rhinolophus affinis foamy virus 1*, with a high posterior probability value of 0.91 (Fig 4). Bat FVs seem to share a common origin with equine, bovine and feline FVs. In addition, this analysis shows that bat FVs are divergent from those infecting nonhuman primates.




Fig 3. Circular genome maps of five putative circoviruses directly recovered from metagenomic data of *M. molossus* and *D. rotundus* bat feces and their phylogenetic relationships with other representative members of the *Circoviridae* family. (A) The inversely arranged open-reading frames encoding the putative replication-associated protein (REP) and capsid protein (CAP) are shown in green and blue boxes, respectively. The genome organization was determined with GENEIOUS R9. (B) The phylogenetic analysis is based on the REP protein sequences (alignment of 109 amino acids). The blue arrows indicate the five REP sequences of bat-sourced circoviruses obtained in the present study. The tree was inferred using the Bayesian method with the blosum62 model. Sequence identifiers include the NCBI accession number and the isolate name. Posterior probabilities of the Bayesian analysis (>50%) are shown next to each node. The scale bar indicates amino acid substitutions per site.

<https://doi.org/10.1371/journal.pone.0186943.g003>




Fig 4. Phylogenetic analysis of partial sequences of the *pol*s region (alignment of 107 amino acids) directly obtained from the metagenomic data of pooled fecal samples of *M. molossus* with other representative members of the *Spumavirus* genus. The tree was inferred using the Bayesian method with the WAG + G model. Sequence identifiers include the NCBI accession number and the isolate name. The blue arrows indicate the sequence of bat-sourced spumavirus obtained in the present study. Posterior probabilities of the Bayesian analysis (>50%) are shown next to each node. The scale bar indicates amino acid substitutions per site.

<https://doi.org/10.1371/journal.pone.0186943.g004>

Bat herpesviruses. Mammalian herpesviruses (HVs) belong to the *Herpesviridae* family (order *Herpesvirales*). HVs are organized in three subfamilies—*Alphaherpesvirinae*, *Betaherpesvirinae* and *Gammapherpesvirinae*. HVs can infect a wide range of hosts, including humans. Most HV-related contigs were found in the saliva sample of *M. molossus* in forest habitats. We identified a 943-nt-length fragment covering the *DNA polymerase* gene (Genbank acc. nb.: KX812446). This fragment displayed from 62.46% to 71.31% and from 68.93% to 80.58% nucleotide and amino acid identities, respectively, with other bat HVs. The highest percentage of identity in amino acids was observed with *Myotis ricketti herpesvirus 1* (Genbank acc. nb.: JN692429). Phylogenetic analysis showed that the HV identified belonged to the *Gammapherpesvirinae* subfamily and was related to the *Tupaia belangeri gammaherpesvirus 1* and the *Myotis ricketti herpesvirus 1*, with a posterior probability of 0.68 (Fig 5).

Bat nairoviruses. The *Nairoviridae* family belongs to the order *Bunyavirales* (ICTV 2016). This family encompasses at least nine serogroups, including the Crimean-Congo hemorrhagic fever serogroup. Nairoviruses (NVs) are primarily tick-borne viruses capable of infecting different vertebrate hosts. Four contigs matching nairoviruses (NVs) were detected in the feces of *M. molossus* in urban areas (Table 2). The contigs covered 1 kb of segment L (Genbank acc. nb.: KX821677) and showed a 43% pairwise homology in amino acids with the RNA-dependent RNA polymerase of the previously described *Rhinolophus pearsoni bunyavirus* (Genbank acc. nb.: KC154063). Phylogenetic analysis revealed that the NV identified clustered with *Rhinolophus pearsoni bunyavirus*, with a posterior probability of 0.74 (Fig 6). These NVs share a




Fig 5. Phylogenetic analysis of partial *pol* gene sequences (alignment of 951 nucleotides) directly obtained from the metagenomic data of pooled fecal samples of *M. molossus* with other representative members of the *Herpesviridae* family. The tree was inferred using the Bayesian method with the GTR + G + I model. Sequence identifiers include the NCBI accession number and the isolate name. The blue arrows indicate the bat-sourced herpesvirus obtained in the present study. Posterior probabilities of the Bayesian analysis (>50%) are shown next to each node. The scale bar indicates amino acid substitutions per site.

<https://doi.org/10.1371/journal.pone.0186943.g005>

common ancestor with mosquito NVs, with whom they clustered with a posterior probability of 1.

Bat papillomaviruses. *Papillomaviridae* are nonenveloped dsDNA viruses. Currently, 49 genera have been identified in this family, including the most species-abundant genera *Gammapapillomavirus*. Papillomaviruses (PVs) are usually asymptomatic and thought to be rarely transmitted between species. In this study, a total of 235 contigs related to PVs were detected in *M. molossus* samples, with most of them identified in the feces collected in forest habitats (Table 2). The longest contig obtained covered the entire genome of known PVs and was named MmoPV1 (Genbank acc. nb.: KX812447). The MmoPV1 genome was 7,869 bp in length with a 40% G+C content. The MmoPV1 genome had the typical organization of PV ORFs (S2 Fig). The phylogenetic tree indicated that MmoPV1 possessed a basal position of a cluster of PVs encompassing mupa PVs, kappa PVs, lambda PVs, sigma PVs, nupa PVs and novel unclassified bat PVs detected in *Miniopterus schreibersii* [12], *Eidolon helvum* (Germany), *Eptesicus serotinus* and *Rhinolophus ferrumequinum* [62], with a high posterior probability value of 0.98 (Fig 7).

Discussion

A wide spectrum of viruses capable of infecting a wide range of animals, plants, insects and bacteria was detected in bats. Since the 1950s, 28 viral families including more than 50 genera




Fig 6. Phylogenetic analysis of partial RdRp protein sequences (alignment of 341 amino acids) directly obtained from the metagenomic data of pooled fecal samples of *M. molossus* with other representative members of the *Nairoviridae* family. The tree was inferred using the Bayesian method with the WAG + G model. Sequence identifiers include the NCBI accession number and the isolate name. The blue arrows indicate the bat-sourced nairovirus sequence obtained in the present study. Posterior probabilities of the Bayesian analysis (>50%) are shown next to each node. The scale bar indicates amino acid substitutions per site.

<https://doi.org/10.1371/journal.pone.0186943.g006>

have been described in bats [18]. With the advent of new-generation technologies, the characterization of bat viromes and the discovery of related variants of known viruses and new bat-borne viruses continue to increase rapidly. The key purpose of the present work was to explore the viral composition of feces and saliva samples of *Desmodus rotundus* and *Molossus molossus* bat species, living in sympatry but using distinct microhabitats in French Guiana.

Overall, this study identified 10,991 viral-associated contigs distributed within 51 known viral families of which 14 are known to infect mammals. New mammalian-related viral sequences/variants were discovered, adding knowledge on the bat-borne viral population. The proportion of eukaryote- and prokaryote-related sequences was low (less than 2% and 20%, respectively). About 0.30% of total contigs were virus-related sequences, with differences in composition depending on the type of sample, the investigated species, habitats, and fine-scale ecological heterogeneity (Table E in S1 File). For instance, of the 30 viral families detected in *D. rotundus*, only 20 were common to both sampling sites (cave F vs. cave M). Likewise, focusing on vertebrate-related viral families, eight of the 11 viral families detected for this species




Fig 7. Phylogenetic analysis of the complete L1 protein sequence (alignment of 459 amino acids) directly obtained from the metagenomic data of pooled fecal samples of *M. molossus* with other representative members of the *Papillomaviridae* family. The tree was inferred using the Bayesian method with the WAG + G model. Sequence identifiers include the NCBI accession number. The blue arrows indicate the bat-sourced papillomavirus sequence obtained in the present study. Posterior probabilities of the

Bayesian analysis (>50%) are shown next to each node. The scale bar indicates amino acid substitutions per site.

<https://doi.org/10.1371/journal.pone.0186943.g007>

were shared between the populations sampled in the two caves. For the *M. molossus* species, of the 46 viral families detected, 29 were common to both habitats (forest areas vs. urban areas). Focusing on vertebrate-related viral families, ten of the 12 viral families detected in this species were common to both habitats. Furthermore, we observed differences in the viral genera and species detected, depending on the sampling sites for both species. These results suggest that habitats may play a role in shaping the viral diversity harbored by the bats investigated. This observation was also supported by Hu *et al.* [16], who showed that the metagenomic approach can reflect differences in viral diversity carried by different bats in different regions. However, a verification of confounding factors (*e.g.*, sampling size, type of sample, capture effort, species, feeding strategies) will be necessary to confirm these results. Likewise, population structure analysis at the species level and studies on interactions with other bat species using the same habitat are needed to determine their respective roles in shaping viral diversity.

Regardless of the bat species and the habitat studied, phage-related sequences represented a significant proportion of the viral-associated sequences identified (Table 2 and Table F in S1 File). Sequences related to the order *Caudovirales* and the *Microviridae* family were the most frequent. Phage-related sequences detected in the feces accounted for most of this proportion, as could be expected considering the type of sample. Despite the constant and rapid expansion of existing databases, sequences lacking counterparts remained the largest part of the data. These global results are in agreement with previous virome studies conducted on North American and Eurasian bat species [8,9,11,12,14,15] and highlight the need to pursue investigations and taxonomic assignments on microorganism communities. Therefore, the underestimation of the total number of bat-borne viruses present in the samples, including highly divergent viruses, cannot be excluded.

Despite a smaller number of samples for *M. molossus*, the number of viral-associated contigs detected in these samples was higher than that found in *D. rotundus*, suggesting a higher number of viruses passing through this species. However, these viruses are likely related to nondigested viruses associated with the diet of their prey but not directly to the host. Indeed, as highlighted in Li *et al.*, insect and plant viruses were mostly represented in the insectivorous species, while they accounted for less than 5% in the vampire species [9]. The diet of the vampire bat (vertebrate blood) may also be reflected in the high proportion of vertebrate-related viral sequences found. Likewise, the divergence observed in the distribution of phage-related viruses might be associated with the diets of the species investigated and/or to their phylogenetic position, Molossidae vs. Phyllostomidae. For instance, *Podoviridae* species were the most frequent in *D. rotundus*, whatever the sampling site, while *Microviridae* species were the most often found in *M. molossus*. In bats, former studies highlighted a strong association between the microbiota, host phylogeny, life history, physiology as well as locality [63,64]. Given that gut microbiota is known to be highly dependent on diet, these results may reflect differences between the dietary ecology of *D. rotundus* and *M. molossus* species.

We found several novel mammalian-related viral sequences/variants from the *Anelloviridae*, *Nairoviridae*, *Circoviridae*, *Hepeviridae*, *Herpesviridae*, *Retroviridae* and *Papillomaviridae* families. These viral sequences were closely related to sequences detected in Old World bat species [17,62,65–68]. Furthermore, most of the families identified in Old World bat species were identified here, even for non-mammalian viruses. Moreover, where divergence could be expected due to geographical isolation, we observed highly supported phylogenetic proximity between Old and New World bat viruses in their respective continents, revealing the existence

of common evolutionary processes and supporting the long evolutionary hypothesis of bats and their viruses [7]. For example, most of the circovirus (CV) sequences detected in the present study shared high sequence identities with bat CVs detected in several bat species trapped in different Chinese provinces [17,69] and clustered in highly supported monophyletic groups. The results reported herein suggest that the associated viruses may be hosted by bats. Conversely, the highly supported clustering of *M. molossus* CV (KX812443) with avian CVs may result from a cross-species jump from birds to bats, as reported by Hu *et al.* [16] and Lima *et al.* [70]. Likewise, foamy virus (FV) sequences, also called spumaviruses, detected in saliva swab samples of *M. molossus* showed a high genomic similarity with the FVs detected in pharyngeal and rectal swab samples of *Rhinolophus affinis* reported by Wu *et al.* [68], but low similarity with other spumaviruses infecting other hosts (*e.g.*, equines, felines, bovines, nonhuman primates). Our findings indicate that FVs can indeed infect bats, which could be considered as potential reservoirs and dispersers of this viral genus.

Several studies reported the detection of herpesviruses (HVs) of the three subfamilies (*Alpha*-, *Beta*- and *Gammaherpesvirinae*) in feces, anal swabs, digestive tract and saliva samples of bats [16,17,68,71–77]. These findings suggested that the oral–fecal transmission route may be important for the transmission of HVs in bats. Furthermore, cases of cross-species transmission of HVs were also reported in bats, but their mode of transmission is still unclear [77]. In this study, we found a high diversity of HVs in the feces and saliva samples of both species investigated. Moreover, some sequences detected in the saliva samples of *M. molossus* from forest habitats were also found in the feces samples, which showed the capacity of these bats to shed HVs in the environment and supported the hypothesis of a potential oral–fecal transmission route. Detected HV-related sequences were novel and showed a high sequence similarity with other bat HVs. Nevertheless, the phylogenetic tree constructed using a Bayesian method revealed that *M. molossus herpesvirus 1* (KX812446) did not cluster in a unique group with other known bat gamma-HVs. Rather, it shares a common ancestor with *Myotis ricketti* and *Tupaia belangeri* gamma-HVs, stressing the need for further characterization of *M. molossus herpesvirus 1* to confirm its taxonomic assignment.

Nairoviruses (NVs) naturally infecting bats were reported in a few studies, with some capable of causing intestine and hepatic disorders in other mammals [15,78–80]. Evidence of neutralizing antibodies against Crimean-Congo hemorrhagic fever (CCHF) was reported by Müller *et al.* [81], but the role of bats regarding the cycle and dispersal of this virus is not clear. Here, we found a novel NV-related sequence in the feces of *M. molossus* sampled in urban areas. Phylogenetic relationships based on the partial segment L sequence showed a highly supported clustering with another NV-related sequence detected from the feces of *Rhinolophus pearsoni*. This analysis also revealed that the detected NV was related to the genus *Phlebovirus*, primarily constituted of tick-, mosquito- and phlebotomine-borne viruses. These findings suggest that this sequence may be related to undigested viruses associated with the bats' diet rather than viruses hosted by bats. However, screening of other organs will be necessary to rule out the role of bats in the virus cycle and dispersal.

Diverse sequences related to papillomaviruses (PVs) were reported in bats [16,17,62,66,67,82,83]. Here, we detected 235 PV-like sequences related to members of genetically diverse genera within the *Papillomaviridae*, indicating that bat species might be associated with a wide diversity of PVs. We described the complete genome of a novel bat PV, MmoPV1 (KX812447), detected in the feces and saliva samples of *M. molossus* sampled in forest and urban habitats. This is the first detection of PV in this species to date. Comparing MmoPV1 with other PVs showed that it contains the typical PV ORFs coding for four putative early proteins (E6, E7, E1, E2) and two putative late capsid proteins (L2 and L1). The putative early protein E4 –present in other phylogenetically closely-related bat PVs (JQ692938 [67], JX123128

[83], KC858263, KC858265, KC858266 [84])—was absent in MmoPV1. Furthermore, the phylogenetic tree revealed that MmoPV1 may be more ancient than these bat PVs, suggesting that they may either have undergone a divergent evolution that allowed them to acquire the early protein or represent different host-specific and area-specific lineages [84].

This study is the very first to explore the virome of fecal and saliva samples obtained from the two common Amazonian bat species, *M. molossus* and *D. rotundus*, with overlapping distribution but using distinct microhabitats in French Guiana. It provides important insight into the feces and saliva viromes of the two bat species investigated within contrasting habitats.

However, to fully understand how viral diversity is shaped in *D. rotundus* and *M. molossus*, further characterization of their geographic range, foraging-induced interactions, species distribution and gregariousness will be necessary. Indeed, larger geographic ranges encompass greater diversity of ecosystems/habitats, which could lead to a spatial population substructure [35]. The overlapping distribution of reservoirs and recipient hosts, the social complexity of bats as well as species-specific evolutionary and life-history traits are known as predictors of viral richness in wildlife [6,35]. Likewise, the contact and proximity between individuals influence the presence, abundance and diversity of the pathogens harbored [85]. Consequently, deciphering the virome of bats should consider the size of the whole community of bats, the genetic structure of populations in a given habitat and the rates of interactions among them, all parameters contributing to the viral diversity observed for bat species.

Supporting information

S1 File. Read and contig processing of *D. rotundus* and *M. molossus* samples. Read data reduction steps and *de novo* assembly of processed reads. Contig and sequential BLAST comparisons, with the total number of contigs and viral families identified in *D. rotundus* and *M. molossus* samples.
(DOCX)

S1 Fig. Map of collecting sites of all samples/individuals used across French Guiana. The total number of collected samples per site is given in parentheses. Pie chart indicates the proportion of feces (orange) and saliva (orange dots) samples collected. Detailed characteristics of the different collecting sites are given in Table 1.
(TIF)

S2 Fig. Predicted genome organization of the putative papillomavirus sequence directly recovered from the metagenomic data of *M. molossus* bat feces. The open-reading frames encoding the putative late (L1 and L2) and early (E1, E2, E6 and E7) proteins are shown in green boxes. The genome organization was determined with GENEIOUS R9.
(TIF)

Acknowledgments

We acknowledge François Catzeffis and Marguerite Delaval for useful discussions on the biology, ecology and dynamics of bats. All field volunteers and owners and/or managers of capture sites are warmly acknowledged for their assistance in captures. We thank the INDA and C3BI teams at the Institut Pasteur and UMR BIOGECO for their technical support and access to the clusters. We thank L. Ma and M. Tichit for technical assistance.

Author Contributions

Conceptualization: Arielle Salmier, Benoit de Thoisy, Vincent Lacoste, Anne Lavergne.

Data curation: Arielle Salmier, Sourakhata Tirera, Alain Franc, Christiane Bouchier.

Formal analysis: Arielle Salmier.

Funding acquisition: Anne Lavergne.

Investigation: Arielle Salmier, Benoit de Thoisy, Edith Darcissac, Damien Donato, Anne Lavergne.

Methodology: Sourakhata Tirera, Alain Franc, Christiane Bouchier.

Project administration: Anne Lavergne.

Supervision: Anne Lavergne.

Writing – original draft: Arielle Salmier, Sourakhata Tirera, Benoit de Thoisy, Alain Franc, Edith Darcissac, Damien Donato, Christiane Bouchier, Vincent Lacoste, Anne Lavergne.

Writing – review & editing: Arielle Salmier, Benoit de Thoisy, Alain Franc, Vincent Lacoste, Anne Lavergne.

References

1. Moratelli R, Calisher CH. Bats and zoonotic viruses: can we confidently link bats with emerging deadly viruses? *Mem Inst Oswaldo Cruz*. 2015; 110: 1–22. <https://doi.org/10.1590/0074-02760150048> PMID: 25742261
2. Olival KJ, Weekley CC, Daszak P. Are Bats Really “Special” as Viral Reservoirs? What We Know and Need to Know. In: Wang L-F, Cowled C, editors. *Bats and Viruses*. John Wiley & Sons, Inc; 2015. pp. 281–294. <https://doi.org/10.1002/9781118818824.ch11>
3. Olival KJ, Hosseini PR, Zambrana-Torrel C, Ross N, Bogich TL, Daszak P. Host and viral traits predict zoonotic spillover from mammals. *Nature*. 2017; 546: 646–650. <https://doi.org/10.1038/nature22975> PMID: 28636590
4. Brook CE, Dobson AP. Bats as ‘special’ reservoirs for emerging zoonotic pathogens. *Trends in Microbiology*. 2015; 23: 172–180. <https://doi.org/10.1016/j.tim.2014.12.004> PMID: 25572882
5. Luis AD, O’Shea TJ, Hayman DTS, Wood JLN, Cunningham AA, Gilbert AT, et al. Network analysis of host–virus communities in bats and rodents reveals determinants of cross-species transmission. *Ecol Lett*. 2015; 18: 1153–1162. <https://doi.org/10.1111/ele.12491> PMID: 26299267
6. Plowright RK, Eby P, Hudson PJ, Smith IL, Westcott D, Bryden WL, et al. Ecological dynamics of emerging bat virus spillover. *Proceedings of the Royal Society B: Biological Sciences*. 2014; 282: 20142124–20142124. <https://doi.org/10.1098/rspb.2014.2124> PMID: 25392474
7. Schneeberger K, Voigt CC. Zoonotic Viruses and Conservation of Bats. In: Voigt CC, Kingston T, editors. *Bats in the Anthropocene: Conservation of Bats in a Changing World*. Cham: Springer International Publishing; 2016. pp. 263–292. http://link.springer.com/10.1007/978-3-319-25220-9_10
8. Donaldson EF, Haskew AN, Gates JE, Huynh J, Moore CJ, Frieman MB. Metagenomic Analysis of the Viromes of Three North American Bat Species: Viral Diversity among Different Bat Species That Share a Common Habitat. *Journal of Virology*. 2010; 84: 13004–13018. <https://doi.org/10.1128/JVI.01255-10> PMID: 20926577
9. Li L, Victoria JG, Wang C, Jones M, Fellers GM, Kunz TH, et al. Bat Guano Virome: Predominance of Dietary Viruses from Insects and Plants plus Novel Mammalian Viruses. *Journal of Virology*. 2010; 84: 6955–6965. <https://doi.org/10.1128/JVI.00501-10> PMID: 20463061
10. Canuti M, Eis-Huebinger AM, Deijs M, de Vries M, Drexler JF, Oppong SK, et al. Two Novel Parvoviruses in Frugivorous New and Old World Bats. Vartanian J-P, editor. *PLoS ONE*. 2011; 6: e29140. <https://doi.org/10.1371/journal.pone.0029140> PMID: 22216187
11. Ge X, Li Y, Yang X, Zhang H, Zhou P, Zhang Y, et al. Metagenomic Analysis of Viruses from Bat Fecal Samples Reveals Many Novel Viruses in Insectivorous Bats in China. *Journal of Virology*. 2012; 86: 4620–4630. <https://doi.org/10.1128/JVI.06671-11> PMID: 22345464
12. Tse H, Tsang AKL, Tsoi H-W, Leung ASP, Ho C-C, Lau SKP, et al. Identification of a Novel Bat Papillomavirus by Metagenomics. Davis T, editor. *PLoS ONE*. 2012; 7: e43986. <https://doi.org/10.1371/journal.pone.0043986> PMID: 22937142

13. Wu Z, Ren X, Yang L, Hu Y, Yang J, He G, et al. Virome Analysis for Identification of Novel Mammalian Viruses in Bat Species from Chinese Provinces. *Journal of Virology*. 2012; 86: 10999–11012. <https://doi.org/10.1128/JVI.01394-12> PMID: 22855479
14. He B, Li Z, Yang F, Zheng J, Feng Y, Guo H, et al. Virome Profiling of Bats from Myanmar by Metagenomic Analysis of Tissue Samples Reveals More Novel Mammalian Viruses. Tse H, editor. *PLoS ONE*. 2013; 8: e61950. <https://doi.org/10.1371/journal.pone.0061950> PMID: 23630620
15. Dacheux L, Cervantes-Gonzalez M, Guigon G, Thiberge J-M, Vandenbogaert M, Maufrais C, et al. A Preliminary Study of Viral Metagenomics of French Bat Species in Contact with Humans: Identification of New Mammalian Viruses. Tse H, editor. *PLoS ONE*. 2014; 9: e87194. <https://doi.org/10.1371/journal.pone.0087194> PMID: 24489870
16. Hu D, Zhu C, Wang Y, Ai L, Yang L, Ye F, et al. Virome analysis for identification of novel mammalian viruses in bats from Southeast China. *Scientific Reports*. 2017; 7. <https://doi.org/10.1038/s41598-017-11384-w> PMID: 28883450
17. Wu Z, Yang L, Ren X, He G, Zhang J, Yang J, et al. Deciphering the bat virome catalog to better understand the ecological diversity of bat viruses and the bat origin of emerging infectious diseases. *ISME J*. 2016; 10: 609–620. <https://doi.org/10.1038/ismej.2015.138> PMID: 26262818
18. Calisher CH. Viruses in bats: a historic review. In: Wang L-F, Cowled C, editors. *Bats and Viruses: A New Frontier of Emerging Infectious Diseases*. 2015. p. 23.
19. Cibulski SP, Teixeira TF, de Sales Lima FE, do Santos HF, Franco AC, Roehe PM. A Novel Anelloviridae Species Detected in *Tadarida brasiliensis* Bats: First Sequence of a Chiropteran Anellovirus. *Genome Announc*. 2014; 2. <https://doi.org/10.1128/genomeA.01028-14> PMID: 25359906
20. de S Lima FE, Cibulski SP, Dall Bello AG, Mayer FQ, Witt AA, Roehe PM, et al. A Novel Chiropteran Circovirus Genome Recovered from a Brazilian Insectivorous Bat Species. *Genome Announcements*. 2015; 3: e01393–15. <https://doi.org/10.1128/genomeA.01393-15> PMID: 26607898
21. de S Lima FE, Cibulski SP, dos Santos HF, Teixeira TF, Varela APM, Roehe PM, et al. Genomic Characterization of Novel Circular ssDNA Viruses from Insectivorous Bats in Southern Brazil. *PLOS ONE*. 2015; 10: e0118070. <https://doi.org/10.1371/journal.pone.0118070> PMID: 25688970
22. Drexler JF, Corman VM, Müller MA, Maganga GD, Vallo P, Binger T, et al. Bats host major mammalian paramyxoviruses. *Nature Communications*. 2012; 3: 796. <https://doi.org/10.1038/ncomms1796> PMID: 22531181
23. Tong S, Zhu X, Li Y, Shi M, Zhang J, Bourgeois M, et al. New World Bats Harbor Diverse Influenza A Viruses. *PLOS Pathogens*. 2013; 9: e1003657. <https://doi.org/10.1371/journal.ppat.1003657> PMID: 24130481
24. de S Lima FE, Cibulski SP, Witt AA, Franco AC, Roehe PM. Genomic characterization of two novel polyomaviruses in Brazilian insectivorous bats. *Arch Virol*. 2015; 160: 1831–1836. <https://doi.org/10.1007/s00705-015-2447-6> PMID: 25963124
25. Catzefflis F. Liste des Mammifères de Guyane française (octobre 2015). 2015; http://www.kwata.net/medias/images/upload/MAMM-GUYANE_Catzefflis-oct2015.pdf
26. Brosset A, Charles-Dominique P, Cockle A, Cosson J-F, Masson D. Bat communities and deforestation in French Guiana. *Can J Zool*. 1996; 74: 1974–1982. <https://doi.org/10.1139/z96-224>
27. Delaval M, Charles-Dominique P. Edge effects on frugivorous and nectarivorous bat communities in a neotropical primary forest in French Guiana. *Revue d'écologie*. 2006; 61: 343–352.
28. Jung K, Threlfall CG. Urbanisation and Its Effects on Bats—A Global Meta-Analysis. In: Voigt CC, Kingston T, editors. *Bats in the Anthropocene: Conservation of Bats in a Changing World*. Cham: Springer International Publishing; 2016. pp. 13–33. http://link.springer.com/10.1007/978-3-319-25220-9_2
29. de Thoisy B, Bourhy H, Delaval M, Pontier D, Dacheux L, Darcissac E, et al. Bioecological Drivers of Rabies Virus Circulation in a Neotropical Bat Community. *PLOS Neglected Tropical Diseases*. 2016; 10: e0004378. <https://doi.org/10.1371/journal.pntd.0004378> PMID: 26808820
30. Gay N, Olival KJ, Bumrungsri S, Siriaronrat B, Bourgarel M, Morand S. Parasite and viral species richness of Southeast Asian bats: Fragmentation of area distribution matters. *International Journal for Parasitology: Parasites and Wildlife*. 2014; 3: 161–170. <https://doi.org/10.1016/j.ijppaw.2014.06.003> PMID: 25161915
31. Wilkinson GS. The social organization of the common vampire bat. *Behav Ecol Sociobiol*. 1985; 17: 123–134.
32. Wilkinson GS. The social organization of the common vampire bat. *Behav Ecol Sociobiol*. 1985; 17: 111–121.
33. Johnson N, Aréchiga-Ceballos N, Aguilar-Setien A. Vampire Bat Rabies: Ecology, Epidemiology and Control. *Viruses*. 2014; 6: 1911–1928. <https://doi.org/10.3390/v6051911> PMID: 24784570

34. Blackwood JC, Streicker DG, Altizer S, Rohani P. Resolving the roles of immunity, pathogenesis, and immigration for rabies persistence in vampire bats. *PNAS*. 2013; 110: 20837–20842. <https://doi.org/10.1073/pnas.1308817110> PMID: 24297874
35. Turmelle AS, Olival KJ. Correlates of Viral Richness in Bats (Order Chiroptera). *EcoHealth*. 2009; 6: 522–539. <https://doi.org/10.1007/s10393-009-0263-8> PMID: 20049506
36. Dechmann DKN, Kranstauber B, Gibbs D, Wikelski M. Group Hunting—A Reason for Sociality in Molossid Bats? *PLOS ONE*. 2010; 5: e9012. <https://doi.org/10.1371/journal.pone.0009012> PMID: 20140247
37. Sikes RS, Gannon WL. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *Journal of Mammalogy*. 2011; 92: 235–253. <https://doi.org/10.1644/10-MAMM-F-355.1>
38. Victoria JG, Kapoor A, Li L, Blinkova O, Slikas B, Wang C, et al. Metagenomic Analyses of Viruses in Stool Samples from Children with Acute Flaccid Paralysis. *Journal of Virology*. 2009; 83: 4642–4651. <https://doi.org/10.1128/JVI.02301-08> PMID: 19211756
39. Prescott LM, Sherwood LM, Woolverton CJ. Les virus : introduction et caractères généraux. Microbiologie. De Boeck Supérieur; 2010.
40. Allander T, Emerson SU, Engle RE, Purcell RH, Bukh J. A virus discovery method incorporating DNase treatment and its application to the identification of two bovine parvovirus species. *Proceedings of the National Academy of Sciences*. 2001; 98: 11609–11614.
41. Berthet N, Reinhardt AK, Leclercq I, van Ooyen S, Batéjat C, Dickinson P, et al. Phi29 polymerase based random amplification of viral RNA as an alternative to random RT-PCR. *BMC Molecular Biology*. 2008; 9: 77. <https://doi.org/10.1186/1471-2199-9-77> PMID: 18771595
42. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 2014; 30: 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170> PMID: 24695404
43. Brown CT, Howe A, Zhang Q, Pyrkosz AB, Brom TH. A Reference-Free Algorithm for Computational Normalization of Shotgun Sequencing Data. *arXiv:12034802 [q-bio]*. 2012; <http://arxiv.org/abs/1203.4802>
44. Crusoe MR, Alameldin HF, Awad S, Boucher E, Caldwell A, Cartwright R, et al. The khmer software package: enabling efficient nucleotide sequence analysis. *F1000Research*. 2015; <https://doi.org/10.12688/f1000research.6924.1> PMID: 26535114
45. Baker KS, Leggett RM, Bexfield NH, Alston M, Daly G, Todd S, et al. Metagenomic study of the viruses of African straw-coloured fruit bats: Detection of a chiropteran poxvirus and isolation of a novel adenovirus. *Virology*. 2013; 441: 95–106. <https://doi.org/10.1016/j.virol.2013.03.014> PMID: 23562481
46. Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome research*. 2008; 18: 821–829. <https://doi.org/10.1101/gr.074492.107> PMID: 18349386
47. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *Journal of Computational Biology*. 2012; 19: 455–477. <https://doi.org/10.1089/cmb.2012.0021> PMID: 22506599
48. Edgar RC. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*. 2010; 26: 2460–2461. <https://doi.org/10.1093/bioinformatics/btq461> PMID: 20709691
49. Kears M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*. 2012; 28: 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199> PMID: 22543367
50. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucl Acids Res*. 2002; 30: 3059–3066. <https://doi.org/10.1093/nar/gkf436> PMID: 12136088
51. Darriba D, Taboada GL, Doallo R, Posada D. jModelTest 2: more models, new heuristics and parallel computing. *Nat Meth*. 2012; 9: 772–772. <https://doi.org/10.1038/nmeth.2109> PMID: 22847109
52. Darriba D, Taboada GL, Doallo R, Posada D. ProtTest 3: fast selection of best-fit models of protein evolution. *Bioinformatics*. 2011; 27: 1164–1165. <https://doi.org/10.1093/bioinformatics/btr088> PMID: 21335321
53. Rambaut A, Suchard M, Xie D, Drummond A. Tracer v1.6. <http://beast.bio.ed.ac.uk/Tracer>
54. Li L, Kapoor A, Slikas B, Bamidele OS, Wang C, Shaikat S, et al. Multiple Diverse Circoviruses Infect Farm Animals and Are Commonly Found in Human and Chimpanzee Feces. *Journal of Virology*. 2010; 84: 1674–1682. <https://doi.org/10.1128/JVI.02109-09> PMID: 20007276
55. Ge X, Li J, Peng C, Wu L, Yang X, Wu Y, et al. Genetic diversity of novel circular ssDNA viruses in bats in China. *Journal of General Virology*. 2011; 92: 2646–2653. <https://doi.org/10.1099/vir.0.034108-0> PMID: 21795473

56. Murray SM, Linial ML. Foamy virus infection in primates. *Journal of Medical Primatology*. 2006; 35: 225–235. <https://doi.org/10.1111/j.1600-0684.2006.00171.x> PMID: 16872286
57. Voevodin AF, Marx PA. Spumaviruses. *Simian Virology*. Blackwell Publishing Ltd.; 2009. pp. 217–233. <https://doi.org/10.1002/9780813809793.ch9>
58. Mouinga-Ondémé A, Caron M, Nkoghé D, Telfer P, Marx P, Saïb A, et al. Cross-Species Transmission of Simian Foamy Virus to Humans in Rural Gabon, Central Africa. *J Virol*. 2012; 86: 1255–1260. <https://doi.org/10.1128/JVI.06016-11> PMID: 22072747
59. Blasse A, Calvignac-Spencer S, Merkel K, Goffe AS, Boesch C, Mundry R, et al. Mother-Offspring Transmission and Age-Dependent Accumulation of Simian Foamy Virus in Wild Chimpanzees. *J Virol*. 2013; 87: 5193–5204. <https://doi.org/10.1128/JVI.02743-12> PMID: 23449796
60. Liu W, Worobey M, Li Y, Keele BF, Bibollet-Ruche F, Guo Y, et al. Molecular Ecology and Natural History of Simian Foamy Virus Infection in Wild-Living Chimpanzees. *PLOS Pathogens*. 2008; 4: e1000097. <https://doi.org/10.1371/journal.ppat.1000097> PMID: 18604273
61. Muniz CP, Jia H, Shankar A, Troncoso LL, Augusto AM, Farias E, et al. An expanded search for simian foamy viruses (SFV) in Brazilian New World primates identifies novel SFV lineages and host age-related infections. *Retrovirology*. 2015; 12: 94. <https://doi.org/10.1186/s12977-015-0217-x> PMID: 26576961
62. García-Pérez R, Ibáñez C, Godínez JM, Aréchiga N, Garin I, Pérez-Suárez G, et al. Novel Papillomaviruses in Free-Ranging Iberian Bats: No Virus–Host Co-evolution, No Strict Host Specificity, and Hints for Recombination. *Genome Biol Evol*. 2014; 6: 94–104. <https://doi.org/10.1093/gbe/evt211> PMID: 24391150
63. Phillips CD, Phelan G, Dowd SE, McDONOUGH MM, Ferguson AW, Delton Hanson J, et al. Microbiome analysis among bats describes influences of host phylogeny, life history, physiology and geography. *Molecular Ecology*. 2012; 21: 2617–2627. <https://doi.org/10.1111/j.1365-294X.2012.05568.x> PMID: 22519571
64. Carrillo-Araujo M, Taş N, Alcántara-Hernández RJ, Gaona O, Schondube JE, Medellín RA, et al. Phyllostomid bat microbiome composition is associated to host phylogeny and feeding strategies. *Front Microbiol*. 2015; 6: 447. <https://doi.org/10.3389/fmicb.2015.00447> PMID: 26042099
65. Drexler JF, Seelen A, Corman VM, Tateno AF, Cottontail V, Zerbini RM, et al. Bats Worldwide Carry Hepatitis E Virus-Related Viruses That Form a Putative Novel Genus within the Family Hepeviridae. *J Virol*. 2012; 86: 9134–9147. <https://doi.org/10.1128/JVI.00800-12> PMID: 22696648
66. Ge X, Li Y, Yang X, Zhang H, Zhou P, Zhang Y, et al. Metagenomic Analysis of Viruses from the Bat Fecal Samples Reveals Many Novel Viruses in Insectivorous Bats in China. *J Virol*. 2012; JVI.06671-11. <https://doi.org/10.1128/JVI.06671-11> PMID: 22345464
67. Tse H, Tsang AKL, Tsoi H-W, Leung ASP, Ho C-C, Lau SKP, et al. Identification of a Novel Bat Papillomavirus by Metagenomics. *PLOS ONE*. 2012; 7: e43986. <https://doi.org/10.1371/journal.pone.0043986> PMID: 22937142
68. Wu Z, Ren X, Yang L, Hu Y, Yang J, He G, et al. Virome Analysis for Identification of Novel Mammalian Viruses in Bat Species from Chinese Provinces. *J Virol*. 2012; 86: 10999–11012. <https://doi.org/10.1128/JVI.01394-12> PMID: 22855479
69. Ge X, Li J, Peng C, Wu L, Yang X, Wu Y, et al. Genetic diversity of novel circular ssDNA viruses in bats in China. *Journal of General Virology*. 2011; 92: 2646–2653. <https://doi.org/10.1099/vir.0.034108-0> PMID: 21795473
70. de S Lima FE, Cibulski SP, dos Santos HF, Teixeira TF, Varela APM, Roehe PM, et al. Genomic Characterization of Novel Circular ssDNA Viruses from Insectivorous Bats in Southern Brazil. *PLOS ONE*. 2015; 10: e0118070. <https://doi.org/10.1371/journal.pone.0118070> PMID: 25688970
71. Wibbelt G, Kurth A, Yasmum N, Bannert M, Nagel S, Nitsche A, et al. Discovery of herpesviruses in bats. *Journal of General Virology*. 2007; 88: 2651–2655. <https://doi.org/10.1099/vir.0.83045-0> PMID: 17872515
72. Molnár V, Jánoska M, Harrach B, Glávits R, Pálmai N, Rigó D, et al. Detection of a novel bat gammaherpesvirus in Hungary. *Acta Veterinaria Hungarica*. 2008; 56: 529–538. <https://doi.org/10.1556/AVet.56.2008.4.10> PMID: 19149107
73. Razafindratsimandresy R, Jeanmaire EM, Counor D, Vasconcelos PF, Sall AA, Reynes J-M. Partial molecular characterization of alpha herpesviruses isolated from tropical bats. *Journal of General Virology*. 2009; 90: 44–47. <https://doi.org/10.1099/vir.0.006825-0> PMID: 19088271
74. Watanabe S, Ueda N, Iha K, Masangkay JS, Fujii H, Alviola P, et al. Detection of a new bat gammaherpesvirus in the Philippines. *Virus Genes*. 2009; 39: 90. <https://doi.org/10.1007/s11262-009-0368-8> PMID: 19452269

75. Sano K, Okazaki S, Taniguchi S, Masangkay JS, Puentes-pina R, Eres E, et al. Detection of a novel herpesvirus from bats in the Philippines. *Virus Genes*. 2015; 51: 136–139. <https://doi.org/10.1007/s11262-015-1197-6> PMID: 25956292
76. Host KM, Damania B. Discovery of a Novel Bat Gammaherpesvirus. *mSphere*. 2016; 1. <https://doi.org/10.1128/mSphere.00016-16> PMID: 27303690
77. Zheng X, Qiu M, Chen S, Xiao J, Ma L, Liu S, et al. High prevalence and diversity of viruses of the sub-family Gammaherpesvirinae, family Herpesviridae, in fecal specimens from bats of different species in southern China. *Arch Virol*. 2016; 161: 135–140. <https://doi.org/10.1007/s00705-015-2614-9> PMID: 26446885
78. Ishii A, Ueno K, Orba Y, Sasaki M, Moonga L, Hang'ombe BM, et al. A nairovirus isolated from African bats causes haemorrhagic gastroenteritis and severe hepatic disease in mice. *Nature Communications*. 2014; 5: ncomms6651. <https://doi.org/10.1038/ncomms6651> PMID: 25451856
79. Walker PJ, Widen SG, Firth C, Blasdel KR, Wood TG, da Rosa APAT, et al. Genomic Characterization of Yague, Kasokero, Issyk-Kul, Keterah, Gossas, and Thiafora Viruses: Nairoviruses Naturally Infecting Bats, Shrews, and Ticks. *The American Journal of Tropical Medicine and Hygiene*. 2015; 93: 1041–1051. <https://doi.org/10.4269/ajtmh.15-0344> PMID: 26324724
80. Lvov DK, Karas FR, Timofeev EM, Tsyarkin YM, Vargina SG, Veselovskaya OV, et al. "Issyk-Kul" virus, a new arbovirus isolated from bats and Argas (Carios) vespertilionis (Latr., 1802) in the Kirghiz S.S.R. Brief report. *Arch Gesamte Virusforsch*. 1973; 42: 207–209. PMID: 4747048
81. Müller MA, Devignot S, Lattwein E, Corman VM, Maganga GD, Gloza-Rausch F, et al. Evidence for widespread infection of African bats with Crimean-Congo hemorrhagic fever-like viruses. *Scientific Reports*. 2016; 6: srep26637. <https://doi.org/10.1038/srep26637> PMID: 27217069
82. Donaldson EF, Haskew AN, Gates JE, Huynh J, Moore CJ, Frieman MB. Metagenomic Analysis of the Viromes of Three North American Bat Species: Viral Diversity among Different Bat Species That Share a Common Habitat. *J Virol*. 2010; 84: 13004–13018. <https://doi.org/10.1128/JVI.01255-10> PMID: 20926577
83. Baker KS, Leggett RM, Bexfield NH, Alston M, Daly G, Todd S, et al. Metagenomic study of the viruses of African straw-coloured fruit bats: Detection of a chiropteran poxvirus and isolation of a novel adenovirus. *Virology*. 2013; 441: 95–106. <https://doi.org/10.1016/j.virol.2013.03.014> PMID: 23562481
84. García-Pérez R, Ibáñez C, Godínez JM, Aréchiga N, Garin I, Pérez-Suárez G, et al. Novel Papillomaviruses in Free-Ranging Iberian Bats: No Virus–Host Co-evolution, No Strict Host Specificity, and Hints for Recombination. *Genome Biol Evol*. 2014; 6: 94–104. <https://doi.org/10.1093/gbe/evt211> PMID: 24391150
85. Duboscq J, Heeb P. Pathogènes, parasites et réseaux dans les sociétés animales. Analyse des réseaux sociaux appliquée à l'éthologie et l'écologie. Editions Matériologiques; 2015. pp. 343–358.