

## Leishmania infantum Lipophosphoglycan-Deficient Mutants: A Tool to Study Host Cell-Parasite Interplay

Milena Lázaro-Souza, Christine Matte, Jonilson Lima, Guillermo Arango Duque, Graziele Quintela-Carvalho, Áislan de Carvalho Vivarini, Sara Moura-Pontes, Cláudio Pereira Figueira, Flávio Jesus-Santos, Ulisses Gazos

Lopes, et al.

### ▶ To cite this version:

Milena Lázaro-Souza, Christine Matte, Jonilson Lima, Guillermo Arango Duque, Graziele Quintela-Carvalho, et al.. Leishmania infantum Lipophosphoglycan-Deficient Mutants: A Tool to Study Host Cell-Parasite Interplay. Frontiers in Microbiology, 2018, 9, pp.626. 10.3389/fmicb.2018.00626. pasteur-01855962

### HAL Id: pasteur-01855962 https://riip.hal.science/pasteur-01855962

Submitted on 8 Aug 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License





## Leishmania infantum Lipophosphoglycan-Deficient Mutants: A Tool to Study Host Cell-Parasite Interplay

#### **OPEN ACCESS**

#### Edited by:

Debora Decote-Ricardo, Universidade Federal Rural do Rio de Janeiro. Brazil

#### Reviewed by:

Fabienne Tacchini-Cottier, Université de Lausanne, Switzerland Leonardo Marques da Fonseca, Universidade Federal do Rio de Janeiro, Brazil

#### \*Correspondence:

Albert Descoteaux albert.descoteaux@iaf.inrs.ca Valéria M. Borges vborges@bahia.fiocruz.br

<sup>†</sup>These authors have contributed equally to this work.

<sup>‡</sup>These authors have contributed equally to this work and as senior author.

#### Specialty section:

This article was submitted to Microbial Immunology, a section of the journal Frontiers in Microbiology

Received: 13 December 2017 Accepted: 19 March 2018 Published: 05 April 2018

#### Citation:

Lázaro-Souza M, Matte C, Lima JB, Arango Duque G, Quintela-Carvalho G, de Carvalho Vivarini Á, Moura-Pontes S, Figueira CP, Jesus-Santos FH, Gazos Lopes U, Farias LP, Araújo-Santos T, Descoteaux A and Borges VM (2018) Leishmania infantum Lipophosphoglycan-Deficient Mutants: A Tool to Study Host Cell-Parasite Interplay. Front. Microbiol. 9:626. doi: 10.3389/fmicb.2018.00626

doi: 10.3389/imico.2018.00626 Leisimuniu against me

Milena Lázaro-Souza<sup>1,2†</sup>, Christine Matte<sup>3†</sup>, Jonilson B. Lima<sup>4</sup>, Guillermo Arango Duque<sup>3</sup>, Graziele Quintela-Carvalho<sup>1,2,5</sup>, Áislan de Carvalho Vivarini<sup>6</sup>, Sara Moura-Pontes<sup>2</sup>, Cláudio P. Figueira<sup>1</sup>, Flávio H. Jesus-Santos<sup>1,2</sup>, Ulisses Gazos Lopes<sup>6</sup>, Leonardo P. Farias<sup>1</sup>, Théo Araújo-Santos<sup>4</sup>, Albert Descoteaux<sup>3\*‡</sup> and Valéria M. Borges<sup>1,2\*‡</sup>

<sup>1</sup> Laboratory of Inflammation and Biomarkers, Gonçalo Moniz Institut, Oswaldo Cruz Foundation, Salvador, Brazil, <sup>2</sup> Department of Legal Medicine, Federal University of Bahia, Salvador, Brazil, <sup>3</sup> Institut National de la Recherche Scientifique –Institut Armand-Frappier, Laval, QC, Canada, <sup>4</sup> Center of Biological Sciences and Health, Federal University of Western of Bahia, Barreiras, Brazil, <sup>5</sup> Instituto Federal de Educação, Ciência e Tecnologia Baiano (IFBaiano), Alagoinhas, Brazil, <sup>6</sup> Laboratory of Molecular Parasitology, Carlos Chagas Filho Biophysics Institute, Center of Health Science, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

Lipophosphoglycan (LPG) is the major surface glycoconjugate of metacyclic Leishmania promastigotes and is associated with virulence in various species of this parasite. Here, we generated a LPG-deficient mutant of Leishmania infantum, the foremost etiologic agent of visceral leishmaniasis in Brazil. The L. infantum LPG-deficient mutant  $(\Delta lpq1)$  was obtained by homologous recombination and complemented via episomal expression of LPG1 ( $\Delta lpg1 + LPG1$ ). Deletion of LPG1 had no observable effect on parasite morphology or on the presence of subcellular organelles, such as lipid droplets. While both wild-type and add-back parasites reached late phase in axenic cultures, the growth of  $\Delta lpg1$  parasites was delayed. Additionally, the deletion of LPG1 impaired the outcome of infection in murine bone marrow-derived macrophages. Although no significant differences were observed in parasite load after 4 h of infection, survival of  $\Delta lpg1$  parasites was significantly reduced at 72 h post-infection. Interestingly, L. infantum LPG-deficient mutants induced a strong NF-kB-dependent activation of the inducible nitric oxide synthase (iNOS) promoter compared to wild type and  $\Delta lpg1 + LPG1$ parasites. In conclusion, the L. infantum  $\Delta lpg1$  mutant constitutes a powerful tool to investigate the role(s) played by LPG in host cell-parasite interactions.

Keywords: Lipophosphoglycan, Leishmania infantum, gene targeting, lipid droplets, macrophage

### INTRODUCTION

Lipophosphoglycan (LPG) is one of the most abundant components of *Leishmania* membranes (Turco and Descoteaux, 1992). In the course of parasite interaction with invertebrate hosts, LPG binds to the midgut epithelium of specific species of the sandfly vectors (Sacks et al., 2000), and protects parasites against the digestive enzymes present in the peritrophic matrix following blood feeding (Sacks and Kamhawi, 2001). In vertebrate hosts, LPG contributes to virulence by shielding *Leishmania* against the complement system (Spath et al., 2003) and by inhibiting phagolysosomal

biogenesis (Desjardins and Descoteaux, 1997; Vinet et al., 2009; Moradin and Descoteaux, 2012). Purified LPG has been considered as a pathogen-associated molecular pattern molecule (PAMP) that triggers Toll-like receptors (TLR) and is also known to interfere with pro-inflammatory and signaling pathways in host cells (Descoteaux et al., 1991; Descoteaux and Turco, 1993; Becker et al., 2003; de Veer et al., 2003; Kavoosi et al., 2009; Rojas-Bernabé et al., 2014; Tavares et al., 2014; Lima et al., 2017).

This complex glycolipid is organized in four domains: a conserved 1-O-alkyl-2-lyso-phosphatidyl(myo)inositol membrane anchor, a conserved diphosphoheptasaccharide core structure, a polymer of repeating phosphodisaccharide units (phosphoglycan or PG) carrying species-specific side chains and variable, often mannose-rich cap structures (Turco and Descoteaux, 1992; McConville and Ferguson, 1993). Although the biosynthesis of LPG has attracted considerable interest, to date only few enzymes and transporters involved in this process have been identified either biochemically, genetically, or both (Ryan et al., 1993; Descoteaux et al., 1995, 1998, 2002).

One of the key enzymes in the biosynthesis of LPG is LPG1, a putative galactofuranosyl transferase specifically involved in the synthesis of the LPG glycan core (Ryan et al., 1993). Consequently, parasites lacking the LPG1 gene  $(\Delta lpg1)$  express a truncated LPG without the PG domain; they nonetheless assemble and secrete other PG-containing molecules (Dermine et al., 2000; Späth et al., 2000). Both L. major and L. donovani require LPG1 for the establishment of infection within macrophages, as evidenced by the elimination of LPG1null mutants following phagocytosis; yet, restoration of LPG expression by genetic complementation restored the capacity to replicate within macrophages (Späth et al., 2000; Lodge et al., 2006). Interestingly, phosphoglycan synthesis does not seem to be an absolute requirement for virulence in all Leishmania species, since L. mexicana phosphoglycan-deficient parasites were found to be similarly virulent to their wild-type (WT) counterparts (Ilg et al., 1999, 2001; Ilg, 2000). This difference in LPG requirement for the establishment of infection within macrophages may be related to the fact that L. mexicana resides in large fusogenic communal vacuoles, as opposed to the non-fusogenic, tight individual vacuoles in which L. major and L. donovani replicate. The role played by LPG1 in L. infantum infectivity in mammals remains to be established.

This report describes the disruption of *LPG1* in *L. infantum*, the main etiological agent of visceral leishmaniasis in Brazil. While deletion of *LPG1* did not alter parasite morphology *in vitro* or the presence of subcellular organelles, e.g., lipid droplets (LD),  $\Delta lpg1$  parasites experienced distinct infection outcomes in comparison to WT parasites. Hence, the *L. infantum LPG1*-null strain described in the present study constitutes a powerful tool to investigate the role of LPG in host-parasite interactions.

#### **METHODS**

#### **Ethics Statement**

This study was carried out in accordance with the recommendations of Institutional Review Board for Animal Experimentation (CEUA), Gonçalo Moniz Institute, Fundação Oswaldo Cruz. The protocol was approved by the Institutional Review Board for Animal Experimentation (CEUA), Gonçalo Moniz Institute, Fundação Oswaldo Cruz (Protocol No. 021/2015).

#### Animals

Inbred male C57BL/6 mice, aged 6–8 weeks, were obtained from the animal care facility of the Gonçalo Moniz Institute, Fundação Oswaldo Cruz (IGM-FIOCRUZ, Bahia, Brazil).

# Targeted Deletion of the *LPG1* Gene and Complementation

The constructs for LPG1 (beta galactofuranosyl transferase) gene targeting were designed based on the L. infantum LPG1 gene sequence (GenBank accession No. GU233511). Homozygous LPG1-null mutants ( $\Delta lpg1$ ) were obtained using two targeting constructs (Figures 1A,B). For the NEO targeting construct, the entire LPG1 gene was amplified by PCR from L. infantum BH46 (MCAN/BR/89/BH46) DNA using Taq DNA polymerase (New England Biolabs) and oligodeoxynucleotides AD-358 (forward) (5'-gtacaagcttccatATGGCGCCGCCTCGCTG-3') and AD-359 (reverse) (5'-gctactcgagTTAGCTGGGGTCAACAG-3'). This fragment was digested with HindIII and XhoI, and then ligated with the *Hind*III-XhoI-digested pBluescript II SK<sup>-</sup> vector, yielding pBS-LPG1. The NEO resistance cassette from pLeishNeo (unpublished) was extracted with NotI and EcoRV, blunted and inserted in the MscI site of pBS-LPG1, within the LPG1 gene, yielding pBS-LPG1::NEO. For the HYG targeting construct, nucleotides 1-437 of the LPG1 gene were amplified by RT-PCR from L. infantum BH46 mRNA using oligodeoxynucleotides AD-53 (forward) (5'-cgggatccatATGGCGCCGCCTCGCTG-3') and AD-357 (reverse) (5'-ggaattcTCGGGGTGGTGAATG-3'). This fragment was digested with BamHI and EcoRI, and then ligated with the BamHI-EcoRI-digested pBluescript II SK<sup>-</sup> vector. A 467-bp fragment containing nucleotides 781-1,247 of the LPG1 ORF was amplified by PCR from L. infantum BH46 genomic DNA using oligodeoxynucleotides AD-355 (forward) (5'gcaagcttGGCATCTATTACACAGACCACAAGG-3') and AD-356 (reverse) (5'-caggtcgacTGGCAGCGAATGTTTTCACC-3'). This fragment was digested with *Hind*III and *Sal*I, then ligated with the same vector, downstream of the first LPG1 sequence, at the HindIII and SalI restriction sites. The HYG resistance cassette from pX63-HYG was excised with SalI and BamHI, blunted and inserted between the two LPG1 sequences, at the EcoRV restriction site, yielding pBS-LPG1::HYG. For genetic complementation of the  $\Delta lpg1$  mutant, the entire LPG1 ORF was amplified by PCR from L. infantum BH46 genomic DNA using Native Pfu polymerase (Stratagene, La Jolla, CA, USA) and oligodeoxynucleotides AD-358 (forward) (5'-gtacaagcttccatATGGCGCCGCCTCGCTG-3') and AD-359 (reverse) (5'-gctactcgagTTAGCTGGGGGTCAACAG-3'). This fragment was digested with HindIII and XhoI, and then ligated with the HindIII-XhoI-digested pBluescript II SK- vector, yielding pBSII-LPG1. The absence of mutations in the amplified LPG1 ORF was verified by Sanger sequencing (Génome Québec; GenBank accession No. GU233511). The LPG1 gene was then excised from pBSII-LPG1 with EcoRV and XhoI, blunted and



LPG1 gene and the targeting constructs. Arrows indicate gene orientation. (C) Western blot analysis of LPG expression in W1,  $\Delta lpg1$ , and  $\Delta lpg1 + LPG1$ promastigotes. Parasite lysates were probed with the anti-phosphoglycan (PG) antibody CA7AE, as described in Materials and Methods. (D) Confocal immunofluorescence analysis of WT,  $\Delta lpg1$  and  $\Delta lpg1 + LPG1$  parasites. Late log-phase promastigotes were adhered on Poly-L-Lysine-coated glass coverslips, fixed and incubated with DAPI to stain DNA (blue), and with the CA7AE antibody to visualize LPG and other Gal( $\beta$ 1,4)Man( $\alpha$ 1-PO4) repeating unit-containing PGs (red), as described in Materials and Methods. Fluorescence staining images merged with differential interference contrast (DIC) are shown in the lower panels. Scale bar, 5 µm.

ligated with the *Eco*RV-digested pLeishZeo vector (unpublished), yielding pLeishZeo-*LPG1*.

## Transfection and Selection of *L. infantum* $\Delta lpg1$ Promastigotes

Log-phase WT *L. infantum* BA262 (MCAN/BR/89/BA262) promastigotes were first electroporated with the purified *LPG1::HYG* targeting construct (excised as a 2.9-kb *XbaI-KpnI* fragment from pBS-*LPG1::HYG*) using 0.2 cm electroporation cuvettes, at 0.45 kV and a high capacitance of 500  $\mu$ F as previously described (Turco et al., 1994). Following electroporation, promastigotes were incubated for 24h in drug-free, complete M199 medium and subsequently grown in the presence of 50  $\mu$ g/mL Hygromycin B (Roche Diagnostics). To generate *LPG1*-null mutants, log-phase *lpg1*+/*HYG* heterozygous *L. infantum* BA262 promastigotes were electroporated with

purified *LPG1::NEO* targeting construct (excised as a 3.8-kb *Hind*III-*Kpn*I fragment from pBS-*LPG1::NEO*) and grown after 24 h in the presence of both 50 µg/mL Hygromycin B and 70 µg/mL G418 (Life Technologies). Absence of LPG in the resulting double drug-resistant  $\Delta lpg1$  promastigotes was verified by Western blot analysis and confocal immunofluorescence. To restore *LPG1* expression, log-phase *L. infantum* BA262  $\Delta lpg1$  cells were electroporated with pLeishZeo-*LPG1*. Complemented mutants ( $\Delta lpg1 + LPG1$ ) were selected with 80 µg/mL Zeocin (in addition to G418 and Hygromycin B at concentrations specified above) and verified by Western blotting and confocal immunofluorescence.

#### Parasite Cultures

L. infantum promastigotes were cultured in HOMEM medium supplemented with 10% inactivated Fetal Bovine Serum (FBS),

100 U/mL penicillin, 100 µg/mL streptomycin and 2 mM Lglutamine in 25 cm<sup>2</sup> flasks at 24°C until late log-phase. For *L. infantum* BA262  $\Delta lpg1$ , Hygromycin (50 µg/mL) and G418 (70 µg/mL) were added to the medium. For *LPG1*-complemented parasites ( $\Delta lpg1 + LPG1$ ), Hygromycin (50 µg/mL), G418 (70 µg/mL), and Zeocin (100 µg/mL) were added to the medium.

#### **Western Blotting**

Late log-phase promastigotes were washed with ice-cold PBS containing 1 mM Na<sub>3</sub>VO<sub>4</sub>, then lysed in 50 mM Tris-HCl pH 8, 150 mM NaCl and 1% Nonidet P-40, containing complete protease inhibitors (Roche Applied Science) and phosphatase inhibitors (1 mM Na<sub>3</sub>VO<sub>4</sub>, 50 mM NaF, 1.5 mM EGTA and 10 mM Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub>). Samples were sonicated briefly, and insoluble material was removed by centrifugation for 10 min at 4°C. Protein concentrations were determined using the Pierce BCA protein assay kit (Pierce). Proteins were separated by SDS-PAGE and then transferred to Hybond-LFP PVDF membranes (GE Healthcare Life Sciences) using a Trans-Blot SD Semi-Dry Transfer Cell apparatus (BioRad). Membranes were blocked with 5% BSA and incubated with the mouse monoclonal antibody CA7AE (MediMabs). For immunodetection, goat anti-mouse IgM Heavy Chain Secondary antibody conjugated with horseradish peroxidase (HRP), and enhanced chemiluminescence (ECL) detection reagents from GE Healthcare Life Sciences were used.

#### **Confocal Immunofluorescence Microscopy**

Late log-phase promastigotes were adhered on Poly-L-Lysinecoated glass coverslips (BD Biosciences, San Jose, CA) by centrifugation, fixed with 4% paraformaldehyde (Canemco and Marivac) for 20 min and simultaneously blocked and permeabilized with a solution of 0.1% Triton X-100, 1% BSA, 6% non-fat dry milk, 20% goat serum and 50% FBS for 20 min. The distribution of LPG and other PGs containing the  $Gal(\beta 1,4)Man(\alpha 1-PO_4)$  repeating unit epitope was visualized using the mouse monoclonal antibody CA7AE (MediMabs, 1:2,000) after 2 h incubation followed by Alexa Fluor 568 goat anti-mouse IgM (Molecular Probes) at 1:500 for 30 min incubation. Parasite nuclei were stained with DAPI (Molecular Probes) at 1:17,000. All steps were performed at room temperature. Coverslips were then mounted in Fluoromount-G (Interscience) and sealed with nail polish. Promastigotes were observed with a Plan APOCHROMAT 63x oil-immersion DIC 1.4 NA objective on a Zeiss LSM780 confocal microscope equipped with a 30 mW 405 nm diode laser, 25 mW 458/488/514 argon multiline laser, 20 mW DPSS 561 nm laser and 5 mW HeNe 633 nm laser, coupled to a Zeiss Axio Observer Z1. Images were acquired in plane scanning mode, and were minimally and equally processed using Carl Zeiss ZEN 2011 software.

#### **Electron Microscopy**

Late log-phase promastigotes were fixed with 2% paraformaldehyde plus 2.5% glutaraldehyde in 0.1 M phosphate buffer (pH 7.4) overnight at  $4^{\circ}$ C. Next, parasites were processed

for Transmission Electron Microscopy (TEM) by post-fixing in 1% osmium tetroxide (OsO<sub>4</sub>) plus 0.8% potassium ferricyanide in 0.1 M cacodylate buffer, pH 7.2, then dehydrated in acetone at increasing concentrations of 50, 70, 90, and 100% followed by processing for resin embedding (PolyBed 812, Polysciences). Sections were mounted on uncoated 200-mesh copper grids and viewed under a TEM microscope (JEOL JEM-1230). Alternatively, parasites were processed for Scanning Electron Microscopy (SEM) by first fixing as described above, then adhered on Poly-L-Lysine-coated glass coverslips and postfixed as described above. Samples were then submitted to critical point-drying under CO<sub>2</sub>, coated with a 20 nm-layer of gold particles and examined under SEM (JSM-6390LV, JEOL).

#### **Parasite Growth Curves**

Early log-phase promastigotes (1  $\times$  10<sup>5</sup>/ml) were cultured and the number of viable promastigotes was determined by daily direct counting performed in a Neubauer chamber.

#### Lipid Droplets Staining and Quantification

Late log-phase promastigotes were fixed with 3.7% formaldehyde and stained with osmium tetroxide. Cell morphology was observed, and LD were counted by light microscopy using a 100X objective lens in 50 consecutively scanned parasites (Araújo-Santos et al., 2014).

## Bone Marrow-Derived Macrophages (BMDM) Macrophage and Infection

Bone marrow-derived macrophages (BMDM) were obtained from C57BL/6 mice as previously described. Briefly, cells were collected from femurs and differentiated in RPMI 1640, 20% inactivated FBS, 30% L929 cell-conditioned media (LCCM), 2 mM L-glutamine, 100 U/mL Penicillin, and 100  $\mu$ g/mL Streptomycin at 36°C under 5% CO<sub>2</sub>. BMDMs were collected after 7 days and seeded on tissue culture plates in RPMI 1640 media, 10% inactivated FBS, 5% LCCM and 2 mM L-glutamine (Araújo-Santos et al., 2014).

Cells  $(2 \times 10^5)$  adhered on coverslips were infected with either WT,  $\Delta lpg1$ , or  $\Delta lpg1+LPG1$  parasites at a 10:1 multiplicity of infection (MOI). After 4 or 72 h of infection, coverslips were fixed and stained with DiffQuik (Wright-Giemsa). Intracellular parasites were counted under light microscopy to determine the infection index under each experimental condition (Araújo-Santos et al., 2014).

#### RAW 264.7 Cell Line, Culture, and Infection

The mouse macrophage leukemia cell line RAW 264.7 (TIB-71; American Type Culture Collection (ATCC), Manassas, VA, USA) was maintained in DMEM medium with high glucose (Vitrocell Embriolife, Campinas, SP, Brazil) supplemented with 10% heat-inactivated fetal bovine serum (Sigma-Aldrich, St. Louis, MO, USA), 100 U/ml penicillin and 100  $\mu$ g/ml streptomycin in an incubator at 37°C under 5% CO<sub>2</sub>. RAW 264.7 cells were infected with either WT,  $\Delta lpg1$  or  $\Delta lpg1+LPG1$  parasites at a 10:1 multiplicity of infection (MOI). After 4 or 8h of infection, cells were processed for quantitative RT-PCR. For the luciferase reporter assay, cultures were washed 2 h post-infection and analyzed 24 h later.

#### **RNA Extraction and RT-qPCR**

For real time quantitative polymerase chain reaction analysis, total RNA of control and infected RAW 264.7 cells (1 imes10<sup>6</sup> cells) was extracted using an Invitrap<sup>®</sup> Spin Cell RNA mini kit (STRACTEC Molecular GmbH, Berlin, Germany). RNA extracts (2 µg) were reverse transcribed into first-strand cDNA with ImProm-II (Promega) and oligo(dT) primers in accordance with manufacturer instructions. The following primer DNA sequences were used to determine iNOS mRNA levels: Forward 5'-CAGCTGGGCTGTACAAACCTT-3' and Reverse: 5'-CATTGGAAGTGAAGCGTTTCG- 3', while GAPDH mRNA levels were quantified using: Forward 5'-TGCACCACCAACTGCTTAGC-3' 5'and Reverse GGCATGGACTGTGGTCATGAG-3'. Amplicon specificity was carefully verified by the presence of a single melting temperature peak in dissociation curves calculated following RT-qPCR, which was performed via the Applied Biosystems StepOne<sup>TM</sup> detection system (Applied Biosystems) using GoTaq® qPCR Master Mix (Promega Corp., Madison, WI, USA). All RT-qPCR analyses were performed in triplicate. RT-qPCR data was normalized using GAPDH primers as an endogenous control. All gene expression ratios were calculated by the  $\Delta\Delta$ Ct method using StepOne software version 2.0 (Applied Biosystems).

## Transient Transfections and Luciferase Assays

To investigate NF-kB transcriptional activity, RAW 264.7 were plated in 48-well polystyrene plates (1  $\times$  10<sup>5</sup> cells per well) and transfected with 1 µg of the p6kB-Luc luciferase reporter construct (kindly provided by Dr. Patrick Baeuerle, Munich University) in the presence of LIPOFECTAMINE 2000 reagent (Invitrogen, Carlsbad, CA, USA). pTK-3XNS luciferase reporter construct was used to measure iNOS promoter activity, provided by Dr. David Geller (University of Pittsburgh, Pennsylvania, EUA). Luciferase activity was normalized using 40 ng of pRL-CMV plasmid (Promega Corp., Madison, WI, USA). Transfected cells were infected with either WT,  $\Delta lpg1$  or  $\Delta lpg1+LPG1$ parasites at a 10:1 MOI. After 24h of infection, cells were washed with PBS, lysed according to the Dual Luciferase System protocol (Promega Corp.), and analyzed in a GloMax<sup>®</sup>-Multi detection system (Promega Corp.). Positive controls consisting of cells stimulated with 1 µg/mL of LPS (Sigma-Aldrich) were used to induce the activation of iNOS gene expression.

#### **Statistical Analysis**

BMDM and RAW 264.7 cell infection assays were performed in triplicate, and each experiment was repeated at least three times. Data are presented as the mean and SE (standard error) of representative experiments, and GraphPad Prism 5.0 software (GraphPad Software) was used for data analysis. Means from different groups were compared by Oneway ANOVA and comparisons between two groups were performed using the Student Newman-Keuls post-test. Differences were considered statistically significant when  $p \le 0.05$ .

### RESULTS

# Generation of a *L. infantum Lpg1*-Null $\Delta lpg1$ Mutant

To generate a *L. infantum* LPG-defective ( $\Delta lpg1$ ) mutant, WT *L. infantum* BA262 promastigotes were transfected with the *LPG1* targeting constructs (**Figures 1A,B**). The resulting *HYG*and *NEO*-resistant  $\Delta lpg1$  parasites were transfected with a *LPG1* expression vector to generate add-back LPG-expressing parasites ( $\Delta lpg1 + LPG1$ ). Loss of LPG expression in the  $\Delta lpg1$ , was determined by comparing LPG levels in WT,  $\Delta lpg1$ , and  $\Delta lpg1 + LPG1L$ . *infantum* promastigotes by Western blot and by confocal immunofluorescence microscopy (**Figures 1C,D**). Together, these data indicate that the *LPG1* gene was successfully deleted in the  $\Delta lpg1$  mutants, resulting in the generation of a LPG-defective *L. infantum* mutant.

# *LPG1-Null* Mutants Retain *L. infantum* Viability and Morphology

To determine the effect of deleting LPG1 on parasite growth and morphology, axenic cultures of the three isolates were monitored and counted daily for 10 days until reaching late log phase. A delayed replication capability of the  $\Delta lpg1$  mutant parasites was noted in comparison to the WT and  $\Delta lpg1$  + LPG1 parasites (Figure 2A). Wild-type L. infantum presented regular growth for 7 days until reaching stationary phase, with a cell density of approximately  $3-4 \times 10^7$  parasites/ml, while the  $\Delta lpg1$  mutant reached the same phase approximately 3 days later, with a cell density of  $1-2 \times 10^7$  parasites/ml (Figure 2A). The  $\Delta lpg1 + LPG1$  mutants presented an intermediate growth profile, reaching stationary phase shortly after the WT parasites. Area Under the Curve (AUC) analysis of the growth curve revealed a significant difference only when comparing WT and  $\Delta lpg1$  parasites (p < 0.05), yet no differences were observed between WT and  $\Delta lpg1 + LPG1$  mutants (Figure 2B). Upon reaching stationary phase, parasites were examined by electron microscopy to assess the presence of morphological alterations. Under both SEM and TEM, no alterations in morphology (Figure 2C) or in ultrastructural characteristics (Figure 3A) were detected among WT,  $\Delta lpg1$ , and  $\Delta lpg1 +$ LPG1 promastigotes. In addition, the absence of LPG1 had no impact on the number of lipid bodies present within the parasites (Figure 3A). Hence, whereas the LPG1 gene had a limited impact on L. infantum promastigotes proliferation, it did not significantly alter morphological features of these parasites (Figure 3).

# *Lpg1-Null* Mutants Exhibit Limited Survival in Macrophages

To evaluate differences in parasite survival among WT and transgenic parasites *in vitro*, BMDMs were infected for 4 or







72 h. No differences were observed between the *Leishmania* parasites after 4 h of infection (**Figure 4**). However, at 72 h post-infection, WT parasites survived more efficiently than the  $\Delta lpg1$  mutant. Expression of *LPG1* in the  $\Delta lpg1$  mutant partially

restored its capacity to survive and replicate within macrophages. These data reinforce the importance of LPG as a virulence factor in the successful maintenance of *Leishmania infantum* infection.



**FIGURE 4** | *LPG1* promotes intraphagosomal survival in infected macrophages. C57BL/6 BMDMs were infected with WT,  $\Delta lpg1$  and  $\Delta lpg1 + LPG1$  promastigotes as described under Materials and Methods. At 4 and 72 h post-infection, cells were fixed and Giemsa-stained. **(A)** Micrographs of infected BMDMs at 4 and 72 h post-infection. The infection index **(B)** and amastigotes per macrophage **(C)** were quantified by light microscopy. Arrows point to amastigotes inside a parasitophorous vacuole. Original magnification × 1000. Statistical differences were evaluated using the Student Newman-Keuls test. \*\*\*p < 0.001 compared to WT or between groups.

#### *LPG1-Null* Mutants Induce NF-κB-Dependent iNOS Expression in Macrophages

To assess the impact of LPG on the expression of inducible nitric oxide synthase by host cells, we first analyzed iNOS transcript levels in RAW 264.7 cells infected with WT or transgenic parasites. The  $\Delta lpg1$  mutant induced a robust (3.5fold increase) expression of iNOS compared to WT and  $\Delta lpg1 +$ LPG1 promastigotes (Figure 5A). We next performed luciferase reporter assays to characterize the modulation of the iNOS promoter by L. infantum promastigotes. RAW 264.7 cells were transiently transfected with either the iNOS promoter reporter construct pTK-3XNS or the NF-κB consensus luciferase reporter construct (p6kB-Luc) prior to infection with either L. infantum WT,  $\Delta lpg1$  or  $\Delta lpg1 + LPG1$ , or stimulation with LPS. As shown in Figures 5B,C, LPG-deficient promastigotes induced stronger activation of the iNOS promoter and of the NF-KB reporter. Collectively, these findings indicate that LPG contributes to the evasion of iNOS expression by L. infantum promastigotes.

#### DISCUSSION

Previous studies using purified LPG were important to unravel its impact on the activation of the immune system. Although purified LPG from different species can activate the release of inflammatory mediators, understanding the role of this response in the context of infection remains a challenge. While some groups have characterized Old World *LPG1*defective *Leishmania* species, the behavior of these parasites when compared to their WT counterparts varies depending on the species under study (Spath et al., 2003; Capul et al., 2007; Forestier et al., 2014). Here, we generated for the first time a LPG-deficient mutant of *L. infantum*, a New World species cluster.

A comparison of parasite growth between axenic cultures containing each of the three isolates showed that the deletion of *LPG1* resulted in a delayed capability of the  $\Delta lpg1$  mutant parasites to replicate in comparison to cultures of WT and  $\Delta lpg1 + LPG1$  parasites. While the deletion of the *LPG1* gene had a limited impact on *L. infantum* promastigote proliferation, no significant morphological or ultrastructural alterations were seen in these parasites, indicating that targeting the *LPG1* gene does not interfere with the intrinsic cell biology of *L. infantum*.

Recently, we have demonstrated that intact LPG from L. infantum promastigotes, but not its glycan and lipid moieties, induced a range of pro-inflammatory responses, including prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) and nitric oxide (NO) release, increased LD formation, and inducible nitric oxide synthase (iNOS) and cyclooxygenase-2 expression (Lima et al., 2017). Consequently, a limitation of using purified LPG is that the physiological conditions present in host cell-parasite interactions are not accurately replicated. LDs are key cytoplasmic organelles involved in production of lipid mediators and pro-inflammatory cytokines in mammalian cells (Bozza et al., 2011). Several intracellular pathogens, including Leishmania, take advantage of LD formation in host cells (Rabhi et al., 2016). Moreover, LDs have also been described in trypanosomatids in association with arachidonic acid metabolism (Araújo-Santos et al., 2014). Our group previously reported an increase in LD formation during L. infantum metacyclogenesis, as well as in the intracellular amastigote form (Araújo-Santos et al., 2014). Here, we showed that the absence of the LPG1 gene in L. infantum did not alter the biogenesis of LDs. In addition, our previous findings showed that parasite-derived  $PGF_{2\alpha}$  produced inside LDs plays a critical role during macrophage infection (Araújo-Santos et al., 2014). We fully intend to comprehensively investigate the potential influence of LPG1 on the release of  $PGF_{2\alpha}$  in infected macrophages using this novel L. infantum  $\Delta lpg1$  mutant.

In *L. major* and *L. donovani*, the specific loss of LPG through the ablation of *LPG1* galactofuranosyl transferase strongly impairs the ability of parasites to survive within the sandfly host, as well as to establish infection in mammalian macrophages and in mice (Sacks et al., 2000; Spath et al., 2003; Secundino et al., 2010). Hence, in these species, LPG impairs the microbicidal mechanisms associated with the biogenesis of phagolysosomes, including assembly of the NADPH oxidase and recruitment of the v-ATPase (Lodge et al.,



control. ##p < 0.0001, #p < 0.001 and #p < 0.05 compared to WT. P < 0.001 compared with control group Student Newman-Keuls post-test.

2006; Vinet et al., 2009). Our results indicate that, similarly to these species, L. infantum LPG1 is required for replication within macrophages. Whether LPG contributes to the ability of L. infantum to successfully infect macrophages through the impairment of phagolysosomal biogenesis remains to be investigated. Interestingly, we observed that our  $\Delta lpg1$  mutant parasites induced robust NF-kB-dependent iNOS expression compared to parental WT L. infantum promastigotes. This seems to suggest that the reduced survival of  $\Delta lpg1$  mutants in mouse macrophages may be related to higher levels of iNOS, which is responsible for the generation of leishmanicidal nitric oxide (Coelho-Finamore et al., 2011; Passero et al., 2015). Further study will involve investigating the contribution of nitric oxide production with respect to the reduced ability of  $\Delta lpg1$  mutants to survive within macrophages. The underlying mechanism by which  $\Delta lpg1$  mutant parasites induce high levels of NF- $\kappa B$ activation remains unknown, and thus represents an additional aspect of host cell-parasite interplay that we intend to further investigate.

With regard to the partial restoration of the WT phenotype observed in  $\Delta lpg1 + LPG1$  parasites, it has been welldocumented that complemented parasites commonly do not fully recover virulence. The inappropriate regulation of *LPG1* expression by the episomal vector may be a possible explanation for this observation (Späth et al., 2000; Spath et al., 2003; Joshi et al., 2002). A previous study demonstrated that *L. major LPG1*deficient mutant promastigotes present an attenuated virulence phenotype, as evidenced by the delayed formation of lesions *in vivo* (Späth et al., 2000). In addition, this delay was associated with a 100-fold decrease in parasite survival within macrophages *in vitro*. The data presented herein are consistent with these results, as well as with other reports in the literature (Privé and Descoteaux, 2000; Sacks et al., 2000; Zhang et al., 2004) propounding LPG as a virulence factor.

Taken together, the present findings support the importance of creating LPG-deficient mutants in various *Leishmania* spp. as a unique tool to investigate the specific impact and contribution of this abundant virulence factor in the complex host cell-*Leishmania* interplay. Hence, we are currently conducting studies to compare the responses of various immune cells to live *L. infantum* promastigotes in the presence or absence of surfaceexpressed LPG, since we feel it is important to thoroughly characterize these isolates to obtain a more comprehensive understanding regarding the role of *L. infantum LPG1* in future *in vitro* and *in vivo* studies.

### **AUTHOR CONTRIBUTIONS**

ML-S, CM, JL, GA, GQ-C, AC, SM-P, UG, LF, TA-S, AD, and VB conceived and designed the study, contributed to the data analysis, drafted, and revised the manuscript. ML-S, CM, JL, GA, GQ-C, AC, SM-P, CF, FJ-S, and TA-S performed the experiments. ML-S, CM, JL, LF, TA-S, AD, and VB wrote and revised the manuscript. All authors read and approved the final version of this manuscript.

### FUNDING

This work was supported by the Brazilian National Research Council (CNPq), (Grant number: 400135/2014-0 to VB), Fundação de Amparo à Pesquisa do Estado da Bahia (FAPESB) (Grant number: 04/2015 to VB), Science without Borders (SwB) (to VB and AD) and the Canadian Institutes of Health Research (CIHR) (grant MOP-125990 to AD). VB and UG are senior investigator funded by CNPq. AD is the holder of the Canada Research Chair on the Biology of intracellular parasitism. GA was partially supported by a CIHR Frederick Banting and Charles Best Doctoral Award. AC: received a fellowship from CNPq. The funders had no role in study design, data collection or analysis, the decision to publish, or preparation of the manuscript.

### ACKNOWLEDGMENTS

The authors thank Andrezza Souza for technical and logistics support. The authors would like to thank Andris K. Walter for English language revision and manuscript copyediting assistance.

#### REFERENCES

- Araújo-Santos, T., Rodríguez, N. E., Moura-Pontes, S., Dixt, U. G., Abánades, D. R., Bozza, P. T., et al. (2014). Role of prostaglandin F2α production in lipid bodies from Leishmania infantum chagasi: insights on virulence. J. Infect. Dis. 210, 1951–1961. doi: 10.1093/infdis/jiu299
- Becker, I., Salaiza, N., Aguirre, M., Delgado, J., Carrillo-Carrasco, N., Kobeh, L. G., et al. (2003). Leishmania lipophosphoglycan (LPG) activates NK cells through toll-like receptor-2. *Mol. Biochem. Parasitol.* 130, 65–74. doi: 10.1016/S0166-6851(03)00160-9
- Bozza, P. T., Bakker-Abreu, I., Navarro-Xavier, R. A., and Bandeira-Melo, C. (2011). Lipid body function in eicosanoid synthesis: an update. *Prostaglandins Leukot Essent Fat Acids* 85, 205–213. doi: 10.1016/j.plefa.2011. 04.020
- Capul, A. A., Hickerson, S., Barron, T., Turco, S. J., and Beverley, S. M. (2007). Comparisons of mutants lacking the golgi UDP-galactose or GDP-mannose transporters establish that phosphoglycans are important for promastigote but not amastigote virulence in Leishmania major. *Infect. Immun.* 75, 4629–4637. doi: 10.1128/IAI.00735-07
- Coelho-Finamore, J. M., Freitas, V. C., Assis, R. R., Melo, M. N., Novozhilova, N., Secundino, N. F., et al. (2011). Leishmania infantum: Lipophosphoglycan intraspecific variation and interaction with vertebrate and invertebrate hosts. *Int. J. Parasitol.* 41, 333–342. doi: 10.1016/j.ijpara.2010.10.004
- Dermine, J. F., Scianimanico, S., Priv,é, C., Descoteaux, A., and Desjardins, M. (2000). Leishmania promastigotes require lipophosphoglycan to actively modulate the fusion properties of phagosomes at an early step of phagocytosis. *Cell. Microbiol.* 2, 115–126. doi: 10.1046/j.1462-5822.2000.00037.x
- Descoteaux, A., Avila, H. A., Zhang, K., Turco, S. J., and Beverley, S. M. (2002). Leishmania LPG3 encodes a GRP94 homolog required for phosphoglycan synthesis implicated in parasite virulence but not viability. *EMBO J.* 21, 4458–4469. doi: 10.1093/emboj/cdf447
- Descoteaux, A., Luo, Y., Turco, S. J., and Beverley, S. M. (1995). A specialized pathway affecting virulence glycoconjugates of Leishmania. *Science* 269, 1869–1872. doi: 10.1126/science.7569927
- Descoteaux, A., Mengeling, B. J., Beverley, S. M., and Turco, S. J. (1998). Leishmania donovani has distinct mannosylphosphoryltransferases for the initiation and elongation phases of lipophosphoglycan repeating unit biosynthesis. Mol. Biochem. Parasitol. 94, 27–40. doi: 10.1016/S0166-6851(98)00047-4
- Descoteaux, A., and Turco, S. J. (1993). The lipophosphoglycan of Leishmania and macrophage protein kinase C. *Parasitol. Today* 9, 468–471. doi: 10.1016/0169-4758(93)90105-O
- Descoteaux, A., Turco, S. J., Sacks, D. L., and Matlashewski, G. (1991). Leishmania donovani lipophosphoglycan selectively inhibits signal transduction in macrophages. J Immunol 146, 2747–2753.
- Desjardins, M., and Descoteaux, A. (1997). Inhibition of phagolysosomal biogenesis by the Leishmania lipophosphoglycan. J. Exp. Med. 185, 2061–2068. doi: 10.1084/jem.185.12.2061
- de Veer, M. J., Curtis, J. M., Baldwin, T. M., DiDonato, J. A., Sexton, A., McConville, M. J., et al. (2003). MyD88 is essential for clearance of Leishmania major: possible role for lipophosphoglycan and Toll-like receptor 2 signaling. *Eur. J. Immunol.* 33, 2822–2831. doi: 10.1002/eji.200324128
- Forestier, C. L., Gao, Q., and Boons, G. J. (2014). Leishmania lipophosphoglycan: how to establish structure-activity relationships for this highly complex and multifunctional glycoconjugate? *Front. Cell. Infect. Microbiol.* 4:193. doi: 10.3389/fcimb.2014.00193
- Ilg, T. (2000). Lipophosphoglycan is not required for infection of macrophages or mice by *Leishmania mexicana*. *EMBO J.* 19, 1953–1962. doi: 10.1093/emboj/19.9.1953
- Ilg, T., Demar, M., and Harbecke, D. (2001). Phosphoglycan Repeat-deficient Leishmania mexicana parasites remain infectious to macrophages and mice. J. Biol. Chem. 276, 4988–4997. doi: 10.1074/jbc.M008030200
- Ilg, T., Montgomery, J., Stierhof, Y. D., and Handman, E. (1999). Molecular cloning and characterization of a novel repeat-containing Leishmania major gene, ppg1, that encodes a membrane-associated form of proteophosphoglycan with a putative glycosylphosphatidylinositol anchor. J. Biol. Chem. 274, 31410–31420. doi: 10.1074/jbc.274.44.31410

- Joshi, P. B., Kelly, B. L., Kamhawi, S., Sacks, D. L., and McMaster, W. R. (2002). Targeted gene deletion in Leishmania major identifies leishmanolysin (GP63) as a virulence factor. *Mol. Biochem. Parasitol.* 120, 33–40. doi: 10.1016/S0166-6851(01)00432-7
- Kavoosi, G., Ardestani, S. K., and Kariminia, A. (2009). The involvement of TLR2 in cytokine and reactive oxygen species (ROS) production by PBMCs in response to Leishmania major phosphoglycans (PGs). *Parasitology* 136, 1193–1199. doi: 10.1017/S0031182009990473
- Lima, J. B., Araújo-Santos, T., Lázaro-Souza, M., Carneiro, A. B., Ibraim, I. C., Jesus-Santos, F. H., et al. (2017). Leishmania infantum lipophosphoglycan induced-Prostaglandin E2 production in association with PPAR-γ expression via activation of Toll like receptors-1 and 2. *Sci. Rep.* 7:14321. doi: 10.1038/s41598-017-14229-8
- Lodge, R., Diallo, T. O., and Descoteaux, A. (2006). Leishmania donovani lipophosphoglycan blocks NADPH oxidase assembly at the phagosome membrane. Cell. Microbiol. 8, 1922–1931. doi: 10.1111/j.1462-5822.2006.00758.x
- McConville, M. J., and Ferguson, M. A. (1993). The structure, biosynthesis and function of glycosylated phosphatidylinositols in the parasitic protozoa and higher eukaryotes. *Biochem. J.* 294, 305–324. doi: 10.1042/bj2940305
- Moradin, N., and Descoteaux, A. (2012). Leishmania promastigotes: building a safe niche within macrophages. *Front. Cell. Infect. Microbiol.* 2:121. doi: 10.3389/fcimb.2012.00121
- Passero, L. F., Assis, R. R., da Silva, T. N., Nogueira, P. M., Macedo, D. H., Pessoa, N. L., et al. (2015). Differential modulation of macrophage response elicited by glycoinositolphospholipids and lipophosphoglycan from Leishmania (Viannia) shawi. *Parasitol. Int.* 64, 32–35. doi: 10.1016/j.parint.2015.01.006
- Privé, C., and Descoteaux, A. (2000). Leishmania donovani promastigotes evade the activation of mitogen-activated protein kinases p38, c-Jun N-terminal kinase, and extracellular signal-regulated kinase-1/2 during infection of naive macrophages. Eur. J. Immunol. 30, 2235–2244. doi: 10.1002/1521-4141(2000)30:8<2235::AID-IMMU2235&gt;3.0.CO;2-9
- Rabhi, S., Rabhi, I., Trentin, B., Piquemal, D., Regnault, B., Goyard, S., et al. (2016). Lipid droplet formation, their localization and dynamics during leishmania major macrophage infection. *PLoS ONE* 11:e0148640. doi: 10.1371/journal.pone.0148640
- Rojas-Bernabé, A., Garcia-Hernández, O., Maldonado-Bernal, C., Delegado-Dominguez, J., Ortega, E., Gutiérrez-Kobeh, L., et al. (2014). *Leishmania mexicana* lipophosphoglycan activates ERK and p38 MAP kinase and induces production of proinflammatory cytokines in human macrophages through TLR2 and TLR4. *Parasitology* 141, 788–800. doi: 10.1017/S00311820130 02187
- Ryan, K. A., Garraway, L. A., Descoteaux, A., Turco, S. J., and Beverley, S. M. (1993). Isolation of virulence genes directing surface glycosylphosphatidylinositol synthesis by functional complementation of Leishmania. *Proc. Natl. Acad. Sci. U.S.A.* 90, 8609–8613. doi: 10.1073/pnas.90. 18.8609
- Sacks, D., and Kamhawi, S. (2001). Molecular aspects of parasite-vector and vector-host interactions in Leishmaniasis. Annu. Rev. Microbiol. 55, 453–483. doi: 10.1146/annurev.micro.55.1.453
- Sacks, D. L., Modi, G., Rowton, E., Späth, G., Epstein, L., Turco, S. J., et al. (2000). The role of phosphoglycans in Leishmania-sand fly interactions. *Proc. Natl. Acad. Sci. U.S.A.* 97, 406–411. doi: 10.1073/pnas.97.1.406
- Secundino, N., Kimblin, N., Peters, N. C., Lawyer, P., Capul, A. A., Beverley, S. M., et al. (2010). Proteophosphoglycan confers resistance of Leishmania major to midgut digestive enzymes induced by blood feeding in vector sand flies. *Cell. Microbiol.* 12, 906–918. doi: 10.1111/j.1462-5822.2010.01439.x
- Späth, G. F., Epstein, L., Leader, B., Singer, S. M., Avila, H. A., Turco, S. J., et al. (2000). Lipophosphoglycan is a virulence factor distinct from related glycoconjugates in the protozoan parasite Leishmania major. *Proc. Natl. Acad. Sci. U.S.A.* 97, 9258–9263. doi: 10.1073/pnas.160257897
- Spath, G. F., Garraway, L. A., Turco, S. J., and Beverley, S. M. (2003). The role(s) of lipophosphoglycan (LPG) in the establishment of Leishmania major infections in mammalian hosts. *Proc. Natl. Acad. Sci.U.S.A.* 100, 9536–9541. doi: 10.1073/pnas.1530604100
- Tavares, N. M., Araújo-Santos, T., Afonso, L., Nogueira, P. M., Lopes, U. G., Soares, R. P., et al. (2014). Understanding the mechanisms controlling

Leishmania amazonensis infection in vitro: the role of LTB4 derived from human neutrophils. J. Infect. Dis. 210, 656–666. doi: 10.1093/infdis/jiu158

- Turco, S., Descoteaux, A., Ryan, K., Garraway, L., and Beverley, S. (1994). Isolation of virulence genes directing GPI synthesis by functional complementation of Leishmania. *Brazilian J. Med. Biol. Res.* 27, 133–138.
- Turco, S. J., and Descoteaux, A. (1992). The lipophosphoglycan of leishmania parasites. Annu. Rev. Microbiol. 46, 65–94. doi: 10.1146/annurev.mi.46.100192.000433
- Vinet, A. F., Fukuda, M., Turco, S. J., and Descoteaux, A. (2009). The Leishmania donovani lipophosphoglycan excludes the vesicular proton-ATPase from phagosomes by impairing the recruitment of Synaptotagmin, V. PLoS Pathog. 5:e1000628. doi: 10.1371/journal.ppat.1000628
- Zhang, K., Barron, T., Turco, S. J., and Beverley, S. M. (2004). The LPG1 gene family of Leishmania major. *Mol. Biochem. Parasitol.* 136, 11–23. doi: 10.1016/j.molbiopara.2004.02.012

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The reviewer LM declared a shared affiliation, with no collaboration, with two of the authors, AC and UG, to the handling Editor.

Copyright © 2018 Lázaro-Souza, Matte, Lima, Arango Duque, Quintela-Carvalho, de Carvalho Vivarini, Moura-Pontes, Figueira, Jesus-Santos, Gazos Lopes, Farias, Araújo-Santos, Descoteaux and Borges. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.