



**HAL**  
open science

## **A Seroepidemiological Study of Serogroup A Meningococcal Infection in the African Meningitis Belt**

Olivier Manigart, Caroline Trotter, Helen Findlow, Abraham Aseffa, Wude Mihret, Tesfaye Moti Demisse, Biruk Yeshitela, Isaac Osei, Abraham Hodgson, Stephen Laryea Quaye, et al.

### ► To cite this version:

Olivier Manigart, Caroline Trotter, Helen Findlow, Abraham Aseffa, Wude Mihret, et al.. A Seroepidemiological Study of Serogroup A Meningococcal Infection in the African Meningitis Belt. PLoS ONE, 2016, 11 (2), pp.e0147928. <10.1371/journal.pone.0147928>. <pasteur-03238501>

**HAL Id: pasteur-03238501**

**<https://riip.hal.science/pasteur-03238501v1>**

Submitted on 27 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

RESEARCH ARTICLE

# A Seroepidemiological Study of Serogroup A Meningococcal Infection in the African Meningitis Belt

Olivier Manigart<sup>1,2</sup>, Caroline Trotter<sup>3</sup>, Helen Findlow<sup>4</sup>, Abraham Assefa<sup>5</sup>, Wude Mihret<sup>5</sup>, Tesfaye Moti Demisse<sup>5</sup>, Biruk Yeshitela<sup>5</sup>, Isaac Osei<sup>6</sup>, Abraham Hodgson<sup>6</sup>, Stephen Laryea Quaye<sup>6</sup>, Samba Sow<sup>7</sup>, Mamadou Coulibaly<sup>7</sup>, Kanny Diallo<sup>7</sup>, Awa Traore<sup>7</sup>, Jean-Marc Collard<sup>8</sup>, Rahamatou Moustapha Boukary<sup>8</sup>, Oumarou Djermakoye<sup>8</sup>, Ali Elhaji Mahamane<sup>8</sup>, Jean-François Jusot<sup>8</sup>, Cheikh Sokhna<sup>9</sup>, Serge Alavo<sup>9</sup>, Souleymane Doucoure<sup>9</sup>, El Hadj Ba<sup>9</sup>, Mariétou Dieng<sup>9</sup>, Aldiouma Diallo<sup>9</sup>, Doumagoum Moto Daugla<sup>10</sup>, Babatunji Omotara<sup>11</sup>, Daniel Chandramohan<sup>1</sup>, Musa Hassan-King<sup>1</sup>, Maria Nascimento<sup>1</sup>, Arouna Woukeu<sup>1</sup>, Ray Borrow<sup>4</sup>, James M. Stuart<sup>1</sup>, Brian Greenwood<sup>1\*</sup>

**1** Faculty of Infectious Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom, **2** Medical Research Council Unit, Fajara, The Gambia, **3** Department of Veterinary Medicine, University of Cambridge, Cambridge, United Kingdom, **4** Public Health England Vaccine Evaluation Unit, Manchester, United Kingdom, **5** Armauer Hansen Research Institute, Addis Ababa, Ethiopia, **6** Navrongo Health Research Centre, Navrongo, Ghana, **7** Centre pour les Vaccins en Développement, Bamako, Mali, **8** Centre de Recherche Médicale et Sanitaire (CERMES), Niamey, Niger, **9** Institut de Recherche pour le Développement, Dakar, Senegal, **10** Centre de Support en Santé International (CSSI), N'Djamena, Chad, **11** Department of Community Medicine, University of Maiduguri, Maiduguri, Nigeria

\* [brian.greenwood@lshtm.ac.uk](mailto:brian.greenwood@lshtm.ac.uk)



CrossMark  
click for updates

**OPEN ACCESS**

**Citation:** Manigart O, Trotter C, Findlow H, Assefa A, Mihret W, Moti Demisse T, et al. (2016) A Seroepidemiological Study of Serogroup A Meningococcal Infection in the African Meningitis Belt. PLoS ONE 11(2): e0147928. doi:10.1371/journal.pone.0147928

**Editor:** I. King Jordan, Georgia Institute of Technology, UNITED STATES

**Received:** September 14, 2015

**Accepted:** January 11, 2016

**Published:** February 12, 2016

**Copyright:** © 2016 Manigart et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Data are available from the Institutional Data Access database <http://datacompass.lshtm.ac.uk/> (DOI: <http://dx.doi.org/10.17037/DATA.34>).

**Funding:** This work was supported by Wellcome Trust (grant no. 086546/Z/08/Z), [www.wellcome.ac.uk](http://www.wellcome.ac.uk); Bill & Melinda Gates Foundation (grant no. 51251), [www.gatesfoundation.org](http://www.gatesfoundation.org). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Abstract

The pattern of epidemic meningococcal disease in the African meningitis belt may be influenced by the background level of population immunity but this has been measured infrequently. A standardised enzyme-linked immunosorbent assay (ELISA) for measuring meningococcal serogroup A IgG antibodies was established at five centres within the meningitis belt. Antibody concentrations were then measured in 3930 individuals stratified by age and residence from six countries. Seroprevalence by age was used in a catalytic model to determine the force of infection. Meningococcal serogroup A IgG antibody concentrations were high in each country but showed heterogeneity across the meningitis belt. The geometric mean concentration (GMC) was highest in Ghana (9.09 µg/mL [95% CI 8.29, 9.97]) and lowest in Ethiopia (1.43 µg/mL [95% CI 1.31, 1.57]) on the margins of the belt. The force of infection was lowest in Ethiopia ( $\lambda = 0.028$ ). Variables associated with a concentration above the putative protective level of 2 µg/mL were age, urban residence and a history of recent vaccination with a meningococcal vaccine. Prior to vaccination with the serogroup A meningococcal conjugate vaccine, meningococcal serogroup A IgG antibody concentrations were high across the African meningitis belt and yet the region remained susceptible to epidemics.

**Competing Interests:** HF and RB report performing contract research on behalf of Public Health England for Novartis Vaccines and Diagnostics, Baxter Biosciences, GlaxoSmithKline, Pfizer, Sanofi Pasteur and Serum Institute of India. CT reports receiving a consulting payment from GlaxoSmithKline. All other authors report no conflict of interest. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials.

## Introduction

Epidemics of meningococcal disease have occurred at irregular intervals across the Sahelian and sub-Saharan regions of Africa, the African meningitis belt, for over 100 years.[1] However, despite many years of research it is still not known why epidemics occur at a particular place at any specific time. An important factor is likely to be the background level of immunity of the population when faced with a potentially epidemic strain. It is known that protective immunity to *Neisseria meningitidis* can be induced by meningococcal carriage,[2] infection with other non-pathogenic *Neisseria* species, such as *N. lactamica*[3] and possibly by other bacteria.[4,5] There is also some evidence that background immunity may be impaired with infection by other bacteria that induce blocking antibodies.[6] The immune response to meningococcal polysaccharide and conjugate vaccines has been studied in the African meningitis belt[7–10] on several occasions but there have been few studies of population levels of antibody to *N. meningitidis* in the African meningitis belt.[11,12] Therefore, we have undertaken a study of community levels of serogroup-specific IgG antibody to *N. meningitidis* serogroup A (NmA) in six countries in the African meningitis belt before the introduction of the serogroup A conjugate vaccine, MenAfriVac™, to investigate heterogeneity in the level of exposure across the meningitis belt and to use age specific antibody titres to measure the force of infection.[13] To ensure that patterns of antibody could be compared across sites, we implemented standardised methods supported by careful quality control.

## Materials and Methods

### Study population

Cross-sectional meningococcal carriage surveys were conducted in seven countries across the meningitis belt during the period July 1<sup>st</sup> 2010 to July 31<sup>st</sup> 2012 as described previously.[14] Ethical approval for the study was obtained from the London School of Hygiene & Tropical Medicine and from an appropriate committee from each African centre. Written, informed consent for study participation was obtained from adults and for the children under their care. Written informed assent was also obtained from participants aged 12 years or more. Oral assent was obtained from younger children.

Subjects were selected randomly from within populations which were part of a routine demographic survey system (DHSS) or in which a census had been performed recently. The study population was recruited from urban and rural populations and stratified into four age groups: < 5 years, 5–14 years, 15–29 years and 30 years or older.

Subjects were asked if they had received a meningitis vaccine in the previous six months. Approximately a year before the survey, a vaccination campaign with an A + C polysaccharide vaccine had been conducted in the study area in Senegal and also in part of the urban study area in Niger.[15] None of the study populations had been vaccinated with MenAfriVac™ at the time of the survey.

Blood samples were collected from the first 100 subjects surveyed within each of the four age bands in both urban and rural study sites, giving an overall target of 800 samples per country. This target was achieved, or nearly achieved, except in Senegal where there was some resistance to the collection of blood samples. A 5 mL sample was collected, serum separated within six hours of collection and then stored at -20°C until assayed.

### Enzyme Linked Immuno-Sorbent Antibody (ELISA) assay

An internationally standardised ELISA, as used at the Vaccine Evaluation Unit (VEU), Public Health England, Manchester, UK was transferred to each of the MenAfriCar centres.

Concentrations of IgG antibody against *N. meningitidis* serogroup A polysaccharide were obtained through a classical sandwich assay ELISA as described previously,[16] except that the standard reference serum CDC1992 was used as the quantification reference and that a monoclonal-PAN anti-human IgG Fc labelled with horseradish peroxidase (HRP)(Hybridoma Reagent Laboratory, Baltimore, MD) was used as conjugate. The lower limit of quantification (LLQ) of the meningococcal serogroup A ELISA was 0.19 µg/mL. Any value lower than the LLQ was assigned a value of 0.095 µg/mL for computational purposes.

## Standardisation of the assay and quality control

To ensure comparability of assay methods between centres, two training sessions were held in Manchester, UK and in Bamako, Mali at the start of the project. Subsequently, approximately 50 samples obtained during a pilot study conducted in each country were selected to allow cross-validation of the technique between each of the centres and the VEU, Manchester. After repeated testing of the 50 samples and adjustment of the technique to ensure that the results obtained fell within a defined range of the results obtained at the VEU, authorization was given to start testing the samples obtained during the cross-sectional survey. During the course of testing, monitoring of key values was performed by a resident scientist: two values representative of the standard curves (average of the duplicate values of the second higher concentration point and midpoint of the slope) as well as the calculated concentration of the local positive control were plotted routinely on Levey-Jennings charts. Regular review of these data was undertaken by the MenAfriCar laboratory manager and advice provided on adjustment of the technique when problems arose, for example detection of degradation in the anti-IgG conjugate used in the assay. It was not possible to complete cross-validation in Nigeria due to increasing insecurity and samples collected in Nigeria were tested in Mali. The laboratory in Chad did not reach the required standard in the validation assay to progress to testing of cross-sectional samples.

## Statistical methods

Pearson's correlation coefficient ( $\rho$ ) was used to compare the results produced at the VEU and by each centre during the validation exercise. The acceptance criterion for passing the cross-validation test was  $\rho$  greater than or equal to 0.9. In addition, Lin's concordance coefficient of correlation, ( $\rho_C$ ), which evaluates the degree to which pairs of observations fall on the 45° line through the origin and which provides a measure of both precision and accuracy of an assay was used.[17]

For analysis of the results from samples obtained during the cross-sectional surveys, geometric mean antibody concentrations (GMCs) were calculated and the percentage of samples reaching the putative protective threshold of  $\geq 2$  µg/mL,[18] together with 95% confidence intervals, was determined. Results by country were analysed graphically using reverse cumulative distribution plots. GMCs were compared by urban /rural residence and by sex in each country using a t-test. Risk factors for seropositivity (i.e. antibody concentration  $\geq 2$  µg/mL) were investigated using logistic regression. A multivariable logistic regression model was developed as follows: all variables with a p-value  $< 0.1$  in univariable analyses were included initially, then any variable with p-value  $< 0.05$  in the multivariable model was retained, with excluded variables re-entered one by one. If any of the re-entered variables had a p-value  $< 0.05$ , they were retained in the final model. Because the survey was designed with the household as the primary sampling unit, and to account for potential household clustering, we used the survey commands in Stata (StataCorp, Texas).

Seroprevalence was stratified into yearly age groups and then analysed using a reverse catalytic modelling approach under a binomial sampling assumption, as described elsewhere.[13] Two key parameters were estimated using this approach (1) the seroconversion rate (SCR), i.e. the annual rate at which individuals change from seronegative to seropositive, also known as the force of infection ( $\lambda$ ) and (2) the seroreversion rate, the annual rate at which seropositive individuals revert to a seronegative state (SRR or  $r$ ). The catalytic model was fitted using a maximum likelihood approach. Analyses were repeated, excluding individuals who reported recent vaccination to estimate ‘natural’ immunity.

All analyses were performed using Stata v12.0.

### Ethics

The purpose and methods of the study were explained to community leaders at community meetings and through the media. Written, informed consent for obtaining a pharyngeal swab and a blood sample was obtained from adults and for the children under their care. Written informed assent was also obtained from participants aged 12 years or more. Oral assent was obtained from younger children. Consent and assent forms were translated into the relevant local language.

The study protocols, consent and assent forms were approved by the LSHTM Ethics Committee and by the ethics committees of each of the African partner institutions with the exception of Chad, which does not have a formal ethical committee, and where approval for the activities of the consortium was granted by a committee set up to oversee MenAfriCar studies by the Ministry of Health.

### Results

#### Cross-validation

Although the cross-validation exercise required several rounds of testing, five centres finally achieved excellent results with Pearson correlation values ( $\rho$ ) between 0.926 and 0.996 (Table 1). Final results obtained in Ghana are shown as an example in S1 Fig. Following the validation exercise, quality control of the results obtained on analysis of the cross-sectional survey samples was ensured by monitoring the key parameters of the standard curve as well as the local positive controls, as shown for Ethiopia in S2A Fig.

#### Meningococcal serogroup A IgG antibodies by country and by sub-group

Sera obtained from 3930 individuals in six African meningitis belt countries were tested. The prevalence of serogroup A specific IgG antibodies for each country is shown as a reverse


**Table 1. Statistical analysis of a comparison of meningococcal serogroup A IgG concentrations obtained at the Vaccine Evaluation Unit (VEU), Public Health England and at five MenAfriCar centres.**

Variable/ Country	Ethiopia	Ghana	Mali 1	Mali 2*	Niger	Senegal
Number of sera tested	50 (29 + 21)	49	50	29	50	60 (39 + 21)
Range of values tested (ug/ml)	0.095–133.38	0.34–133.38	0.43–67.42	0.095–3.42	0.095–79.36	0.28–133.38
Pearson’s correlation coefficient ( $\rho$ )	0.996	0.994	0.859	0.947	0.938	0.996
Lin’s concordance coefficient ( $\rho_c$ )	0.988	0.991	0.858	0.825	0.757	0.966
Slope	1.135	0.930	0.970	1.338	0.577	1.281
Intercept	-0.957	-0.372	0.010	0.159	0.966	-0.703

Footnote

\* Two sets of validation samples were used for Mali because the correlation co-efficient from the first 50 samples did not reach the required threshold of 0.9. The second set of samples was selected from a set tested at the VEU which gave consistent results when tested in Manchester, UK.

doi:10.1371/journal.pone.0147928.t001



**Fig 1. Reverse cumulative distribution curves of meningococcal serogroup A IgG antibodies by country.**

doi:10.1371/journal.pone.0147928.g001

cumulative plot in Fig 1. There were significant differences GMC by country with Ghana having the highest mean GMC and Ethiopia the lowest (Table 2).

The GMC increased with age in all countries with the exception of Senegal where there was a drop in GMC in those over 30 years of age (Table 3). This was less apparent when subjects with a history of recent vaccination were excluded. In four countries, no differences in GMCs by sex were observed (Ethiopia, Niger, Nigeria, Senegal,  $p > 0.2$  in each country); in Ghana and Senegal GMCs were higher in females compared to males ( $p = 0.0002$  and  $p = 0.0203$ , respectively). A comparison was made in each country between GMCs according to whether the

**Table 2. Geometric mean meningococcal serogroup A IgG antibody concentrations and seroprevalence by country.**

A. All individuals			
Country	Number	GMC (95% CI)	% $\geq 2\mu\text{g/ml}$ (95% CI)
Ethiopia	619	1.43 (1.31, 1.57)	33.8% (30.0, 37.5%)
Ghana	765	9.09 (8.29, 9.97)	87.6% (85.2, 89.9%)
Mali	756	2.04 (1.77, 2.34)	52.1% (48.5, 55.7%)
Niger	826	4.98 (4.37, 5.66)	65.6% (62.3, 68.8%)
Nigeria	584	2.24 (1.91, 2.62)	52.1% (48.1, 56.2%)
Senegal	380	5.90 (5.00, 6.96)	81.1% (77.1, 85.0%)
B. Excluding individuals with a history of recent meningococcal vaccination			
Country	Number	GMC (95% CI)	% $\geq 2\mu\text{g/ml}$ (95% CI)
Ethiopia	617	1.43 (1.31, 1.57)	33.7% (30.0, 37.4%)
Ghana	650	8.70 (7.89, 9.59)	87.2% (84.6, 89.8%)
Mali	706	2.10 (1.82, 2.43)	52.7% (49.0, 56.4%)
Niger	757	5.24 (4.59, 5.99)	66.8% (63.5, 70.2%)
Nigeria	559	2.20 (1.88, 2.58)	52.0% (47.9, 56.2%)
Senegal	178	3.96 (3.05, 5.15)	74.7% (68.3, 81.1%)

doi:10.1371/journal.pone.0147928.t002

**Table 3. Geometric mean meningococcal serogroup A IgG antibody concentrations (95% CI) by age and country.**

A. All individuals				
Country	Age group (years)			
	< 5	5–14	15–29	> = 30
Ethiopia	0.62 (0.53, 0.72)	0.89 (0.78, 1.03)	1.86 (1.59, 2.17)	2.68 (2.30, 3.14)
Ghana	2.48 (1.92, 3.19)	6.25 (5.06, 7.72)	11.59 (9.97, 13.47)	12.06 (10.55, 13.80)
Mali	0.28 (0.13, 0.34)	1.89 (1.46, 2.46)	5.75 (4.67, 7.08)	5.86 (4.84, 7.10)
Niger	0.89 (0.72, 1.10)	3.39 (2.67, 4.31)	10.99 (8.90, 13.57)	14.36 (11.89, 17.35)
Nigeria	0.39 (0.30, 0.51)	1.00 (0.78, 1.29)	5.07 (3.95, 6.50)	8.88 (7.25, 10.88)
Senegal	1.84 (1.18, 2.87)	7.40 (5.46, 10.02)	11.95 (9.01, 15.87)	4.26, 7.07)
B. Excluding individuals with a history of recent meningococcal vaccination				
Country	Age group (years)			
	< 5	5–14	15–29	> = 30
Ethiopia	0.62 (0.53, 0.72)	0.89 (0.78, 1.03)	1.86 (1.59, 2.17)	2.68 (2.30, 3.14)
Ghana	2.41 (1.81, 3.22)	5.79 (4.64, 7.23)	11.36 (9.65, 13.38)	11.29 (9.81, 13.00)
Mali	0.26 (0.22, 0.32)	1.92 (1.46, 2.53)	5.67 (4.59, 7.00)	6.04 (4.98, 7.33)
Niger	0.94 (0.75, 1.19)	3.50 (2.72, 4.50)	10.89 (8.73, 13.57)	14.26 (11.77, 17.28)
Nigeria	0.39 (0.30, 0.51)	0.89 (0.70, 1.14)	4.88 (3.81, 6.26)	8.59 (7.02, 10.52)
Senegal	0.94 (0.41, 2.19)	3.68 (1.41, 9.60)	6.21 (3.28, 11.78)	5.10 (3.86, 6.74)

doi:10.1371/journal.pone.0147928.t003

study site was urban or rural and by the sex of the participant. In five countries, NmA-specific IgG GMCs were higher in urban than in rural areas ( $p < 0.0001$  for Ethiopia, Ghana, Mali and Niger,  $p = 0.0012$  for Senegal); in Nigeria, GMC was higher in the rural study site ( $p = 0.0123$ ).

Factors associated with NmA-specific IgG concentrations above the putative protective threshold of 2 µg/ml are shown in Table 4. The multivariable model included the following factors shown to be associated with a higher odds of seropositivity in addition to observed country-level differences: urban location, increasing age and reported receipt of a meningitis vaccine which is likely to have contained the serogroup A polysaccharide. Current carriage of meningococci or other *Neisseria* species were not associated with seropositivity in the multivariable model.

The influence of age on the putative protective threshold of 2 µg/mL is further shown by country in Figs 2 and 3.

### Force of infection

The force of infection was calculated based on the age prevalence of seropositivity as described above. Unexpectedly, this was highest in Senegal, even when subjects with a recent history of vaccination were excluded, followed by Ghana and Niger; Ethiopia had the slowest seroconversion rate (Table 5).

### Discussion

Although a number of studies of meningococcal serology have been performed previously in countries of the African meningitis belt, most of these have been undertaken in the context of evaluation of the response to vaccination [7–10] or in disease survivors [19]. Comparisons between the results obtained in individual studies can be difficult when standardised methods are not used. Therefore, for this comparative study, substantial efforts were made to standardise the ELISA technique used at each of the collaborating centres with support from the Vaccine Evaluation Unit at PHE, Manchester, UK. This proved more challenging than anticipated

**Table 4. Logistic regression analysis of factors associated with a meningococcal serogroup A IgG concentration  $\geq 2\mu\text{g/ml}$ , a putative correlate of protection.**


Variable	N	Crude Odds Ratio (95% CI)	Adjusted Odds Ratio (95% CI)
<i>Country</i>			
Ethiopia	619	0.27 (0.21, 0.33)	0.15 (0.11, 0.20)
Ghana	765	3.72 (2.83, 4.91)	2.79 (2.03, 3.85)
Mali	756	0.57 (0.46, 0.71)	0.49 (0.38, 0.64)
Niger (baseline)	826	1.0	1.0
Nigeria	583	0.57 (0.46, 0.71)	0.44 (0.34, 0.58)
Senegal	380	2.24 (1.65, 3.00)	2.02 (1.31, 3.12)
<i>Age group</i>			
0 to 4 years	746	0.09 (0.07, 0.11)	0.06 (0.05, 0.08)
5 to 14 years	957	0.32 (0.27, 0.39)	0.26 (0.21, 0.32)
15 to 29 years (baseline)	1023	1.0	1.0
30+ years	1203	1.59 (1.29, 1.96)	1.68 (1.33, 2.12)
<i>Sex</i>			
Female	2241	1.0	Not included
Male	1668	0.82 (0.72, 0.93)	
Not known	20		
Urban	1871	1.0	1.0
Rural	2058	0.64 (0.56, 0.73)	0.54 (0.45, 0.64)
<i>Recently vaccinated with a meningitis vaccine</i>			
No	3459	1.0	1.0
Yes	463	2.09 (1.65, 2.66)	1.71 (1.23, 2.38)
<i>Meningococcal carrier</i>			
No	3771	1.0	Not included
Yes	158	0.94 (0.67, 1.31)	
<i>Carrier of other <i>Neisseria</i> species</i>			
No	3706	1.0	Not included
Yes	224	0.40 (0.30, 0.53)	
<i>Crowded living conditions*</i>			
No	1438	1.0	Not included
Yes	2484	0.70 (0.60, 0.81)	

\* crowding was defined as  $> = 2$  people per room as used previously[27]

doi:10.1371/journal.pone.0147928.t004

and satisfactory results were not obtained at one out of six centres prior to the end of the project. Problems encountered included the short shelf-life of some of the reagents required, difficulties in clearing reagents through customs and difficulties in shipping frozen samples to the UK for validation. However, despite these challenges, excellent results were eventually obtained at five centres using Pearson coefficient of correlation ( $r > 0.92$ ) and satisfactory ones using Lin's concordance coefficient of correlation, allowing adjustment of some issues which were not identified using the correlation coefficient alone.

Overall, serogroup A meningococcal IgG antibody concentrations were high in the African populations investigated, with the highest GMCs being obtained in countries in the centre of the meningitis belt. This has been noted previously.[10,11] This was the case despite the fact that there was little circulation of the serogroup A meningococcus in the African meningitis belt at the time of the study.[20] These high antibody concentrations probably reflect prior exposure to the serogroup A meningococcus and other cross-reactive bacteria such as *Bacillus*




**Fig 2. Statistical analysis of seropositivity data for all individuals.** Age-adjusted seroprevalence (blue solid lines) using appropriate reversible catalytic models. The observed seroprevalences (red-filled triangles) were pooled according to the 10%-centiles of the underlying age distribution.

doi:10.1371/journal.pone.0147928.g002

*pumilis* [5,21] and *Escherichia coli* capsule types K51 or K93.[22] Similarly high serogroup A meningococcal IgG antibody concentrations have been found in industrialised countries where there is no group A disease and no evidence of carriage. For example, in England and Wales a GMC of 4.75 µg/mL was found in the general population with 87% of individuals with antibody levels  $\geq 2$  µg/mL,[23] an observation which supports the role for cross-reactive bacteria in inducing antibodies against the serogroup A meningococcus.

None of the populations in the MenAfriCar cross-sectional surveys had been vaccinated previously with MenAfriVac™ but meningococcal vaccines containing A polysaccharide have been used quite extensively in some of the study countries in the past. Subjects were asked whether they had received a meningitis vaccine in the previous six months and if this was the case, they are likely to have been vaccinated with a vaccine containing serogroup A polysaccharide. Analyses of antibody distribution by country and by age were undertaken excluding these subjects but this had little impact on the pattern of results. However, earlier vaccination campaigns may have resulted in some persistent antibody. Prior vaccination may be the reason for



**Fig 3. Statistical analysis of seropositivity data excluding vaccinated individuals.** Age-adjusted seroprevalence (blue solid lines) using appropriate reversible catalytic models. The observed seroprevalences (red-filled triangles) were pooled according to the 10%-centiles of the underlying age distribution.

doi:10.1371/journal.pone.0147928.g003

the rapid seroconversion rate in the Senegalese population where a mass vaccination campaign with an A + C polysaccharide vaccine had been undertaken in 2010, about 12 months prior to the study and also for the reason why the GMC was lower in subjects over 30 years of age in Senegal than in younger subjects who would not have been targeted in the immunisation campaign. Thus, the antibody distribution seen in each country may reflect a combination of responses induced by both natural exposure and vaccination.

An Ig meningococcal serogroup A polysaccharide antibody concentration of  $> 2 \mu\text{g/mL}$  as measured by radioimmunoassay has been suggested as a correlate of protection against serogroup A meningococcal disease based on the results of a trial of a meningococcal polysaccharide vaccine conducted in Finland in the 1970s.[18] However, it seems unlikely that this is the case in countries of the meningitis belt where a high proportion of the population has antibody concentrations above this value and yet, until the introduction of MenAfriVac™, the region remained peculiarly susceptible to large serogroup A epidemics. It is likely that much of the antibody detected by ELISA is non-functional, perhaps because it is induced by cross-reacting

**Table 5. Estimates of the annual force of infection ( $\lambda$ ) and seroreversion ( $r$ ) by country.**

A. All individuals		
Country	$\lambda$ (95% CI)	$r$ (95% CI)
Ethiopia	0.028 (0.022, 0.036)	0.015 (0.007, 0.033)
Ghana	0.240 (0.192, 0.299)	0.015 (0.009, 0.024)
Mali	0.077 (0.064, 0.091)	0.017 (0.011, 0.027)
Niger	0.110(0.093, 0.130)	0.009 (0.005, 0.018)
Nigeria	0.052 (0.044, 0.062)	0.005 (0.000, 0.485)
Senegal	0.434 (0.282, 0.670)	.033, 0.124)
B. Excluding individuals with a history of recent meningococcal vaccination		
Country	$\lambda$ (95% CI)	$r$ (95% CI)
Ethiopia	0.028 (0.022, 0.036)	0.015 (0.007, 0.333)
Ghana	0.233 (0.183, 0.295)	0.015 (0.008, 0.025)
Mali	0.075 (0.063, 0.090)	0.016 (0.010, 0.027)
Niger	0.114 (0.095, 0.137)	0.010 (0.005, 0.020)
Nigeria	0.050 (0.042, 0.060)	0.0002 (0, 1)
Senegal	0.457 (0.160, 1.30)	0.109 (0.031, 0.380)

doi:10.1371/journal.pone.0147928.t005

bacteria. Bactericidal antibodies are the accepted correlate of protection for meningococcal disease [24] and measurement of serogroup A serum bactericidal antibodies may give a better reflection of the background level of immunity of a community. However, these are technically more difficult to perform reliably than the ELISA, and were only undertaken at two of the centres that participated in this study before and after the introduction of MenAfriVac™. These results, together with studies of the correlates of protection against invasive meningococcal disease and meningococcal pharyngeal carriage will be reported subsequently.

In this study, we have shown that serology can be used to show differences in exposure to meningococcal infection between countries and age groups and we have shown how age dependent variations in seropositivity can be used to measure the force of infection. The force of infection was generally highest in countries in the centre of the meningitis belt, with the exception of Senegal where the situation may have been confounded by vaccination, and lowest in Ethiopia on the margin of the belt. Measurement of seroconversion by age has proved to be a valuable approach to study of the epidemiology of malaria and of the impact of control interventions on this infection [25,26] and, as shown here, is valuable for the study of other infectious diseases. Further study of antibody kinetics in the African meningitis belt is important for understanding the epidemiology of meningococcal infection and monitoring control measures including widespread deployment of conjugate vaccines.

## Supporting Information

**S1 Fig.**  
(DOCX)

**S2 Fig.**  
(DOCX)

## Acknowledgments

Firstly, we thank the tens of thousands of individuals who participated in the surveys reported in this paper. The work described here also relied upon many staff, including fieldworkers and laboratory technicians whom we thank for their contribution. We acknowledge the Directors

of the African research centres for their support and the following individuals who provided clinical monitoring: Richard Nare (Chad), Frank Baiden (Ghana), Workeabeba Taye (Ethiopia), Haoua Amadou (Niger and Mali), and Pierre Ndiaye (Senegal). Lesley Mabey and Daniel Holme (Vaccine Evaluation Unit, Manchester) supported transfer of the ELISA assay to Africa and its evaluation. The guidance provided by the MenAfriCar Advisory Committee (Fred Binka, Mamadou Djingarey, Robert Heyderman, Marie-Paule Kiene, Marie-Pierre Preziosi, David Stephens and Marcel Tanner [chairman]) has been much appreciated.

We also thank the following individuals who contributed to the establishment of the MenAfriCar Consortium and to its activities in various ways—Eric Bertherat and William Perea (WHO, Geneva, Switzerland), Dominique Caugant (NIPH, Oslo, Norway), Mamadou Djingarey (WHO, Ouagadougou, Burkina Faso), Rana Hajjeh and Nancy Messonnier (CDC, Atlanta, Georgia), Marc LaForce (PATH, Seattle, USA), Judith Mueller (EHESP, Rennes, France), Pierre Nicolas (IMTSSA, Marseilles, France), Gerd Pluschke (STPI, Basle, Switzerland) and Muhamed-KheirTaha (Institut Pasteur, Paris, France). The work of the consortium across Africa would not have been possible without the strong logistic support provided by members of the MenAfriCar secretariat in London—Amit Bhasin, Elizabeth Huntley, Karen Williams, Karen Slater and Lyanne Wylde. Studies conducted in each country received full support from the national health and local authorities and this is gratefully acknowledged.

## Author Contributions

Conceived and designed the experiments: OM CT HF AA AH SS J-MC J-FJ CS DMD BO DC MH-K MN RB JMS BG. Performed the experiments: WM TMD BY IO SLQ MC KD AT RMB OD AEM SA SD EHB MD AD. Analyzed the data: OM CT AW JMS BG. Wrote the paper: OM CT HF AA WM TMD BY IO AH SLQ SS MC KD AT JMC RMB OD AEM JFJ CS SA SD EHB MD AD DMD BO DC MHK MN AW RB JMS BG.

## References

1. Greenwood B. 1999. Manson Lecture. Meningococcal meningitis in Africa. *Trans R Soc Trop Med Hyg* 93: 341–53. PMID: [10674069](#)
2. Goldschneider I, Gotschlich EC, Artenstein MS. 1969. Human immunity to the meningococcus. II. Development of natural immunity. *J Exp Med* 129: 1327–48. PMID: [4977281](#)
3. Gold R, Goldschneider I, Lepow ML, Draper TF, Randolph M. 1978. Carriage of *Neisseria meningitidis* and *Neisseria lactamica* in infants and children. *J Infect Dis* 137: 112–21. PMID: [415097](#)
4. Robbins JB, Schneerson R, Glode MP, Vann W, Schiffer MS, Liu TY, et al. 1975. Cross-reactive antigens and immunity to diseases caused by encapsulated bacteria. *J Allergy Clin Immunol* 56: 141–51. PMID: [239050](#)
5. Vann WF, Liu TY, Robbins JB. 1976. *Bacillus pumilus* polysaccharide cross-reactive with meningococcal group A polysaccharide. *Infect Immun* 13: 9.
6. Griffiss JM. 1975. Bactericidal activity of meningococcal antisera. Blocking by IgA of lytic antibody in human convalescent sera. *Journal of immunology* 114: 1779–84.
7. Greenwood BM, Bradley AK, Blakebrough IS, Whittle HC, Marshall TF, Gilles HM. 1980. The immune response to a meningococcal polysaccharide vaccine in an African village. *Trans R Soc Trop Med Hyg* 74:340–6. PMID: [6776665](#)
8. Ceesay SJ, Allen SJ, Menon A, Todd JE, Cham K, Carlone GM, et al. 1993. Decline in meningococcal antibody levels in African children 5 years after vaccination and the lack of an effect of booster immunization. *J Infect Dis* 167: 1212–6. PMID: [8486957](#)
9. MacLennan J, Obaro S, Deeks J, Williams D, Pais L, Carlone G, et al. 1999. Immune response to revaccination with meningococcal A and C polysaccharides in Gambian children following repeated immunisation during early childhood. *Vaccine* 17: 3086–93. PMID: [10462244](#)
10. Sow SO, Okoko BJ, Diallo A, Viviani S, Borrow R, Carlone G, et al. 2011. I. Immunogenicity and safety of a meningococcal A conjugate vaccine in Africans. *N Engl J Med* 364: 2293–304. doi: [10.1056/NEJMoa1003812](#) PMID: [21675889](#)

11. Trotter CL, Yaro S, Njanpop-Lafourcade BM, Drabo A, Kroman SS, et al. 2013. Seroprevalence of bactericidal, specific IgG antibodies and Incidence of meningitis due to Group A *Neisseria meningitidis* by age in Burkina Faso 2008. *PLoS One* 8: e55486. doi: [10.1371/journal.pone.0055486](https://doi.org/10.1371/journal.pone.0055486) PMID: [23457471](https://pubmed.ncbi.nlm.nih.gov/23457471/)
12. Mueller JE, Yaro S, Njanpop-Lafourcade BM, Drabo A, Idohou RS, et al. 2011. Study of a localized meningococcal meningitis epidemic in Burkina Faso: incidence, carriage, and immunity. *J Infect Dis* 204: 1787–95. doi: [10.1093/infdis/jir623](https://doi.org/10.1093/infdis/jir623) PMID: [21998478](https://pubmed.ncbi.nlm.nih.gov/21998478/)
13. Drakeley CJ, Corran PH, Coleman PG, Tongren JE, McDonald SL, Carneiro I, et al. 2005. Estimating medium- and long-term trends in malaria transmission by using serological markers of malaria exposure. *Proc Natl Acad Sci USA* 102: 5108–13. PMID: [15792998](https://pubmed.ncbi.nlm.nih.gov/15792998/)
14. MenAfriCar Consortium. 2013. Meningococcal carriage in the meningitis belt. *Trop Med Int Health* 19:968–78.
15. Collard JM, Issaka B, Zaneidou M, Hugonnet S, Nicolas P, Taha MK, et al. 2013. Epidemiology changes in meningococcal meningitis strains in Niger from 2008 to 2011; attempt for an explanation and perspectives for vaccination strategy. *BMC Infect Dis* 13:576. doi: [10.1186/1471-2334-13-576](https://doi.org/10.1186/1471-2334-13-576) PMID: [24313998](https://pubmed.ncbi.nlm.nih.gov/24313998/)
16. Carlone GM, Frasch CE, Siber GR, Quataert S, Gheesling LL, Turner SH, et al. 1992. Multicenter comparison of levels of antibody to *Neisseria meningitidis* group A capsular polysaccharide measured by using an enzyme-linked immunosorbent assay. *J Clin Microbiol* 30: 154–9 PMID: [1734048](https://pubmed.ncbi.nlm.nih.gov/1734048/)
17. Lin LI. 1989. A concordance correlation coefficient to evaluate reproducibility. *Biometrics* 45: 255–68. PMID: [2720055](https://pubmed.ncbi.nlm.nih.gov/2720055/)
18. Peltola H, Makela H, Kayhty H, Jousimies H, Herva E, Hällström K, et al. 1977. Clinical efficacy of meningococcus group A capsular polysaccharide vaccine in children three months to five years of age. *N Engl J Med* 297: 686–91. PMID: [408682](https://pubmed.ncbi.nlm.nih.gov/408682/)
19. Norheim G, Aseffa A, Yassin MA, Mengistu G, Kassu A, Fikremariam D, et al. 2008. Specificity of sub-capsular antibody responses in Ethiopian patients following disease caused by serogroup A meningococci. *Clin Vaccine Immunol* 15: 863–71. doi: [10.1128/CVI.00252-07](https://doi.org/10.1128/CVI.00252-07) PMID: [18337382](https://pubmed.ncbi.nlm.nih.gov/18337382/)
20. MenAfriCar Consortium. 2015. The diversity of meningococcal carriage across the African meningitis belt and the impact of vaccination with a Group A meningococcal conjugate vaccine. *J Infect Dis pii: jiv211*. [Epub ahead of print].
21. Robbins JB, Myerowitz L, Whisnant JK, Argaman M, Schneerson R, Handzel ZT, et al. 1972. Enteric bacteria cross-reactive with *Neisseria meningitidis* groups A and C and *Diplococcus pneumoniae* types I and 3. *Infect Immun* 6: 651–6. PMID: [4118016](https://pubmed.ncbi.nlm.nih.gov/4118016/)
22. Guirguis N, Schneerson R, Bax A, Egan W, Robbins JB, Shiloach J, et al. 1985. *Escherichia coli* K51 and K93 capsular polysaccharides are cross reactive with the group A capsular polysaccharide of *Neisseria meningitidis*. Immunochemical, biological, and epidemiological studies. *J Exp Med* 162: 1837–51. PMID: [3934317](https://pubmed.ncbi.nlm.nih.gov/3934317/)
23. Trotter CL, Borrow R, Findlow J, Holland A, Frankland S, Andrews NJ, et al. 2008. Seroprevalence of antibodies against serogroup C meningococci in England in the post vaccination era. *Clin Vaccine Immunol* 15: 1694–8. doi: [10.1128/CVI.00279-08](https://doi.org/10.1128/CVI.00279-08) PMID: [18827191](https://pubmed.ncbi.nlm.nih.gov/18827191/)
24. Frasch CE, Borrow R, Donnelly J. 2009. Bactericidal antibody is the immunologic surrogate of protection against meningococcal disease. *Vaccine* 27 Suppl 2:B112–6. doi: [10.1016/j.vaccine.2009.04.065](https://doi.org/10.1016/j.vaccine.2009.04.065) PMID: [19464093](https://pubmed.ncbi.nlm.nih.gov/19464093/)
25. Wong J, Hamel MJ, Drakeley CJ, Kariuki S, Shi YP, Lal AA, et al. 2014. Serological markers for monitoring historical changes in malaria transmission intensity in a highly endemic region of Western Kenya, 1994–2009. *Malar J* 13:451. doi: [10.1186/1475-2875-13-451](https://doi.org/10.1186/1475-2875-13-451) PMID: [25416454](https://pubmed.ncbi.nlm.nih.gov/25416454/)
26. Cook J, Kleinschmidt I, Schwabe C, Nseng G, Bousema T, Corran PH, et al. 2011. Serological markers suggest heterogeneity of effectiveness of malaria control interventions on Bioko Island, equatorial Guinea. *PLoS One* 6: e25137. doi: [10.1371/journal.pone.0025137](https://doi.org/10.1371/journal.pone.0025137) PMID: [21980386](https://pubmed.ncbi.nlm.nih.gov/21980386/)
27. Ferraro CF, Trotter CL, Nascimento MC, Jusot JF, Omotara BA, Hodgson A, et al. 2014. Household crowding, social mixing patterns and respiratory symptoms in seven countries of the African meningitis belt. *PLoS One* 9: e101129. doi: [10.1371/journal.pone.0101129](https://doi.org/10.1371/journal.pone.0101129) PMID: [24988195](https://pubmed.ncbi.nlm.nih.gov/24988195/)