



**HAL**  
open science

## Editorial: Exploring the unique biodiversity of the Western Pacific to identify novel anti-infectious and anti-inflammatory compounds of natural origin

Philippe Georgel, Nicolas Lebouvier, Mariko Matsui

### ► To cite this version:

Philippe Georgel, Nicolas Lebouvier, Mariko Matsui. Editorial: Exploring the unique biodiversity of the Western Pacific to identify novel anti-infectious and anti-inflammatory compounds of natural origin. *Frontiers in Pharmacology*, 2023, 14, pp.1154627. 10.3389/fphar.2023.1154627. pasteur-03984042

**HAL Id: pasteur-03984042**

**<https://riip.hal.science/pasteur-03984042v1>**

Submitted on 12 Feb 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## OPEN ACCESS

## EDITED BY

Javier Echeverria,  
University of Santiago, Chile

## REVIEWED BY

Adolfo Andrade-Cetto,  
National Autonomous University of  
Mexico, Mexico

## \*CORRESPONDENCE

Mariko Matsui,  
✉ mmatsui@pasteur.nc

## SPECIALTY SECTION

This article was submitted to  
Ethnopharmacology, a section of the  
journal Frontiers in Pharmacology

RECEIVED 30 January 2023

ACCEPTED 10 February 2023

PUBLISHED 20 February 2023

## CITATION

Georgel P, Lebouvier N and Matsui M  
(2023), Editorial: Exploring the unique  
biodiversity of the Western Pacific to  
identify novel anti-infectious and anti-  
inflammatory compounds of  
natural origin.

*Front. Pharmacol.* 14:1154627.

doi: 10.3389/fphar.2023.1154627

## COPYRIGHT

© 2023 Georgel, Lebouvier and Matsui.  
This is an open-access article distributed  
under the terms of the [Creative  
Commons Attribution License \(CC BY\)](#).  
The use, distribution or reproduction in  
other forums is permitted, provided the  
original author(s) and the copyright  
owner(s) are credited and that the original  
publication in this journal is cited, in  
accordance with accepted academic  
practice. No use, distribution or  
reproduction is permitted which does not  
comply with these terms.

# Editorial: Exploring the unique biodiversity of the Western Pacific to identify novel anti-infectious and anti-inflammatory compounds of natural origin

Philippe Georgel<sup>1</sup>, Nicolas Lebouvier<sup>2</sup> and Mariko Matsui<sup>2,3\*</sup>

<sup>1</sup>Laboratoire d'ImmunoRhumatologie Moléculaire, Faculté de Médecine, Institut National de la Santé et de la Recherche Médicale (INSERM) UMR\_S 1109, Institut thématique interdisciplinaire (ITI) de Médecine de Précision de Strasbourg, Transplantex NG, Fédération Hospitalo-Universitaire OMICARE, Fédération de Médecine Translationnelle de Strasbourg (FMTS), Université de Strasbourg, Strasbourg, France, <sup>2</sup>Institut des Sciences Exactes et Appliquées (ISEA) EA7484, Université de la Nouvelle Calédonie, Noumea, New Caledonia, <sup>3</sup>Group Bioactivities of Natural compounds and derivatives (BIONA), Institut Pasteur of New Caledonia, Member of the Pasteur Network, Noumea, New Caledonia

## KEYWORDS

Western Pacific, biodiversity, bioactivity, phytochemistry, traditional medicine, natural products

## Editorial on the Research Topic

Exploring the unique biodiversity of the Western Pacific to identify novel anti-infectious and anti-inflammatory compounds of natural origin

Inflammation and many infectious diseases, such as sepsis, are still ongoing major health concerns. Although considerable therapeutic advances were made over the last decades, resulting in increased life expectancy, several hurdles remain to be addressed. Namely, antimicrobial resistance (AMR)—that led to more than 1.2 million deaths in 2019 (Murray et al., 2022)—or the adverse effects of several anti-inflammatory compounds are of utmost importance. Therefore, continuous efforts aiming at the discovery and development of novel molecules to deal with these challenges are required. For the past couple of years, research on natural products (NPs) used in traditional, herbal medicine, has been of high interest (Thomford et al., 2018) because such ancestral knowledge may reveal less toxic active compounds with fewer side effects.

The Western Pacific Region (WPR) comprises 37 countries and territories from China to New Zealand. Including 9 of the 36 established biodiversity hotspots (Kobayashi et al., 2019), the WPR therefore represents territories where medicinal plants belong to ecosystems threatened by human activities. Moreover, state members of the WPR show a strong acknowledgment of Traditional and Complementary Medicine (T&CM), with 93% of them reporting use of T&CM by their populations (WHO, 2019). Interaction between T&CM and conventional medicine, especially herb-drug interactions (HDI), can possibly cause ineffective treatment or adverse effects (Rosenkranz et al., 2012). Precise mechanisms of action and description of bioactive molecules participate in the prevention of HDI. In this Research Topic, bioactivities and phytochemistry of several traditional plants used in T&CM from the WPR countries are introduced, focusing on their chemical components and the related anti-infectious, immunoregulatory and healing properties.

Native from South-East Asia, *Calotropis gigantea* is an Apocyanaceae found in several Pacific Islands up to Australia and Central and South America. The whole plant is traditionally used to cure common skin diseases, including burns and small wounds. Roots and bark extracts are also used against leprosy, scabies and other infectious diseases. In their original research, Alafnan et al. studied the chemical composition and the biological properties of the *C. gigantea* leaves extracts. Phytochemical analyses showed that leave extracts contain 17 different compounds, mostly sesquiterpene, alkaloids and flavonoids. Ethanol extracts of *C. gigantea* leaves exhibited considerable antioxidant activities. This paper also reports that plant extracts inhibited acetylcholinesterase (AChE) and tyrosinase enzymatic activities. *in vivo* upon topical application, these extracts improved wound healing following surgical- or burn-induced skin damage in Wistar rats.

*Helminthostachys zeylanica* is a fern-like plant belonging to the Ophioglossaceae family and native from the South-East Asia and Australia. This plant is traditionally used to treat diabetes, inflammatory and hepatic disorders and various infectious diseases. Shah et al. revealed that bacterial neuraminidase (BNA), which is an important player in several microbial infections and biofilm formation, is inhibited by the six known ugonin metabolites isolated from the rhizome of *H. zeylanica*. Interestingly, inhibitory activities of the compounds were higher than the reference (quercetin) and structurally related compounds (luteolin and eriodictiol). Biofilm formation by *E. coli* was also inhibited by the ugonin J in a dose-dependent manner, highlighting a potential antibacterial interest of the plant and these molecules.

Widespread in the South-West Pacific Islands, *Coleus forsteri* is an herbaceous Lamiaceae traditionally used to treat flu-like symptoms; it is also utilized as analgesic following ecchymosis. Only few studies characterized the chemical composition and biological activities of the plant. In this Research Topic, Nicolas et al. showed evidence of anti-inflammatory activities of *C. forsteri* whole extracts on human macrophages. Ethanol and cyclohexane extracts of *C. forsteri* significantly decreased LPS-induced inflammatory cytokine IL-6 and TNF- $\alpha$  and quinolinic acid production. Seven known abietane diterpenes were characterized from *C. forsteri* cyclohexane extracts, including coleon U, horminone and family-related compounds which potentially contribute to the plant anti-inflammatory activities.

Among traditional Chinese medicine, Danggui Buxue Tang (DBT) is a formulae of two medicinal herbs containing root extracts of *Angelica sinensis* (Apiaceae) and *Astragalus membranaceus* (Fabaceae). DBT is recommended for patient

recovery after a bone fracture, especially in aged women. Calycosin is the major active flavonoid found in *A. membranaceus* roots and in DBT. Using multi-omics approaches, the team of Kwan et al. showed that DBT and calycosin orchestrate osteoblastic differentiation and proliferation by regulating signaling pathways at RNA and protein levels, impacting on energy metabolism. Moreover, they showed that calycosin controls interaction with other DBT components during these processes.

In conclusion, this Research Topic reveals new biological properties and chemical characterization of four traditional plants and T&CM used within the WPR. It also highlights further explorations that are expected in this field to deepen our knowledge of their mechanisms of action.

## Author contributions

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

## Acknowledgments

We acknowledge the French Ministry of Higher Education, Research, and Innovation (MESRI), the Institut Pasteur of New Caledonia (IPNC), the Pasteur Network, and the Government of New Caledonia for funding MM position. We would like to thank all the authors, reviewers and frontiers editorial team for their valuable input and contributions.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Kobayashi, Y., Okada, K.-i., and Mori, A. S. (2019). Reconsidering biodiversity hotspots based on the rate of historical land-use change. *Biol. Conserv.* 233, 268–275. doi:10.1016/j.biocon.2019.02.032
- Murray, C. J. L., Ikuta, K. S., Sharara, F., Swetschinski, L., Robles Aguilar, G., Gray, A., et al. (2022). Global burden of bacterial antimicrobial resistance in 2019: A systematic analysis. *Lancet* 399 (10325), 629–655. doi:10.1016/S0140-6736(21)02724-0
- Rosenkranz, B., Fasinu, P., and Bouic, P. (2012). An overview of the evidence and mechanisms of herb-drug interactions. *Front. Pharmacol.* 3, 69. doi:10.3389/fphar.2012.00069
- Thomford, N. E., Senthelane, D. A., Rowe, A., Munro, D., Seele, P., Maroyi, A., et al. (2018). Natural products for drug discovery in the 21st century: Innovations for novel drug discovery. *Int. J. Mol. Sci.* 19 (6), 1578. doi:10.3390/ijms19061578
- WHO (2019). *WHO global report on traditional and complementary medicine*. Available at: <https://www.who.int/publications/i/item/978924151536>.